


INDIGENOUS HIGHER PLANTS (LYCOPODS, QUILLWORTS, FERNS, GYMNO-SPERMS, FLOWERING PLANTS) IN UPPER WAIRAU VALLEY (WAIRAU-RAINBOW FORKS<sup>1</sup> TO ISLAND PASS<sup>2</sup>), 825-2200 M / 2700-7225 FT

<sup>1</sup>S40 26-38-  
<sup>2</sup>S40 17-19-

A.P. Druce, Botany Division, D.S.I.R., 1973  
(Revised May 1974, July 1975, Aug. 1976, May 1977, June 1979,  
June 1980, July 1985, Sept. 1985, Aug. 1986, Aug. 1987, Nov. 1990, June 1991,  
March 1992, May 1993)  
(Based on visits 1972, 1973, 1976, 1980, 1987, 1988, 1991)



(unc) - uncommon or local

**GYMNOSPERM TREES AND SHRUBS**

- Halocarpus bidwillii
- Phyllocladus alpinus s.s.
- Podocarpus nivalis
- P. hallii X P. nivalis (unc)

- C. depressa (unc)
- C. linariifolia (unc)
- C. microcarpa (unc)
- C. niphophila
- C. parpusilla ssp. parpusilla
- C. petriei
- C. propinqua ssp. propinqua

**DICOT TREES**

- Fuchsia excorticata (unc)
- Griselinia littoralis (unc)
- Hoheria lyallii var. lyallii (unc)
- Nothofagus solandri var. cliffortioides
- Pittosporum patulum (unc)
- P. tenuifolium ssp. tenuifolium (unc)
- Pseudopanax sp. (unnamed; included in P. colensoi, as var. ternatum, and as var. fiordensis, by Wardle 1968) (unc)

- C. pseudocuneata ssp. (lvs narrower than in ssp. pseudocuneata) (unc)
- C. rhamnoides (incl. C. polymorpha) (unc)
- C. sp. (i) (unnamed; included in C. parviflora by Oliver, 1935, and others) (If undersurface not minutely hairy as is usual in C. parviflora; drupes usual unpigmented, occ. purplish red) (unc)

**DICOT SHRUBS**

- Aristotelia fruticosa var.
- Brachyglottis bidwillii var. viridis (unc)
- B. cassinioides
- B. grayi var. (B. laxifolius) (unc)
- B. montoi var. montoi (incl. B. m. var. elongatus) (unc)
- Cassinia vauvilliersii
- Coprosma atropurpurea
- C. cheesemanii s.s. (unc)
- C. ciliata ssp. (unc)

- C. sp. (ii) (unnamed; aff. C. pseudocuneata) (low-growing shrubs with small, thick, arcuate lvs)
- Cornaria angustissima (unc)
- C. plumosa (unc)
- C. sarmentosa
- C. angustissima X C. sarmentosa (unc)
- Corokia cotoneaster var. cotoneaster (unc)
- Cyathodes sp. (C. juniperina agg.) (lvs us. < 1 cm X < 1 mm) (unc)
- Discaria toumatou
- Dracophyllum prunum

- D. uniflorum
- D. longifolium X D. uniflorum (unc)
- D. pronom X D. uniflorum
- Exocarpus bidwillii (unc)
- Gaultheria antipoda (unc)
- G. crassa
- G. sp. (unnamed; included in G. depressa, as var. novae-zelandiae, by Franklin 1962)
- G. nubicola
- G. macrostigma (unc)
- G. crassa X G. sp. (unnamed) (unc)
- G. c. X G. macrostigma (unc)
- G. sp. (unnamed) X G. macrostigma (unc)
- Hebe anomala
- H. armstrongii (?) (see Moore 1961)
- H. canterburyensis
- H. cheesemanii
- H. cochayneana
- H. cupressoides (Cheeseman 1882 and 1925, as Veronica cupressoides)
- H. decumbens
- H. epactidea var. epactidea
- H. haastii var. (H. ramosissima) (unc)
- H. lycopodioides
- H. mactantha var. bradyphylla
- H. pauciramosa
- H. pimeleoides s.s. (incl. H. p. var. minor)
- H. pinguifolia

- H. rakaiensis (?)
- H. rupicola (Cheeseman 1925, as Veronica rupicola)
- H. salicifolia var. salicifolia (unc)
- H. salicornioides (Cheeseman 1925, as Veronica salicornioides)
- H. subalpina
- H. tumida
- H. venustula (incl. H. brachysiphon)
- H. vernicosa
- H. sp. (i) (unnamed; included in H. raoulii; as var. pentasepala, by Moore 1961)
- H. sp. (ii) (unnamed; aff. H. rigidula) (lvs glaucous beneath, conspicuously boat-shaped, widest above mid-point)
- Helictysum depressum
- H. parvifolium
- H. sp. (unnamed; aff. H. coralloides) (branchlets smaller in diam. than in H. coralloides; fls creamy, not yellow)
- H. depressum X H. parvifolium (unc)
- H. parvifolium X Raoulia glabra (?) (unc)
- Kunzea ericoides agg.
- Leptospermum scoparium
- Leucopogon fraseri s.s.
- L. suaveolens (incl. Cyathodes colensoi)
- L. sp. (unnamed; included in L. fraseri, as var. muscosus, by Simpson 1945)
- Melicytus alpinus s.s.

*M. sp.* (unnamed; aff. *M. alpinus*)

(low-growing shrubs with re- FERNS  
curved, spinous branches, and *Asplenium flabellifolium* agg. (unc)  
narrowly linear-lanceolate (vs) *A. richardii*

*Myrsine divaricata* s.s. (unc)

*M. nummularia*

*Oleatia arborescens* (unc)

*O. arcennifolia* (unc)

*O. cymbifolia*

*O. odorata*

*Pentachondra pumila*

*Peraxilla tetrapetala* (unc)

*Pimelea concinna* (Allan 1961)

*P. orzophila*

*P. sericeovillosa*

*P. traversii*

*Pittosporum anomalum*

*P. divaricatum*

*Traversia baccharoides*

*A. sp.* (unnamed; aff. *A. trichomanes*)

*Blechnum penna-marina*

*B. sp.* (unnamed) (mountain sp.;  
lower pinnae not reduced in length)  
("The Dunedin-Cass Variety" of  
Crookes 1963) (unc)

*Cystopteris tasmanica*

*Grammitis patagonica* (unc)

*G. poeppigiana*

*Histiopteris incisa* (unc)

*Hymenophyllum multifidum*

*H. villosum*

*Hypolepis millefolium*

*H. rufobarbata* (unc)

*Ophioglossum cotiaceum* agg.

*Pilularia novae-zelandiae*

*Polystichum cystostegia*

*P. vestitum*

*Pteridium esculentum* (unc)

## DICOT LIANS AND RELATED

### TRAILING PLANTS

*Clematis forsteri*

*C. marata* (unc)

*Muehlenbeckia australis* (unc)

*M. axillaris*

*Rubus cissoides* var. (leaflets  
broader than in var. *cissoides*) (unc)

### ORCHIDS

*Aporostylis bifolia* (unc)

*Microtus unifolia* (unc)

*Prasophyllum edensei*

*Pterostylis sp.* (*P. tristis* or *P. sp.*  
aff. *P. cyenocephala*)

*Thelymitra formosa* (unc)

*T. hatchii* (unc)

*T. longifolia* (unc)

### LYCOPODS

*Lycopodium australianum*

*L. fastigiatum*

### QUILLWORTS

*Isoetes kirkii*

- Agrostis muallieriana*  
*A. muscosa* (unc)  
*A. personata*  
*A. sp.* (unnamed; aff. *A. dyetii*)  
 (panicle spreading, not contracted as in *A. dyetii*)
- Chionochloa australis*  
*C. maera*  
*C. oreophila*  
*C. pallens* ssp.
- C. rubra* ssp.  
*C. sp.* (unnamed) (*C. flavescens* auct. South Island, non *C. flavescens* Zotov)
- C. australis* × *C. pallens* (unc)  
*C. oreophila* × *C. pallens*  
*C. rubra* × *C. sp.* (unnamed)
- Deschampsia novae-zelandiae*  
*Deyeuxia aucklandica*  
*D. avenioides* (incl. *D. a.* var. *brachyantha*)
- Dichelachne crinita*  
*Elymus enysii*  
*E. sp.* (i) (aff. *E. rectisetus*) (lf blade flat, ± glaucous)  
*E. sp.* (ii) (unnamed; aff. *E. rectisetus*) (lf blade flat to rolled)
- Festuca sp.* (i) (unnamed; aff. *F. rubra*)  
*F. sp.* (ii) (*F. matthewsii* sensu Connor 1968)
- Microchloa equisetata* (unc)  
*H. novae-zelandiae*  
*H. recurvata* (unc)
- Koeleria cheesemani*  
*K. sp.* (i) (*K. novozelandica* agg.)  
 (plants tufted; lvs up to 3 mm wide, flat)
- K. sp.* (ii) (*K. novozelandica* agg.)  
 (plants slender, rhizomatous; lvs up to 1.5 mm wide) (unc)
- Lachnagrostis sp.* (i) (unnamed; included in *L. filiformis*, as var. *semiglabra*, by Zotov 1965)
- Microlaena colensoi* (unc)
- Poa breviglumis* (unc)  
*P. buchananii*  
*P. cita* ssp. *cita*  
*P. colensoi*  
*P. dipsacea*  
*P. imbecilla*  
*P. kirkii*  
*P. lindsayi*
- P. pusilla* (unc)  
*P. subvestita*  
*P. sp.* (unnamed; aff. *P. novae-zelandiae*)
- Rytidosperma australe*  
*R. buchananii*  
*R. clavatum*

*R. gracile*  
*R. pumilum*  
*R. setifolium*  
*R. tenue* (unc)  
*Trisetum youngii*  
*T. sp. (i)* (*T. spicatum* agg.)  
*T. sp. (ii)* (unnamed) (plants slender;  
 lf sheath vs. glabrous; lf blade  
 up to 2.5 mm wide, margins  
 smooth) (unc)

*C. sinclairii*  
*C. wakatipu* ss.  
*C. sp.* (undetermined)  
*Carpha alpina* ss.  
*Eleocharis acuta*  
*E. gracilis*  
*E. pusilla*  
*Isolepis aucklandica*  
*I. basilaris* (unc)  
*I. caliginis* (unc)  
*I. habra* (unc)  
*I. subtilissima*  
*Oreobolus pectinatus*  
*Schoenus pauciflorus* ss. (= "slender  
 form" of Edgar (1970, p. 194))  
*Uncinia clavata* (unc)  
*U. divaricata*  
*U. fuscovaginata* var.  
*U. nervosa* ss. (unc)  
*U. tubra* (unc)

### SEDGES

*Carex berggrenii* (unc)  
*C. breviculmis*  
*C. capillacea* (unc)  
*C. carsei* (unc)  
*C. cockayneana* (unc)  
*C. colensoi* (unc)  
*C. comans*  
*C. coriacea*  
*C. diandra*  
*C. echinata* var. (*C. stellulata* var.  
*australis*)  
*C. enysii* (unc)  
*C. flagellifera* (unc)  
*C. flaviformis*  
*C. gaudichaudiana*  
*C. goyenii* (unc)  
*C. kaloides*  
*C. muelleri*  
*C. petrici*  
*C. pyrenaica* var. *cephalotes*  
*C. resectans* (unc)  
*C. secta* ss.

### RUSHES AND ALLIED PLANTS

*Centrolepis ciliata*  
*Gammaria setacea* (Cheeseman  
 1882)  
*Juncus antarcticus*  
*J. novae-zelandiae*  
*J. pusillus*  
*Luzula leptophylla*  
*L. migrata*  
*L. pumila*  
*L. rufa* ss.  
*L. subclavata* (unc)

- L. traversii s.s.
- L. sp.(i) (unnamed; included in L. tufa, as var. albicomans, by Edgar 1966)
- L. sp.(ii) (unnamed; included in L. crinita, as var. petricana, by Edgar 1966)
- L. sp.(iii) (unnamed; included in L. picta, as var. lunosa, by Edgar 1966)
- L. sp.(iv) (unnamed; aff. L. tufa) (plants stoloniferous; infl head smaller than in L. tufa)
- Marsippospermum gracile

- Calmisia alpina s.s.
- C. bellidioides
- C. gracilentata ssp.
- C. haastii ssp. haastii (unc)
- C. incana (incl. C. discolor)
- C. laticifolia
- C. lateralis (incl. C. l. var. villosa)
- C. monroi ssp. monroi
- C. m. ssp. (C. semicordata ssp. semicordata)
- C. sessiliflora
- C. sinclairii (incl. C. allanii and C. dorietzii)

**MONOCOT HERBS OTHER THAN ORCHIDS, GRASSES, SEDGES, RUSHES AND ALLIED PLANTS**

- Astelia nervosa s.s. (unc)
- A. petrici (unc)
- Bulbinella hookeri
- Lemna sp. (L. minor auct.)
- Phormium cookianum (unc)
- Potamogeton cheesemanii
- P. ochreatus
- P. suboblongus
- Typha orientalis

- C. spectabilis ssp. spectabilis
- C. viscosa
- C. sp. (unnamed; aff. C. gracilentata) (plants strongly rhizomatous)
- C. monroi X C. spectabilis (unc)
- C. sessiliflora X C. spectabilis (unc)
- Craspedia lanata s.s.
- C. sp.(i) (unnamed; included in C. lanata, as var. elongata, by Allan 1961)
- C. sp.(ii) (unnamed) (plants summer-green; lvs green, both surfaces densely clad in multicellular non-glandular hairs with cottony tips, over short glandular hairs; margins ± white with cottony hairs; fls white)
- C. sp.(iii) (unnamed) (plants summer-green; lvs green, both surfaces clad in short glandular hairs; margins ± faintly white with cottony hairs; fls white)
- C. sp.(iv) (aff. C. sp. iii, but lvs smaller and narrower) (dried-up taro)

**COMPOSITE HERBS**

- Abrotanella caespitosa (unc)
- Anaphalis sp. (Helichrysum bellidioides s.s.)
- Brachyglottis bellidioides s.s.
- B. lagopus var. lagopus
- Bradyscome radicata (unc)
- B. sinclairii s.s.

*C. lanata* X *C. sp.* (iii)

*Dolichoglottis lyallii*

*Gnaphalium avdax*

*G. laterale*

*G. limosum* s.s. (unc)

*G. mackayi* s.s.

*G. nitidulum*

*G. polylepis* (unc)

*G. sphaericum* (unc)

*G. traversii*

*G. mackayi* X *G. nitidulum* (unc)

*Haastia pulvinatis* s.s.

*H. recurva*

*H. sindarii* s.s. (unc)

*Melichrysom filicaule*

*Lagenifera barkeri* (unc)

*L. cuneata*

*L. petiolata* (unc)

*L. strangulata* (unc)

*Leptinella dandyi*

*L. dioica* ssp. *dioica*

*L. pusilla*

*L. pyrrhifolia* s.s.

*L. sp.* (i) (unnamed; included in

*L. squalida*, as ssp. *mediana*, by  
Lloyd and Webb, 1987)

*L. sp.* (ii) (unnamed) (lvs dimpled)

*L. dandyi* X *L. pyrrhifolia* (unc)

*Leucogenes grandiceps*

*L. sp.* (unnamed; aff. *L. leontopodium*)

*Microseris scapigera* var.

*Pseudognaphalium sp.* (i) (unnamed; aff. *P. luteoalbum*) (lvs mostly oblong spatulate to narrowly oblong, rounded; involueral bracts light brown)

*P. sp.* (ii) (unnamed; included in *Gnaphalium luteoalbum*, as var. *compactum*, by Kirk 1899) (plants small, depressed, much branched, grey) (unc)

*Raoulia apicimgra* s.s.

*R. australis* agg.

*R. bryoides*

*R. eximia* var. *eximia* (unc)

*R. glabra*

*R. grandiflora* var. *grandiflora*

*R. parkii*

*R. subsericea*

*R. subulata*

*R. tenuicaulis* (incl. *R. t.* var. *dimorpha* and *R. t.* var. *pusilla*)

*R. sp.* (unnamed) (stems forming open patches; lvs rigid, recurved, subacute; upper surface grey-green, lower white)

*Senecio glaucophyllus* ssp. (aff. ssp. *toa*) (unc)

*S. wairauensis*

*Taraxacum magellanicum*  
*Vittadinia australis* agg. (unc)

*C. sp. (iii)* (*C. debilis* agg.) ("Glossy Leaf" of Pritchard 1957) (unc)  
*C. sp. (iii)* (*C. debilis* agg.) (cf. "Long Style" of Pritchard 1957) (unc)  
*C. sp. (iv)* (*C. debilis* agg.) ("Narrow Petal" of Pritchard 1957) (unc)

**DICOT HERBS OTHER THAN COMPOSITES**

- Acaena anserinifolia*
- A. caesuglavca*
- A. dunicola*
- A. fissistipula*
- A. glabra*
- A. inermis*
- A. juvenca*
- A. profundeincisa*
- A. saccaticupula*
- A. anserinifolia* X *A. inermis*
- Aciphylla aurea*
- A. colensor* var. *colensor* (unc)
- A. monroi*
- A. subflabellata* (unc)
- A. sp. (i)* (unnamed; aff. *A. glaucescens*) (pinnae rigid, not flaccid as in *A. glaucescens*) (unc)
- Anisotome aromatica*
- A. filifolia*
- A. pilifera*
- A. sp.* (unnamed; included in *A. imbricata*, as var. *pratrata*, by Dawson 1961)
- Callitriche petriei* ssp. *petriei*
- Cardamine bilobata* (unc)
- C. sp. (i)* (unnamed) ("Scree Race" of Pritchard 1957)
- C. sp. (ii)* (aff. *C. corymbosa*) (unc)
- A. sp. (ii)* (unnamed)

*C. sp. (v)* (unnamed; aff. *C. corymbosa*) (plants small; stems and petioles clad in hispid hairs; lvs simple or pinnate; fls up to 2mm diam. solitary; siliques c. 1cm long) (unc)

- Centella uniflora*
- Cheesemania fastigata*
- Chionohebe pulvinaris*
- Colobanthus aciculatis*
- C. apetalus* (incl. *C. a. var. alpinus*)
- C. bochananii*
- C. strictus*
- Crassula sinclairii*
- Drosera arcturi*
- Elatine gratioloides*
- Epilobium alsinoides* s.s. (unc)
- E. angustum* (unc)
- E. atriplicifolium* s.s.
- E. brevipes* (Cheeseman 1925)
- E. brunnescens* s.s.
- E. chionanthum* (unc)
- E. chlorifolium*
- E. cockayneianum* (unc)
- E. crassum* (unc)
- E. elegans* (unc)
- E. forbesii* (unc)
- E. glabellum* s.s.
- E. hectorii* s.s.
- E. insulare* (unc)

- E. komarovianum
- E. krullozanum (unc)
- E. macropus
- E. melanocaulon
- E. microphyllum
- E. netteroides (unc)
- E. porphyrium
- E. pubens
- E. pycnostachyum
- E. tenuipes ss.
- E. tasmanicum
- E. sp.(i) (unnamed; included in E. brunnescens, as ssp. minutiflorum, by Raven and Engelman, 1971)
- E. sp.(ii) (unnamed; aff. E. glabellum) (lvs broader than in E. glabellum, very glossy; capsules vs. glabrous, rarely with erect non-glandular hairs) (unc)
- Euphrasia laingii (unc)
- E. zelandica agg. (unc)
- E. sp. (unnamed; aff. E. petrici)

- Forstera tenella
- F. sp. (unnamed; aff. F. sedi-folia)
- Galium parpusillum ss.
- G. sp.(i) (G. propinquum or G. trilobum) (unc)
- G. sp.(ii) (unnamed; aff. G. parpusillum) (plants forming moss-like turf; lvs narrowly elliptic, rounded to subacute) (unc)

- Gentiana corymbifera ss.
- G. grisebachii
- G. sp.(i) (unnamed; included in G. ballidifolia, as var. australis, by Petrie in Cheeseman 1925)
- G. sp.(ii) (unnamed; aff. G. montana) (plants vs. biennial, occ. perennial, much branched from the base)
- Geranium sessiliflorum var. novae-zelandiae
- G. sp. (unnamed; aff. G. microphyllum)
- Geum leiospermum (unc)
- G. sp. (G. parviflorum agg.) (stock short; terminal leaflet about as long as broad; fls up to 13mm diam.)
- Gingidia decipiens (unc)
- G. montana
- Glossostigma diandrum
- Gonocarpus aggregatus (unc)
- G. micranthus ssp. micranthus
- Hydrocotyle heterometra (unc)
- H. microphylla (unc)
- H. sulcata
- H. sp. (unnamed; included in H. novae-zelandiae, as var. montana, by Kirk 1899)
- H. sulcata X H. sp. (?)
- Hypericum japonicum agg. (unc)
- Hypsela rivalis (unc)
- Ischnocarpus novae-zelandiae (Hooker 1864-7 and Cheeseman 1925, as Sisymbrium novae-zelandiae; Allan 1961)

- Kelleria croizatii*  
*K. dieffenbachii*  
*K. laxa*  
*K. villosa* var. *villosa*  
*Lachnocarpa carnosula* (Cheeseman 1882, as *Ligusticum carnosulum*; 1925, as *Anisotome carnosula*)  
*L. diversifolia*  
*Lilaeopsis ruthiana*  
*Limosella lineata* (unc)  
*Lobelia roughii*  
*Mazus radicans* (unc)  
*Mentha cunninghamii* (unc)  
*Montia fontana* s.s.  
*Myosotis laingii* (Cheeseman 1925)  
*M. macrantha* (unc)  
*M. traversii* s.s.  
*M. sp. (i)* (*M. australis* agg.) (*M. saxatilis* Petrie 1918, non *M. saxatilis* Pallas 1795) (fls white)  
*M. sp. (ii)* (*M. australis* agg.) (fls yellow)  
*M. sp. (iii)* (*M. australis* agg.) (fls small, white)  
*Myrtophyllum pedunculatum* ssp. *novae-zealandiae*  
*M. propinquum*  
*M. triphyllum*  
*M. votschii*  
*Neopaxia* sp. (incl. *Montia calycina* and *Claytonia australasica* var. *biflora*)  
*Nertera balfouriana* (unc)
- N. setulosa*  
*Notothlaspi australe* s.s. (incl. *N. rosulatum*)  
*Oreomyrrhis colensoi* s.s. (incl. *O. e.* var. *hispida* and *O. e.* var. *multifida*)  
*O. ramosa* agg.  
*O. sp. (i)* (unnamed; included in *O. colensoi*, as var. *delicatula*, by Allan 1961) (unc)  
*O. sp. (ii)* (unnamed; aff. *O. colensoi*) (unc)  
*Oreostylidium subulatum* (Cheeseman 1882)  
*Ourisia caespitosa* ssp. *caespitosa*  
*O. lactea* ssp. *lactea*  
*O. simpsonii*  
*Oxalis exilis* (unc)  
*O. magellanica*  
*Parahebe decora*  
*P. linifolia* ssp. *linifolia* (unc)  
*P. lyallii*  
*P. decora* × *P. lyallii* (unc)  
*Phyllachne colensoi* var.  
*Plantago lanigera* s.l. (incl. *P. novae-zealandiae*)  
*P. raoulii*  
*P. triandra* s.s.  
*P. uniflora*  
*Potentilla anserinoides* (unc)  
*Pratia angulata* agg.  
*P. macrodon* s.s.  
*Psychrophila obtusa*

*Ranunculus brevis*  
*R. cheesemanii* var. *cheesemanii*  
*R. crithmifolius* ssp. *crithmifolius* (unc)  
*R. foliosus*  
*R. glabrifolius*  
*R. gracilipes*  
*R. insignis*  
*R. limosella*  
*R. verticillatus*  
*R. sp.* (unnamed; included in *R.*  
*lappaceus*, as var. *macrophyllus*,  
 by Kirk 1899)  
*R. crithmifolius* × *R. gracilipes*  
 (Fisher 1965)  
*R. foliosus* × *R. glabrifolius* (unc)  
*R. gracilipes* × *R. insignis* (unc)  
*Schuzerilema nitens*  
*S. pallidum*  
*S. roughii*  
*S. trifoliolatum* (unc)  
*Scleranthus brockiei*  
*S. uniflorus*  
*Stachhousia minima*  
*Stellaria decipiens* (unc)  
*S. gracilentia*  
*S. roughii*  
*Utricularia novae-zelandiae*  
*Viola cunninghamii*  
*Wahlenbergia albomarginata*  
 var. *albomarginata* (ind. *W. pyg-*  
*maea* var. *laxa*, *W. brockiei*,  
 and *W. simpsonii*)  
*W. cartilaginea*

No. of species: 431.

DOUBTFUL AND ERRONEOUS RECORDS

<u>Record</u>	<u>Probable identity of plant on which record based</u>
<i>Acaena adscendens</i> (Cheeseman 1882)	
<i>Cardamine depressa</i> (Cheeseman 1882, 1925)	
<i>Coprosma parviflora</i> (Manson & Guest, 1975)	<i>Coprosma</i> sp. (i) (unnamed)
<i>Epilobium confertifolium</i> (Cheeseman 1882)	
<i>E. vernicosum</i> (Cheeseman 1925)	<i>Epilobium</i> sp. (ii) (unnamed)
<i>Hebe rigidula</i> (Cheeseman 1925, as Veron. Fig.)	<i>Hebe</i> sp. (ii) (unnamed)
<i>H. tetrasticha</i> (Cheeseman 1882, as <i>Veronica tetrasticha</i> )	
<i>Kelleria lyallii</i> (Cheeseman, 1882 and 1925, as <i>Drapetes lyallii</i> )	<i>Kelleria croizatii</i>
<i>Leucogenes leontopodium</i> (Kirk 1899, as <i>Helichrysum leontopodium</i> ; Cheeseman 1925)	<i>Leucogenes</i> sp. (unnamed)
<i>Oursia colensoi</i> (Cheeseman 1882)	<i>Oursia lactea</i> ssp. <i>lactea</i>
<i>Plantago triantha</i> (Cheeseman 1882, as <i>P. brownii</i> )	<i>Plantago lanigera</i>
<i>Ranunculus pinguis</i> (Cheeseman 1882)	
<i>Uncinia filiformis</i> (Manson and Guest, 1975)	

REFERENCES

- Allan, H.H. 1961: "Flora of N.Z.". Govt. Printer, Wgtn. Vol. 1. 1885 pp
- Cheeseman, T.F. 1882: Contributions to a flora of the Nelson provincial district. Trans. N.Z. Inst. 14: 301-329.
- 1925: "Manual of the N.Z. Flora". Ed. 2. Edited by W.R.B. Oliver. Govt. Printer, Wellington. 1163 pp.
- Connor, H.E. 1968: Interspecific hybrids in hexaploid N.Z. Festuca. N.Z. J. Bot. 6: 295-308.
- Crookes, M.W. 1963: "N.Z. Ferns". Whitcombe & Tombs. 408 pp.

- Dawson, J.W. 1961: Univ. Calif. Pubes Bot. 33: 1-98.
- Edgar, E. 1966: Luzula in N.Z. N.Z. J. Bot. 4: 159-84.  
 ——— 1970: In "Flora of N.Z." Vol. 2. 354 pp.
- Fisher, F.J.F. 1965: "The Alpine Ranunculi of N.Z." N.Z. D.S.I.R. Bull. 165.  
 192 pp.
- Franklin, D.A. 1962: The Ericaceae in N.Z. (Gaultheria and Pernettya)  
Trans. R. Soc. N.Z. Bot. 1: 155-73.
- Hooker, J.D. 1864-7: "Handbook of the N.Z. Flora." London.
- Kirk, T. 1899: "The Students' Flora of New Zealand." Government  
 Printer, Wellington. 408 pp.
- Lloyd, D.G.; Webb, C.J. 1987: N.Z. J. Bot. 25: 99-105.
- Manson, B.R.; Guest, R. 1975. Protection forests of the Wairau  
 catchment. N.Z. J. For. Sci. 5: 123-42.
- Moore, L.B. 1961: In "Flora of N.Z." Govt. Printer. Vol. 1. 1085 pp.
- Oliver, W.R.B. 1935: "The genus Coprosma." Bishop Mus. Bull. 132. 207 pp.
- Pritchard, G.G. 1957: Experimental taxonomic studies on species of  
Cardamine Linn. in New Zealand. Trans. R. Soc. N.Z. 85: 75-89.
- Raven, P.H.; Engelhorn, T. 1971: New taxa and new combinations in  
 Australasian Epilobium (Onagraceae). N.Z. J. Bot. 9: 345-50.
- Wardle, P. 1968: The taxonomy and distribution of the stipitate species  
 of Pseudopanax in New Zealand. N.Z. J. Bot. 6: 226-36.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New  
 Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.