


INDIGENOUS VASCULAR PLANTS OF SPENSER MOUNTAINS,
2500-7550 FT

[†]Highest peak: Mt Una, grid ref. S40 980158

A.P.Druce, Botany Division, D.S.I.R., Feb. 1979 (revised June 1979)
 (Based on visit Jan. 1979)
 (unc-uncommon)

June 1978
 Feb. 1978
 June 1979
 Aug. 1985
 Aug. 1986
 Oct. 1988
 Sept. 1990
 July 1991
 July 1992
 Apr. 1993)



0 5 Miles

[GYMNOSPERM TREES AND SHRUBS]

Dacrydium bidwillii
Phyllocladus alpinus agg.
Podocarpus nivalis
P. hallii × *P. nivalis* (unc)

C. linariifolia (unc)
C. propinquua ssp. *propinquua*
C. perpusilla ssp. *perpusilla*
C. pseudocuneata ssp.
C. rugosa ssp. *rugosa* (unc)
C. serrulata

[DICOT TREES]

Dracophyllum traversii (include
D. pyramidalis) (unc)

C. sp. (i) (unnamed; included
in *C. parviflora* by Oliver,
1935, and others)

Fuchsia excorticata (unc)

C. sp. (ii) (unnamed; aff *C.*
pseudocuneata) (low-growing
shrub with small, thick, arcuate vs)

Griselinia littoralis

Cotatia angustissima

Hoheria lyallii var. (*H. glabrata*)

C. plumosa

Nothofagus fusca

C. satmentosa × *C. sp.* (unc)
var. *ternatus* and as var. *flordensis*,
Cyathodes dealbata (incl.
by Wardle 1968)
C. punica

N. menziesii

N. solandri var. *cliffortioides*

Pseudopanax sp. (unnamed;
included in *P. colensoi*, as

var. *ternatus* and as var. *flordensis*,
Cyathodes dealbata (incl.
by Wardle 1968)

[DICOT SHRUBS]

Aristotelia fruticosa var.

Discaria toumatou

Brachyglottis bidwillii var. *viridis*

Diacophyllum palustre

↗ *B. rotundifolia* (incl. *B. buchananii*
and *B. cockaynei*)

D. primum

Cassinia vauvilliersii

D. uniflorum

Coprosma atropurpurea

Gaultheria crassa

C. cheesemanii ss.

G. depressa ss.

C. ciliata ssp.

↗ *G. sp.* (unnamed; included

C. colensoi ss. (unc)

in *G. depressa*, as var.

C. depressa

novae-zelandiae, by

C. foetidissima

Franklin 1962)

↗ *B. grayi* (Cheeseman 1925, and Allan 1961,
as *Senecio laxifolius*)

G. nubicola

Traversia baccharoides

Gaultheria classa X *G. sp.* (unnamed)
G. macrostigma X *G. sp.* (unnamed)

Hebe canterbutiensis
H. ciliolata
H. cockayneana
H. cressoides (Laing 1912)
H. epactidea var.
H. haastii var. (*cf.* *V. humilis*)
H. lycopodioides
H. paucitamosa ss.
H. takaiensis
H. subalpina var. *subalpina*
H. topiaria
H. sp. (*H. odora* agg.)
H. lycopodioides X *H. sp.* (*H. odora* agg.) (unc) *L. scariosum*
Helichrysum intermedium var. (unc)
Leucopogon floscaerulei ss.
Melicytus alpinus s.s.
Myrsine divaricata s.s.
M. nummularia

Olearia arborescens (unc)
O. avicenniifolia (unc)
O. lacunosa var. *lacunosa*
O. nummularifolia var. (unc)
Perilachnandra punica
Paraxilla tetrapetala (Laing 1912,
as *Elytranthe tetrapetala*)
Pimelea oreophila
Pittosporum rigidum var. (*P. crassicaule*)

DICOT LIANES AND RELATED

TRAILING PLANTS

Clematis forsteri

Muehlenbeckia axillaris

Rubus schmidelioides var.
schmidelioides (unc)

LYCOPODES

Lycopodium australiun
L. fastigiatum

FERNS

Blechnum fluitatile agg.
B. perfoliatum
B. procerum
B. sp. (unnamed)
(*mountain* sp.; lower
pinnae not reduced in length
("The Dunedin-Caw Variety" of
Crookes 1963))
B. procerum X *B. sp.* (*B. capense* agg.)
Grammitis billardierei
G. patagonica (unc)
G. poeppigiana
G. sp. (undetermined)

Histiopteris incisa (unc)
Hydrophyllum multifidum
H. villosum

C. crassiuscula ssp. *torta*
C. oreophila
C. pallens ssp. *pilosa*


Hypolepis millefolium
Polystichum cystostegia
P. vestitum

C. rubra ssp. *occulta*
C. sp. (unnamed) (*C. flaves-*
cans auct South Island, non
C. flavescentis Zотов)

ORCHIDS

Adenochilus gracilis
Aporostylis bifolia
Caladenia lyallii
Chiloglottis cornuta
Cetybas macranthus
C. trilobus ss.
Gastrolezia cunninghamii
Prasophyllum colensoi
Pterostylis australis
P. irsoniana

C. crassiuscula × *C. pallens*
C. pallens × *C. rubra*
Deschampsia chapmanii
D. tenella
Deyauxia aucklandica
D. avencoides (incl. *D. a.* var.
brachyantha)



Festuca sp. (i) (unnamed; aff. *F. rubra*)
F. sp. (ii) (*F. matthewsii* sensu
 Connor 1968)

Hierochloe cuprea
H. novae-zelandiae

Koeleria cheesemanii
Lachnagrostis sp. (unnamed;
 included in *L. filiformis*, as
 var. *semiglabra*, by Zотов 1965)

Microlaena colensoi

Poa braviglumiis
P. cita ssp. *cita*
P. colensoi
P. dipsacea
P. imbecilla
P. Kirkii

GRASSES

Elymus enysii
*E. sp. (aff. *E. tectorum*)*
E. enysii × *E. sp. (?)* (unc)
Agrostis dyeri s.s.
A. muelleriana
A. personata
*A. sp. (unnamed; aff. *A. dyeri*)*
 (panicle spreading, not contracted
 as in *A. dyeri*)
Chionochloa australis
C. conspicua ssp. *conspicua*

P. maia

P. pusilla (unc)

P. sublimis ssp. *sublimis*

P. sp(i) (*P. novae-zelandiae* or *P. subvestita*)

P. sp(ii) (unnamed; aff. *P. novae-zelandiae*)
(Ivs much narrower than in *P. n.z.*)

Rytidosperma australe (unc)

R. buchananii

R. gracile

R. nigricans

R. setifolium

Trisetum youngii

T. sp. (unnamed) (plants slender; lf sheath glabrous, or nearly so)

C. petriei (unc)

C. pyrenaica var. *cephalotes*

C. secta s.s.

C. sinclairii

C. sp. (unnamed; aff. *C. testacea*)

Carpha alpina s.s.

Elaeocharis acuta

E. gracilis

Oreobolus impar

O. pectinatus

O. strictus

Schoenus pauciflorus ss

Isolepis aucklandica

I. caligenis (unc)

I. habra

Uncinia caespitosa

U. clavata

U. divaricata

U. dTuca (unc)

U. involuta

U. rubra

U. rupestris (incl. *U. angustifolia*)

U. sp. (unnamed) (*U. affinis* Hamlin 1959, non *U. riparia* var. *affinis* Col. ex Clarke)

RUSHES

SEDGES

Carex acicularis

C. breviculmis

C. comans

C. coriacea

C. dipsacea var. (*C. falhoata*) (unc)

C. dissita s.s.

C. echinata var. (*C. stellulata* var. *australis*)

C. gaudichaudiana

C. goyanii

C. kalooides

Centrolepis ciliata

Gaimardia setacea

Juncus antarcticus

J. novae-zelandiae

J. pusillus (unc)

<i>Luzula colensoi</i>	<i>Celmisia alpina</i>
<i>L. migrata</i>	<i>C. angustifolia</i> (Cheeseman 1925 as <i>C. novae-zelandiae</i>)
<i>L. pumila</i> (Brockie 1946)	<i>C. bellidiooides</i>
<i>L. rufa</i> s.s.	<i>C. glandulosa</i> ssp.
<i>L. sp.(i)</i> (unnamed; included in <i>L.</i> <i>picta</i> , as var. <i>limosa</i> , by Edgar 1966)	<i>C. haastii</i> ssp. <i>haastii</i>
<i>L. sp.(ii)</i> (unnamed; included in <i>L.</i> <i>erinita</i> , as var. <i>petrieana</i> , by Edgar 1966)	<i>C. incana</i> (incl. <i>C. discolor</i>)
<i>L. colensoi</i> × <i>L. sp.(b)</i> (unc)	<i>C. laricifolia</i>
<i>Marsippospermum gracile</i>	<i>C. montroi</i> ssp. (<i>C. semicordata</i>)

MONOCOT HERBS (OTHER THAN ORCHIDS.) /

/GRASSES, SEDGES, RUSHES/

Astelia lineatis var.

A. nervosa s.s.

A. nivicola s.s.

A. petriæ

Bulbinella hookeri (unc)

Phormium cookianum

Potamogeton cheesemanii (unc)

C. viscosa

C. sp. (unnamed; aff. *C. gracilenta*)
(plants strongly rhizomatous)

Claspedia sp.(i) (included in
C. lanata, as var. *elongata*,
by Allan 1961)

C. sp.(ii) (unnamed) (lvs green,
fls white)

C. sp.(iii) (?)

Dolichoglottis lyallii

D. scorzonerooides

D. lyallii × *D. scorzonerooides*

Gnaphalium audax

G. mackayi s.s.

G. paludosum s.s.

G. traversii

Haastia pulvinaris agg.

COMPOSITE HERBS

Abrotanella caespitosa

Anaphalis sp. (*Helichrysum bellidiooides* s.s.)

Brachyglottis bellidiooides ss.

B. lagopus (Laurig 1912, as *Senecio*
lagopus)

Brachyscome radicata

B. sinclairii

<i>H. sinclairii</i> var. <i>sinclairii</i> (tomentum whitish)	<i>A. profundifolia</i>
<i>H. s.</i> var. (tomentum yellowish-white, looser than in var. <i>sinclairii</i>)	<i>A. anserinifolia</i> × <i>A. inermis</i>
<i>H. pulvinaris</i> × <i>H. sinclairii</i> var. <i>sinclairii</i> (unc)	<i>Aciphylla colensoi</i> var. <i>colensoi</i>
<i>H.p.</i> × <i>H. sinclairii</i> var. (unc)	<i>A. ferox</i>
<i>Helichrysum filicaule</i>	<i>A. montoi</i>
<i>Lagenifera strangulata</i>	<i>Anemone tenuicaulis</i>
→ <i>Leucogenes grandiceps</i>	<i>Anisotome aromatica</i>
<i>Microsetis scapigera</i> var.	<i>A. filifolia</i>
<i>Raoulia australis</i> s.s.	<i>A. haastii</i>
<i>R. btyoides</i>	<i>A. pilifera</i>
<i>R. eximia</i> (Brockie 1946)	<i>A. sp. (unnamed; included in A. imbricata, as var. prostrata, by Dawson 1961)</i>
<i>R. glabra</i>	<i>A. pilifer</i> × <i>A. sp. (unnamed)</i> (unc)
<i>R. grandiflora</i> var. <i>grandiflora</i>	
<i>R. subulata</i>	<i>Callitricha petriei</i> ssp. <i>petriei</i> (unc)
<i>R. tenuicaulis</i> (incl. <i>R.t.</i> var. <i>dimorpha</i> and <i>R.t.</i> var. <i>pusilla</i>)	
<i>Senecio lautavensis</i>	<i>Cardamine</i> sp. (i) (<i>C. debilis</i> agg.) (cf. "Glossy Leaf" of Pritchard 1957)
<i>Leptinella dendyi</i>	<i>C. sp. (ii)</i> (<i>C. debilis</i> agg.) ("Scree Race" of Pritchard)
<i>L. dioica</i> ssp. <i>dioica</i>	<i>C. sp. (iii)</i> (<i>C. debilis</i> agg.) ("Tussock Race" of Pritchard)
<i>L. sp. (i)</i> (<i>L. pyrifolia</i> agg.)	<i>C. sp. (iv)</i> (<i>C. debilis</i> agg.) (var. b of Allan 1961) (unc)
<i>L. sp. (ii)</i> (unnamed; included in <i>L. Squalida</i> , as ssp. <i>mediana</i> , by Lloyd and Webb, 1987)	
<i>L. sp. (iii)</i> (aff. <i>L. sp. ii</i> but lf pinnae entire or nearly so)	
DICOT HERBS (OTHER THAN COMPOSITES)	
<i>Acaena anserinifolia</i>	<i>Cheesemania emysii</i> (Cheeseman 1925, as <i>Nasturtium emysii</i>)
<i>A. fissistupula</i>	
<i>A. inermis</i> (incl. <i>A. microphylla</i> var. <i>robusta</i>)	<i>Chionolobus pulvinaris</i>

- Colobanthus aciculatus*
- C. apetalus* (incl. *C.a. var. alpinus*)
- C. sp.* (unnamed; aff. *C. buchananii*)
(discussed by Moore 1961, p. 212)
- Crassula sinclairii* (unc)
- Drosera arcturi*
- D. spathulata*
- Epilobium atriplicifolium* s.s.
- E. brunneans* s.s.
- E. chlorifolium*
- E. cockayneum*
- E. ciliatum* (unc)
- E. glabellum* s.s.
- E. insulare* (Raven & Raven, 1976)
- E. komarovianum*
- E. macropus*
- E. melanocaulon*
- E. microphyllum*
- E. nerteroides*
- E. linnaeoides*
- E. parnitens*
- E. porphyrium*
- E. pubens*
- E. pycnostachyum*
- E. rotundifolium*
- E. rubromarginatum*
- E. tasmanicum*
- E. tenuipes* s.s.
- E. sp.* (unnamed; included in *E. brunneans*, as *ssp. minutiflorum*, by Raven & Engelkorn, 1971)
- E. macropus* X *E. sp.* (*E. nerteroides*?)
- Euphrasia montoi*
- E. sp.* sp. (unnamed; aff. *F. Forstera*/*sedifolia*)
- F. tenella*
- Galium patposillum* ss.
- Gentiana divisa* (Cheeseman 1925)
- G. grisebachii* (unc)
- G. sp. (i)* (unnamed; included in *G. bellidifolia*, as var. *australis*, by Petrie in Cheeseman 1925)
- G. sp. (ii)* (unnamed; aff. *G. montana*) (stems tall, robust; caudine lf pairs (2)-4-10; infl umbelliform)
- G. sp. (iii)* (cf. *G. bellidifolia*)
- G. sp. (iv)* sp. (unnamed; aff. *G. Geranium*/*microphyllum*)
- G. sessiliflorum* var. *novaeglandiae*
- Geum uniflorum*
- G. sp. (i)* (*G. parviflorum* agg.)
(stock short; terminal leaflet about as long as broad; fls upto 13mm diam.)

G. sp.(ii)(*G. parviflorum* agg.) (stock
creeping and branching; terminal
leaflet slightly longer than broad;
fls c. 8 mm diam.)

Gingidia decipiens

G. montana

Gonocarpus aggregatus

G. micranthus ssp. *micranthus*

Gunnera dentata var. *dentata*

Hydrocotyle microphylla (unc)

H. sulcata

H. sp.(unnamed; included in *H.*
novae-zealandiae, as var.*montana*,
by Kirk (899))

Lilaeopsis ruthiana (unc)

Mazus radicans

Mentha cunninghamii

Neopaxia sp. (incl. *Montia calycinia* and

Claytonia austrolasica var. *biflora*)

Montia fontana ss.

Myosotis forstieri

M. traversii ss. (unc)

M. sp. (unnamed; included in *M.*
pygmaea, as var. *drucei*, by Moore
1961)

Myriophyllum triphyllum (unc)

Nertera batfouriana

N. ciliata

{ *Kelliera croizatii*

K. dieffenbachii

K. laxa

K. multiflora

K. villosa

N. sp.(unnamed; aff. *N. dichondrifolia*) (hairs present
on both surfaces of lf, straight,
not hooked as in *N. dichondrifolia*)

Notothlaspi australe s.s. (unc)

N. sp.(unnamed) (*N. australe* avet;
non *Thlaspi australe* Hook.f.)

Oreomyrtus colensoi s.s.

O. sp.(unnamed; aff. *O. ramosa*)

Oreostylidium subulatum

Ottelia caespitosa ssp. *caesp.*

O. lactea ssp. *lactea*

O. sessilifolia ssp. *sessilifolia*

O. simpsonii

O. caespitosa × *O. sessilifolia*
(unc)

Oxalis magellanica

Patahebe cheesemanii

P. decora

P. linifolia ssp. *linifolia* (unc)

P. lyallii

Phyllachne colensoi var.

Plantago lanigera s.l.

P. raoulii

P. uniflora

Pratia angulata agg.

P. macrodon s.s.

Psychrophila obtusa

Ranunculus cheesemanii var. *ches.* (unc) *Viola cunninghamii*

R. foliosus *V. filicaulis*

R. glabrifolius (unc)

R. gracilipes

R. insignis

R. membranifolius

R. reflexus

R. sp. (unnamed; included in *R. lappaceus*, as var. *macrophyllus*, by Kitch 1899)

R. gracilipes × *R. insignis* (unc)

V. lyallii

Wahlenbergia albomarginata

var. *albomarginata* (incl. *W.*

pygmaea var. *laxa*, *W. brockiei*,
and *W. simpsonii*)

No. of species: 375.

R. gracilipes × *R. insignis* (unc)

Rumex flexuosus

Schizeilema nitens

S. pallidum

S. trifoliolatum (Laing 1912)

S. sp. (unnamed; included in *S.*

haastii, as ssp. *cyanopetalum*, by
Domin (1908)

Scleranthus brockiei (unc)

S. uniflorus (unc)

Stackhousia minima (unc)

Stellaria decipiens (incl. *S. parviflora*)

S. gracilenta

Urtica incisa

Utricularia novae-zelandiae

Doubtful or erroneous records not included in main list

<u>Record</u>	<u>Probable identity of plant on which record based</u>
<i>Acaena microphylla</i> (L ¹ , B ²)	<i>Acaena inermis</i>
<i>Cardamine</i> sp. (B, as "C. heterophylla var. uniflora")	
<i>Carex wakatipu</i> (B)	<i>Carex</i> sp. (unnamed)
<i>Chionochloa flavescens</i> (B, as <i>Danthonia flavescens</i>)	<i>Chionochloa</i> sp. (unnamed)
<i>Coprosma virescens</i> (?) (L)	
<i>Corybas rivularis</i> (B, as <i>Cotysanthes rotundifolia</i>)	<i>Corybas macranthus</i> (?)
<i>Craspedia uniflora</i> (B)	<i>Craspedia</i> sp. (i) and/or sp. (ii)
<i>Forstera bidwillii</i> (L, B)	<i>Forstera tenella</i>
<i>Galium propinquum</i> (B, as <i>G. umbrosum</i>)	<i>Galium perpusillum</i> s.s.
<i>Gaultheria rupestris</i> (B)	<i>Gaultheria crassa</i>
<i>Gentiana setifina</i> (B)	
<i>Hebe traversii</i> (B)	<i>Hebe rakaiensis</i> (?)
<i>Isolepis cernua</i> (B as <i>Scirpus cernuus</i>)	<i>Isolepis habra</i>
<i>Lageniphora cuneata</i> (B as <i>Lageniphora cuneata</i>)	<i>Lageniphora strangulata</i> (?)
<i>Myosotis pygmaea</i> (B)	<i>Myosotis</i> sp. (unnamed)
<i>Nertera depressa</i> (B)	<i>Nertera ciliata</i>
<i>N. dichondrifolia</i> (B)	<i>N. sp.</i> (unnamed)
<i>Oreisia crosbyi</i> (B)	
<i>Phyllachne clavigera</i> (B)	<i>Phyllachne colensoi</i>

(cont. next page)

¹ Laing 1912 ² Brockie 1946

RecordProbable identity of plant on which record based*Plantago triandra* (B)*Plantago uniflora**P. triantha* (B, as *P. brownii*)*P. lanigera**Pseudopanax arboreus* (L.as *Noltopanax arboreum*)*P. colensoi* (B, as *Noltopanax*
colensoi)*P. sp. (unnamed)**Ranunculus lyallii* (Kirk 1899)*Senecio minimus* (L, as*Erechtites pteranthoides*)*Senecio waiauensis* (?)

REFERENCES

Allan, H.H. 1961: "Flora of New Zealand." Government Printer, Wellington. Vol. I. 1085 pp.

Brockie, W.B. 1946: Notes on the indigenous vegetation at the head of the Manuā River and Ada Pass, with lists of species. Trans. R. Soc. NZ. 76: 255-61.
1,2) Dawson, J.W. 1961: A revision of the genus Anisotome (Umbrelliferae). Univ. Calif. Publs Bot. 33: 1-98.

Domin, K. 1908: Engl. Bot. Jb. 40: 573-85.

Druy, D.G. 1972: The cluster and solitary-headed cudweeds native to N.Z.: (Gnaphalium Section Eudeton - Composite). N.Z. Jl. Bot. 10: 112-79.

Edgar, E. 1966: Luzula in New Zealand. N.Z. Jl. Bot. 4: 197-84.

Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria + Pernettya). Trans. R. Soc. NZ. Bot. 1: 155-73

Hacker, E. 1903: On new species of grasses in New Zealand. Trans. N.Z. Inst. 35: 377-85

Kirk, T. 1899: "The Students' Flora of New Zealand." Government Printer, Wellington. 408 pp.

Laing, R.M. 1912: Some notes on the botany of the Spenser Mountains, with a list of the species collected. Trans. N.Z. Inst. 44: 60-75.
3)

Moor, L.B. 1961: Caryophyllaceae, Boraginaceae, Orobanchaceae. Pp. 205-19, 806-70, 861-70 in "Flora of New Zealand". Government Printer, Wellington. Vol. I. 1085 pp.

Pritchard, G.G. 1957: Experimental taxonomic studies on species of Carex amne Linn. in New Zealand. Trans. R. Soc. NZ. 85: 75-89.

Raven, P.H.; Englehorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. Jl. Bot. 9: 345-50.

Raven, P.H.; Raven, T.E. 1976: The genus Epilobium (Onagraceae) in Australasia: a systematic and evolutionary study. N.Z. D.S.R. Bull. 216: 321 pp

Wardlaw, P. 1968: The taxonomy and distribution of the stipitate species of Pseudopanax in New Zealand. N.Z. Jl. Bot. 6: 226-36.

Iotov, V.D. 1965: Grases of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46

3) Loyd, D.C.; Webb, C.J. 1987: The reinstatement of Leptinella at generic rank, and the status of the Cotulaceae (Asteraceae, Anthemidinae). N.Z. Jl. Bot. 25: 99-105

1) Cheeseman, T.F. 1925: "Manual of the New Zealand Flora." Ed. 2. (Editor, W.R. G. Oliver). Government Printer, Wellington. 1163 pp.

2) Crookes, M.W. 1963: "N.Z. Ferns." Ed. 6. Whitcombe & Tombs. 408 pp.

