

INDIGENOUS VASCULAR PLANTS OF KEEBLE'S BUSH

MANAWATU

Grid ref. N 149 090295; alt. c. 50m.

A.P.Druce, Botany Division, D.S.I.R., and
 R.M.Greenwood, Applied Biochemistry Division, D.S.I.R.,
 July 1979 (revised May 1980, Aug. 1982, Oct. 1983,
 (Based on observations 1943-1985) July 1984,
 (unc) - uncommon April 1985,
 (Esler 1962, 1978) - recorded by Esler but not Oct. 1988,
 seen by A.P.D. or R.M.G. Sept. 1990
 [] - apparently no longer present

GYMNOSPERM TREES

Dacrycarpus dacrydioides
Dacrydium cupressinum
Podocarpus totara s.s.
Prumnopitys ferruginea
P. taxifolia

Beilschmiedia tawa

Catocalatus senatus (unc)
Elaeocarpus dentatus (incl. E. cl.
 var. *obovatus*)
E. hookerianus (unc)

MONOCOT TREES

Cordyline australis
Rhopalostylis sapida (unc - juv.
 only)

[*Fuchsia excorticata*]

Griselinia lucida (unc)
Hedycarya arborea
Hohoria populea var. *sextostylosa*
Knightia excelsa

DICOT TREES

Alectryon excelsus var. *excelsus*
Aristotelia serrata (unc)

Lauridia novae-zelandiae (unc)
Lophomyrtus bullata (unc)
L. obcordata (Esler 1962)
L. bullata × *L. obcordata* (unc)

Macropiper excelsum var. *excelsum* [DICOT SHRUBS]

Melicytus ramiflorus ssp. *ramiflorus* *Baccharis repanda* s.s.

Metrosideros robusta (unc)

Mida salicifolia (incl. M.S. var
myrtifolia) (unc)

Myoporum laetum var. *laetum* (unc)

Myrsine australis

M. salicina (unc)

Nestegis cunninghamii (unc)

N. lanceolata

Olearia rani var. *colorata* (unc)

Coprosma areolata

C. classifolia (unc)

C. grandifolia

C. lucida s.s. (unc)

C. propinqua ssp. *propinqua* (incl.)

C.p. var. latiuscula (unc)

C. rhamnoides ssp. *rhamnoides*

(incl. *C. polymorpha*) (unc)

C. robusta

C. rotundifolia

C. tenuicaulis (unc)

C. propinqua × *C. robusta*

Cotatia arborea (unc)

Fuchsia excorticata × *F. perscandens* (unc)

Pseudopanax arboreus var. *alboeius*
(unc)

Geniostoma rupestre var.

ligustrifolium

Hebe stricta var. *stricta* (unc)

Korthalsella lindsayi s.s. (unc)

Kunzea ericoides var. (unc)

Leptospermum scoparium var.
(unc)

Metcalfa simplex

Melicytus micranthus (incl. M.m.
var. *microphyllus*)

M.m. × *M. ramiflorus* (unc)

Neomyrtus pedunculata (unc)

Wainmannia racemosa var.
tacamosa

Olearia solandri (unc)
(incl. *O.v. var. ramuliflora*)
O.virgata var. *virgata* (unc)
Pittosporum cornifolium (unc)
Pseudowintera axillaris (unc)
Solanum aviculare var. *aviculare*

Urtica ferox (unc)

[MONOCOT LIANES]

Freylinia baueriana ssp.
banksii

Ripogonum scandens

[DICOT LIANES]

Calystegia tuguriorum

Clematis foetida

C.paniculata

Fuchsia petiolaris

Mettasideros colensoi (incl. *M.c.*
var. *pendens*) (unc)

M. diffusa

M. fulgens (Esler 1962, as *M. scandens*)

M. perforata

Muehlenbeckia australis

M.a. X M.c. complexa (unc)

Patsencia heterophylla

Passiflora tetrandra

Rubus australis (unc)

R.cissoides (Esler 1962, 1978)

R.schmidtioides var. *schmidtioides*

R.squarrosum (Esler 1962?)

[PSILOPSIDS AND LYCOPDS]

Thesium elongata (incl. *T.e.*, ssp.
robusta) (unc)
Lycopodium variatum (incl. *L.billard-*
ieri and *L.novae-zelandicum*) (unc)

[FERNS]

Adiantum cunninghamii (unc)
A.formosum

Anathropteris lanceolata (unc)

Asplenium bulbiferum

A.flaccidum

A.gracillimum

A.oblongifolium

A.polyodon

A.flaccidum X *Asp.* (*A.bulbiferum*
or *A.gracillimum*) (unc)

Blechnum chambersii

B.discolor

B.filiforme

B.fluviale agg.

B.membranaceum

B.sp. (unnamed) (common lowland
sp.; lower pinnae reduced in length)

B.chambersii X *B.membranaceum*

Cyathea dealbata

C.medullaris (unc)

Dicksonia squarrosa

Diplazium australe (unc)

[GRASSES]

- Hymenophyllum denissimum*
H. dilatatum (Esler 1962)
H. flabellatum
H. flexuosum (unc)
H. revolutum (unc)
H. sanguinale (L.) agg.
H. scabrum (Esler 1962, 1978)
Hypolepis ambigua

- Lastreopsis glabella*
L. hispida (unc)
L. microsora ssp. *pentangularis*
Leplopteris hymenophylloides
Poa scaberula (unc)
Pellaea falcata
P. rotundifolia
Phyllocladus diversifolius
P. scandens
Pneumatopteris pennigera
Polystichum richardii
Pteridium esculentum
Pteris tremula (unc)
Rumohra adiantiformis (unc)

Trichomanes venosum**[ORCHIDS]**

- Dendrobium cunninghamii* (unc)
Drymoanthus adversus (unc)
Earia autumnalis
E. mucronata s.s.
Gastrolechia cunninghamii (unc)
Microtis unifolia (unc)
Orthoceras novae-zeelandiae (unc)
Thelemitra longifolia (unc)

[SEDGES]

- Carex dipsacea* var. *dipsacea*
C. dissita s.s.
C. lambertiana
C. maorica
C. secta s.s.
C. solandri
C. virgata
C. sp. (i) (cf. *C. geminata*) (unc)
C. sp. (ii) (unnamed; aff. *C. geminata*) (much larger than *C. geminata*; up to 2.5 cm wide)

- Cyperus ustulatus* (unc)
Eleocharis acuta
Gahnia pauciflora (unc)
G. setifolia (unc)
Isolepis inundata
I. reticulata

- Schoenus maschalinus* (unc)
Uncinia banksii

- U. distans*
U. scabra (unc)
U. uncinata

RUSHES

- Juncus australis*
J. graciliflorus
J. scaphophorus
Luzula picta s.s. (unc)

- Myriophyllum propinquum* (unc)
Oxalis exilis
Parietaria debilis
Piatia angulata agg.
Ranunculus amphitrichus (unc)

MONOCOT HERBS (OTHER THAN ORCHIDS, GRASSES, SEDGES, RUSHES)

- Arthropodium candidum* (unc)
Astelia fragrans
A. solandri (unc)
Colospermum hastatum
Dianella nigra
Lemna sp. (L. minor var.?)

Stellaria decipiens (incl. *S. parviflora*)

Urtica incisa

No. of species: 187.

COMPOSITE HERBS

- Senecio minimus* (Esler 1962, as
Erechtites prenanthoides)

REFERENCES

- Esler, A.E. 1962: Forest remnants of the Manawatu lowlands. *J. Roy. N.Z. Inst. Hort.* 4: 255-68.
— 1978: "Botany of the Manawatu". *D.S.I.R. Inf. Ser.* 127. 206 pp.

DICOT HERBS (OTHER THAN COMPOSITES)

- Acaena anserinifolia* (unc)
Australinia pusilla (unc)
Cardamine sp. (*C. debilis* agg) ("Long-Style" of Pritchard 1956)
Dictionaria sp. (unnamed) (slender; fls small) (unc)
Epilobium heterolepis (unc)
E. rotundifolium (unc)
Haloragis erecta ssp. *erecta* (unc)
Hydrocotyle elongata
H. heteromorpha
H. moschata (Esler 1962)

Pritchard, G.C. 1957: Experimental taxonomic studies on species of *Cardamine* Linn. in N.Z. *Trans. Roy. Soc. N.Z.* 85: 75-

DOUBTFUL AND ERRONEOUS RECORDS OF ESLER (1962)

- Astelia nervosa* (= *A. fragrans*)
Carex tetanaria (= *C. sp. a* &/or *C. sp. b*)
Hebe salicifolia (= *H. stricta* var. *stricta*)
Hydrocotyle dissecta (= *H. elongata*)
Scirpus carolinus (= *Isoloma reticulatum*)
Thesium tannensis (= *T. elongata*)
Uncinia clavata
U. laxiflora

SUPPLEMENT TO LIST 188 (INDIGENOUS VASCULAR PLANTS
OF KEEBLE'S BUSH, MANAWATU)

IMPORTANT WEEDS

(i.e. those actually or potentially of concern in the management of the reserve)

- Acacia dealbata* — tending to spread
 - * *Allium triquetrum* — a couple of patches
 - * *Berberis glaucocarpa* — occasional young plants
 - Calystegia sepium* ssp. — in-floodable areas near willows
 - * *Clematis vitalba* — several plants
 - * *Iris foetidissima* — occasional
 - * *Lonicera japonica* — many plants on leach side of bush
 - Pittosporum crassifolium* — several young plants
 - Rubus fruticosus* agg. — several patches, and appears to be increasing in some fairly open areas in bush
 - Salix fragilis* — spreading along stream where it enters the bush
 - Sambucus nigra* — many plants, particularly under old podocarps
 - Solanum dulcamara* — occasional
 - S. pseudocapsicum* — throughout
 - * *Tradescantia fluminensis* — many patches in floodable areas throughout
- * removal of these species is underway (April 1985)