

VASCULAR PLANTS OF THE PEAKS-TARAMEA², AND KARETI³-
NGAMAI A⁴ AREAS, NORTHERN TARARUA RANGE, 2300⁵-3700 FT

A.P.Druce, Botany Division, D.S.I.R., May 1979 (revised June 1979)
(Based on visits 1955, 1962, 1968, 1979) May 1980
June 1980

¹ Grid ref. N 152 999031, alt. 2866 ft; and N 153 005032, alt. 2920 ft

² Grid ref. N 153 035028, alt. 2246 ft June 1980

→ ³ Grid ref. N 153 005965, alt. 3700 ft Sept 1990
July 1991

⁴ Grid ref. N 153 028992, alt. 3221 ft

⁵ Lower limit of *Chiroschiza flavesans* shrub-tussock land

unc - uncommon; T,N - see footnote

INDIGENOUS PLANTS

[GYMNOSPERM TREES AND SHRUBS]

Dacrydium cupressinum (unc)
(upper limit 2350 ft)

Halocarpus biformis

Pittosporum alpinus

Podocarpus hallii

Prumnopitys ferruginea

Griselinia littoralis

Myrsine salicina

Nothofagus fusca (unc - only (small stand seen)) (T)

Pseudopanax colensoi s.s. (unc - only (plant seen, May 1979)) (N)

P. simplex (incl. *P.s. var. sinclairii*)

Weinmannia racemosa var.
racemosa

[MONOCOT TREES]

Cordyline indivisa (unc)

[DICOT SHRUBS]

Alseuosmia pusilla

Brachyglossis rotundifolia var.
(*B. glauca* (Ait.) (unc))

Coprosma colensoi s.s.

C. depressa (unc) (N)

C. foetidissima

[DICOT TREES]

Carpodetus serratus (unc)

Elaeocarpus hookerianus

Fuchsia excorticata (unc) (T)

T - The Peaks-Taramea area only

N - Kareti-Ngamaia area only

- 2
- C. perpusilla* ssp. *perpusilla* *Pentachondria pumila*
C. pseudocuneata ssp. *pseudocuneata* *Pimelea longifolia* var.
C. sp.(i) (unnamed; included in *C.*) *Pittosporum rigidum* var. *maius*(unc)
 parviflora by Clivs! (1935, and others)
 (If undersurface not minutely hairy
 as is usual in *C. parviflora*; drupe vs.
 unpigmented, occ. purplish red) *Pseudowintera axillaris*
C. sp.(ii) (unnamed; *C. parviflora* var.
dumosa sensu Allan (1961, non *C.p.*
 var. *dumosa* Cheeseman)

Cyathodes empetrifolia

DICOT LIANES

Dracophyllum longifolium var.
 (incl. *D. filifolium* and *D. oliveri*)

D. uniflorum (N)

Gaultheria antipoda (unc)

G. rupestris (N)

G. sp. (unnamed; included in *G.*
dopressa, us var. *novaeseelandiae*,
 by Franklin 1962)

G. antipoda X *G. sp.* (unnamed)(T)

Habe odora s.s. (unc) (T)

H. stricta var. *atkinsonii* (unc) (T)

Laukopogen fasciculatus

Myrsine divaricata s.s.

M. nummularia (N)

Neomyrtus pedunculata

Olearia arborascens (unc)

O. colensoi var. *colensoi*

O. lacunosa var. (*O. alpina*)(unc)

O. virgata var. *migata* (incl. *O.v.*
 var. *tamniflora*)(unc)(T)

Metrosideros diffusa (unc)

Muehlenbeckia australis (unc)

Rubus cissoides var. *cissoides* (unc)

LYCOPODE

Lycopodium austriatum (unc) (N)

L. fastigiatum

L. scariosum

L. vatum (incl. *L. bellidifolii*)

L. novae-zelandicum (unc) (N)

L. volubile (unc) (T)

FERNS

Asplenium flaccidum (unc)

Blechnum colensoi (unc) (T)

B. discolor

B. fluvialite egypt.

B. minus (unc)

B. pann-mariina (T)	[ORCHIDS]
B. procerum (unc)	<i>Aporostylis bifolia</i>
B. vulcanicum (unc) (N)	<i>Clitaglottis cornuta</i> (unc) (N)
B.sp. (unnamed) (common lowland sp.; C. trilobus s.s. (unc) (T) lower primae reduced in length)	<i>Corybas oblongos</i> (unc) (T)
B. procerum X B. sp. (unnamed)	<i>C. trilobus</i> s.s. (unc) (T)
<i>Ctenopteris heterophylla</i>	<i>Lypetanthus antarcticus</i> (N)
<i>Cyathea colensoi</i>	<i>Prasophyllum colensoi</i>
<i>C. smithii</i>	<i>Pterostylis venosa</i> (unc)
<i>Grammitis billardieri</i>	<i>Thelymitra cyanea</i>
<i>G. magellanica</i> ssp. <i>nothofageti</i>	<i>T. hatchii</i> (unc)
<i>G. poeppigiana</i> (unc) (N)	<i>T. longifolia</i> (unc) (T)
<i>Histeropteris incisa</i> (unc)	[GRASSES]
<i>Hymenophyllum revolutum</i>	<i>Agrostis muscosa</i> (N)
<i>H. bivalve</i>	<i>A. personata</i>
<i>H. demissum</i>	<i>Chiococloa conspicua</i> ssp. <i>cunninghamii</i> (unc) (N)
<i>H. multifidum</i>	<i>C. flavescens</i> s.s.
<i>H. peltatum</i> (unc) (N)	<i>C. pallens</i> var. <i>pallens</i> (N)
<i>H. pulcherrimum</i> (unc) (N)	<i>C. rubra</i> var. (T)
<i>H. ratum</i>	<i>C. flavescens</i> X <i>C. pallens</i> (unc) (N)
<i>H. sanguinolentum</i> ags.	<i>C. f. X C. rubra</i> (T)
<i>Hypolepis rufobarbata</i> (unc)	<i>Cottadaria fulvida</i> (T)
<i>Laptopterus superba</i> (unc) (T)	<i>Deyeuxia aucklandica</i>
<i>Paspalum scaberrula</i> (unc)	<i>Microlaena avenacea</i>
<i>Polystichum vestitum</i>	<i>M. colensoi</i> s.s. (unc) (N)
<i>Sticherus cunninghamii</i>	<i>Poa breviglumis</i> (N)
<i>Trichomanes reniforme</i>	<i>P. colensoi</i> (N)
	<i>Rytidosperma gracile</i>
	<i>R. nigriceps</i> (N)

SEDGES

- Carex acicularis* (unc) (N)
C. dissita ss.
C. sp. (unnamed; Section Acutae)
Carex alpina ss.
Gahnia pauciflora
Isolepis aucklandica
I. habra
Lepidosperma austiale (unc) (T)
Oreobolus pectinatus
Schoenus concinnus (unc) (N)
S. sp. (unnamed; aff. *S. pauciflorus*) ("robust form" of Edgar 1970, p. 194)

RUSHES

- Centrolepis ciliata* (N)
Juncus antarcticus
J. gregiflorus
J. novae-zelandiae
J. planifolius
Luzula sp. (unnamed; included in *L. picta*, as var. *limosa*, by Edgar 1966)

MONOCOT HERBS (OTHER THAN)

[ORCHIDS, GRASSES, SEDGES, RUSHES]

- Uncinia astonii*
U. caespitosa
U. clavata (?) (unc)
U. filiformis
U. fuscovaginata var. *fusco-vaginata* (N)

- A. sp.* (unnamed; aff. *A. nervosa*)
(lvs not conspicuously white ad-
axially as in *A. nervosa*)

- U. gracilenta*
U. rupestris (incl. *U. angustifolia*)
U. silvestris (incl. *U. affinis*) (T)
U. uncinata
U. viridis (N)
U. sp. (unnamed; aff. *U. nervosa*)
(more slender than *U. nervosa*;
see Edgar 1970, p. 231) (N)

- Bulbinella gibbsii* var. *balanifera* (unc) (N)

- Libertia peltigera*

- Luzuriaga parviflora*

- Phormium cookianum*

COMPOSITE HERBS

- Abrotanella caespitosa*
A. pusilla (N)
Anaphalis kirkii s.s. (T)
A. sp. (*Helichrysum bellidioides*
s.s.)
Brachyglottis lagopus var. *lagopus*
Celmisia hieracifolia var. (lvs
broader than in var. *hieracifolia* (N))

- C.spectabilis* ssp. *spectabilis* (N) *Forsterotanella* (N)
- C.sp.(unnamed; aff. C.gracilenta)(plants tufted; lvs linear to linear-lanceolate; upper surface and margins thinly covered in tomentum; lower surface glabrous, except for midrib)* *Gentiana montana* (incl. *G.m.* var. *stolonifera*, *G.patula*, and *G.bellidifolia*)
- C.spectabilis* X *C.sp.(unnamed (unc))* (N) *G. sp.(unnamed; aff. G.grisebachii)*
- Gonocarpus aggregatus*
- Craspedia minor* var. *minor* (incl. *C.major*) (unc) *Hydrocotyle dissecta* (T)
- H. heterometa* (unc)
- Gnaphalium limosum* s.s. (unc) (T) *H. micropylla* (T)
- G.mackayi* (unc) (T) *H.meschata* (unc)
- G.wahinicicum* (unc) (T) *Kelleria dieffenbachii*
- Halictysum filicaule* (T) *Liparophyllum gunnii* (N)
- Leucogenes leontopodium* s.s. (unc) *Nertera ciliata*
- (N) *N.depressa* (incl. *N.cunninghamii*) (T)
- Raoulia glabra* (T) *N.sp.(unnamed; aff. N.dichondrifolia)* (hairs present on both surfaces of lf, straight, not hooked as in *N.dichondrifolia*) (unc) (T)
- R.grandiflora* var. *grandifolia* (N) *Oreomytthis colensoi* s.s. (incl. *O.c.* var. *hispieta* and *O.c.* var. *multifida*) (T)

DICOT HERBS (OTHER THAN COMPOSITES)

- Acaena anserinifolia* (unc) *Phyllachne colensoi* (N)
- Aciphylla colensoi* var. *colensoi* (unc) (N) *Plantago lanceolata* s.l. (N)
- A. dissecta* (N) *P.uniflora* (N)
- Anisotome aromatica* (N) *Pratia angulata* agg. (T)
- Drosera stenopetala* (N) *Psychrophila novae-zealandiae* (N)
- Epilobium alsinoides* s.s. (T) *Ranunculus reflexus* (unc) (T)
- E. cockayneanum* (unc) *R.verticillatus*
- E.linnaeoides* (unc) (T) *Schizolejeunea* sp. (unnamed) (lvs cut $\frac{1}{3}$ - $\frac{2}{3}$ to base; scattered hairs on upper surface; involucral bracts obtuse; fr = ol < its pedicel)
- E.parvitens* (unc) (N)
- Euphrasia cuneata*
- E.revoluta* (N)

Stellaria decipiens (incl. S. paniciflora) (unc)

Viola filicaulis

ADVENTIVE PLANTS

GRASSES

Agrostis capillaris

Holcus lanatus (unc) (T)

Poa annua (unc) (T)

SEDGES

Carex demissa (unc) (N)

RUSHES

Juncus articulatus

J. effusus (N)

Luzula congesta (unc) (N)

DICOT HERBS

Cirsium arvense (unc) (T)

C. vulgare (unc) (T)

Hypochaeris radicata

Leontodon tataricoides (unc) (T)

Sagina procumbens (T)

No. of indigenous species: 193. (145 The Peaks-Taramea area)
 (165 Karetí-Ngamaia area)

No. of adventive species: 12.