

(20)

INDIGENOUS VASCULAR PLANTS OF GUNNER DOWNS[†],
N.W. NELSON, 3000-4000 FT (900-1200 M)

A.P. Druce, Botany Division, D.S.I.R., April 1980

(revised May 1980, May 1981, May 1982, Sept 1982

Based on visit Nov. 1979

June 1983

June 1984

July 1985

Oct. 1986

[†]Highest point: Mt Barr, grid ref. ST 627618

(rocks: granite)


Nov. 1988

(unc) - uncommon

July 1990

July 1991

March 1992



GYMNOSPERM TREES AND SHRUBS

- Halocarpus biformis*
Lepidothamnus laxifolius
Libocedrus bidwillii
Phyllocladus sp. (unnamed; aff. *P. alpinus*) ("lowland" ecotype of Wardle 1969)
Podocarpus nivalis (unc)
P. hallii × *P. nivalis* (unc)

DICOT SHRUBS

- Archetia traversii* (incl. *A.t. var. australis*)
Brachyglottis rotundifolia var. (incl. *B.buchananii* and *B. cockaynei*)
Cassinia vauvilliersii

DICOT TREES

- Dracophyllum traversii* (incl. *D. pyramidale*)
Elaeocarpus hookerianus (unc)
Griselinia littoralis (unc)
Metrosideros umbellata
Nothofagus menziesii
N. solandri var. *cliffortioides*

- Coprosma cheesemanii* ss.
C. ciliata ssp. *ciliata*
C. colensoi ss.
C. crenulata
C. foetidissima
C. perpusilla ssp. *perpusilla*
C. pseudocuneata ssp.
C. rhomboides (incl. *C. polymorpha*)

- Pseudopanax linearis*
P. simplex (incl. *P.s. var. sinclairii*)
P. sp. (unnamed; included in *P. colensoi*, as var. *fiordensis* and as var. *tarnatus*, by Wardle 1968)
Quintinia serrata (incl. *Q. acutifolia* and *Q. elliptica*)

- C. sp. (i)* (unnamed) (*C. parviflora* var. *dumosa* sensu Allan 1961, non *C.p. var. dumosa* Cheeseman)
C. sp. (ii) (unnamed; aff. *C. colensoi*) (low-growing shrubs with trailing branches)

- Weinmannia racemosa* var. *racemosa*

- Cyathodes dealbata* (incl. *C. pumila*)
C. sempervirens
Dracophyllum longifolium var. (incl. *D.filifolium*, *D. diveri*)
D. palustre
D. pubescens var. *pubescens*

D. uniflorum var. *uniflorum*
D. longifolium × *D. uniflorum*

Epactis alpina (unc)

Gaultheria macrostigma
G. rupestris (incl. *G.*
subcorymbosa) (unc)

G. sp. (unnamed; included in *G.*
depressa, as var. *novae-*
zelandiae, by Franklin 1962)

Gaultheria macrostigma ×
G. sp. (unnamed)

Hebe canterburiensis

H. odora s.s.

H. topiaria (unc)

H. sp. (*Leionchebe mooreae*)

Leptospermum scoparium

Leucopogon fraseri s.s. (unc)

Melicytus alpinus s.s. (unc)

Myrsine divaricata s.s.

M. nummularia

Olearia colensoi var. *colensoi*

O. lacunosa var. *lacunosa*

Pentachondra pumila

Pimelea suteri var. (*P. oreophila*)

Pittosporum rigidum var. (*P. crassi-*
caule)

Pseudopanax anomalus

Pseudowintera colorata

P. traversii (unc)

PSILOPSIDS AND LYCOPODS

Tmesipteris tannensis

Lycopodium australianum (unc)

L. fastigiatum

L. scariosum

L. varium (incl. *L. billardierei* and
L. novae-zelandicum)

FERNS

Asplenium flaccidum

Blechnum discolor

B. fluviatile agg.

B. pennata-marinia

B. procerum

B. sp.(i) (unnamed) (common lowlands
lower pinnae reduced in length
(unc))

B. sp.(ii) (unnamed) (in mountaini
sp.; lower pinnae not reduced in
length) ("The Dunedin-Cass Variety
of Crookes 1963")

Grammitis billardierei

G. givenii (unc)

G. magellanica ssp. *nootrofageti*

G. poeppigiana (unc)

Hymenophyllum armstrongii
H. malingii
H. multifidum
H. farum
H. revolutum
H. rufescens
H. villosum
Leptopteris superba
Polystichum vestitum(unc)

Hierochloe norae-zelandiae
Microlepia avenacea
M. colensoi
M. thomsonii(unc)
Poa colensoi
Rytidosperma gracile
R. nigricans

ORCHIDS

Aporostylis bifolia
Chiloglottis cornuta
Corybas acominatus(unc)
Liparophyllum antarcticus
Prasophyllum colensoi
Pterostylis venosa
Thelymitra cyanea(unc)

SEDGES

Carex acicularis(unc)
C. libera
C. sp.(i)(aff. C. dissita)
C. sp.(ii)(cf. C. gaudichaudiana)
C. sp.(iii)(aff. C. coriacea)
Carex alpina
Gahnia procera
Oreobolus impar
O. pectinatus
O. strictus

GRASSES

Agrostis dyeri s.s.
A. personata

Chionochloa australis
C. conspicua ssp. conspicua
C. rubra ssp. occulta
Deyeuxia aucklandica
D. avenoides (incl. D. a. var.
brachyantha)(unc)

Schoenus pauciflorus s.s.
Isolepis aucklandica
Uncinia astonii
U. filiformis
U. fuscovaginata var. (glaucoous) plants
U. involuta (unc)
U. rupestris (incl. U. angustifolia)
U. uncinata
U. sp. (unnamed; aff. U. nervosa) (more
slender than U. nervosa; see Edgar 1970, p. 23)

RUSHES AND ALLIED PLANTS

- Centrolepis ciliata*
Empodisma minus
Gaimardia setacea
Juncus antarcticus
J. novae-zelandiae
J. pusillus
Luzula migrata (unc)
L. sp. (unnamed; included in *L. picta*, as var. *linosa*, by Edgerton 1966)

- Brachyglottis* sp. (unnamed; included in *B. bellidifloroides*, as var. *crassa*, by Nordenstam 1978)
Celmisia alpina s.s.
C. armstrongii
C. dallii
C. incana (incl. *C. discolor*)
C. laticifolia
C. lateralis (incl. *C. l. var. villosa*)
C. patra
C. similis
C. sp. (unnamed; aff. *C. gracilenta*) (lvs linear to linear-lanceolate; lower surface glabrous)
C. armstrongii × *C. patra* (unc)
C. a. × *C. similis* (unc)

MONOCOT HERBS (OTHER THAN ORCHIDS)

GRASSES, SEDGES, RUSHES AND ALLIED PLANTS

- Astelia linearis*
A. nivicola s.s.
A. skottsbergii
A. sp.(i) (unnamed; included in *A. nivicola*, as var. *moriceae*, by Moore 1966)

- A. sp.(ii) (unnamed; aff. *A. nervosa*)
Harpotrichion novae-zelandiae (unc)
Libertia pulchella
Luzuriaga parviflora
Phormium cookianum

Craspedia sp. (plants summer-green; lvs small, elliptic-spathulate, glabrous; blade rather abruptly narrowed to slender petiole; both surfaces clad in hispid hairs up to c.1 mm long; fls white)

- Gnaphalium mackayi* s.s.
G. paludosum s.s.
Leptinella sp. (unnamed; included in *L. squalida*, as ssp. *mediana*, by Lloyd and Webb, 1987)

- Raoulia grandiflora* var. *grandiflora*

DICOT HERBS (OTHER THAN COMPOSITES)

- Aciphylla colensoi* var. *colensoi* (unc)
A. hookeri (unc)
Actinotis novae-zelandiae

- Anisotome aromatica*

- A. deltoidea* (unc)

- A. sp. (unnamed) (turf-forming in seepages; infls short, up to 5 cm) (unc)

COMPOSITE HERBS

- Abrotanella caespitosa*
A. linearis s.s.
Anaphalis sp. (*Helichrysum bellidioides* s.s.)

Cardamine sp. (*C. dubilis* agg.)
("Glossy Leaf" of Pritchard
(1957) (unc)

Colobanthus affinis

Donatia novae-zelandiae

Drosera arcturi

D. spathulata

D. stenopetala

Epilobium cockayneanum

E. petnites

Euphrasia townsonii s.s.

Forstera mackayi

F. tenella

Gentiana montana (incl. *G. m. var. stolonifera*, *G. patula* and *G. bellifolia* var. *pulchella*)

G. spenceri

Gentianum sp. (aff. *G. microphyllum*)

Geum uniflorum

Gonocarpus aggregatus

Hydrocotyle sp. (undetermined) (unc)

Kelleria dieffenbachii

Klaxa

Liparophyllum gunnii (unc)

Mitrasacme novae-zelandiae (unc)

Montia fontana s.s.

Nertera balfouriana

N. sp. (unnamed; aff. *N. dichondrifolia*)

(hairs present on both
surfaces of lf, straight, not hooked
as in *N. dichondrifolia*)

Oreomythris colensoi s.s. (incl. *O.c.*
var. *hispida* and *O.c. var. multifida*)

Oreostylidium subulatum

Ouria macrocarpa ssp. *calycina*

O. simpsonii

O. macrocarpa X *O. simpsonii* (unc)

Oxalis magellanica

Plantago novae-zelandiae agg.

P. uniflora

Pratia angulata ass.

Psychrophila novae-zelandiae (unc)

Ranunculus verticillatus

Schizolema nitens

Viola filicaulis (unc)

No. of species: 198.

REFERENCES

- Allan, H.H. 1961: "Flora of New Zealand." Government Printer, Wellington. Vol. 1. 1085 pp.
- Crookes, M. 1963: "New Zealand Ferns." Whitcombe & Tombs. 408 pp.
- Edgar, E. 1966: Huzula in New Zealand. N.Z. Jl Bot. 4: 159-84.
- 1970: Cyperaceae. Pp. 167-285 in "Flora of New Zealand". Government Printer, Wellington. Vol. 2. 354 pp.
- Franklin, D. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya) Trans R.Soc. N.Z. Bot. 1: 155-73.
- Moore, L.B. 1961: Habe (except whipcord species). In "Flora of New Zealand." Government Printer, Wellington. Vol. 1. 1085 pp.
- 1966: Australasian asteliads (Liliaceae). N.Z. Jl Bot. 4: 201-40.
- Nordenstam, B. 1978: Opera Bot. 44: 1-84
- Pritchard, G. G. 1957: Experimental taxonomic studies on species of Cardamine Linn. in New Zealand. Trans R. Soc. N.Z. 85: 75-89.
- Wardle, P. 1968: The taxonomy and distribution of the stipulate species of Pseudopanax in New Zealand. N.Z. Jl Bot. 6: 226-36.
- Lloyd, D.G.; Webb, C.J. 1987: The reinstatement of heptinella as generic rank, and the status of the 'Cotulaceae' (Asteraceae, Anthemideae). N.Z. Jl Bot. 25: 99-105.
- Wardle, P. 1969: Biological flora of New Zealand. Phyllocladus alpinus Hoch f. (Podocarpaceae); mountain toatoa, celery pine. N.Z. Jl Bot. 7: 76-95.

