


INDIGENOUS VASCULAR PLANTS OF DENNISTON AND STOCKTON PLATEAUX (INCLUDING MTS AUGUSTUS, FREDERICK, AND ROCHFORT)
N.W. NELSON, 2000-3620 FT (600-1100 M)

A.P.Druce, Botany Division, D.S.I.R., June 1981
 (revised Jan 1982, May 1982, June 1982, Sept. 1982,

June 1983
 June 1984
 Nov. 1984
 May 1985
 July 1985
 June 1986
 Aug. 1986
 June 1987
 June 1988
 Aug. 1990
 Dec. 1990
 May 1991
 Aug. 1991
 Aug. 1992
 Aug. 1993)

(Based on visits Jan. 1980 and Jan 1987)

- 1 Grid ref. S24 325820; alt. 3316'
- 2 Grid ref. S24 315786; alt. 3620'
- 3 Grid ref. S31 218698; alt. 3417'



seen or recorded:

- (D) — in Denniston Plateau - Mt Rochfort area
(S) — in Stockton Plateau - Mt Augustus - Mt Frederick area

GYMNOSPERM TREES AND SHRUBS

Dacrydium cupressinum (D,S)

Halocarpus bidwillii (D,S)

H. biformis (D,S)

Lepidothamnus intermedius (D,S)

L. laxifolius (D,S)

L. intermedius x *L. laxifolius* (D,S)

Libocedrus bidwillii (D,S)

Phyllocladus alpinus agg. (D,S)

Podocarpus hallii (D,S)

Quintinia serrata (incl. *Q. acutifolia* and *Q. elliptica*) (D,S)
Weinmannia racemosa var. *racemosa* (D,S)

DICOT SHRUBBS

Archeia traversii (incl. *A.t.* var. *australis*) (D,S)

Brachyglottis rotundifolia var. (incl. *B. buchananii* + *B. cockaynei*) (D,S)

Cassinia vauvillietii (D,S)

MONOCOT TREES

Cordyline australis (S)

C. indivisa (D,S)

Coprosma cheesemanii s.s. (D)

C. ciliata ssp. *ciliata* (D)

C. colensoi s.s. (D,S)

C. crenulata (D,S)

C. foetidissima (D,S)

C. pseudocuneata ssp. (D,S)

C. sp. (i) (unnamed; aff. *C. colensoi*)

(low-growing shrubs with trailing branches) (D,S)

C. sp. (ii) (unnamed; included in

C. parviflora by Oliver, 1935,

and others) (If undersurface

not minutely hairy as is usual

in *C. parviflora*; oltypes vs. un-

pigmented, occ. purplish red) (D)

C. sp. (iii) (unnamed) (*C. parviflora*

var. *dumosa* sensu Allan 1961,

non *C. p. var. dumosa* Cheeseman) (S)

DICOT TREES

Dracophyllum traversii (incl. *D. pyramidale*) (D)

Elaeocarpus hookerianus (D,S)

Griselinia littoralis (D,S)

Metrosideros parkinsonii (D,S)

M. umbellata (D,S)

Nothofagus fusca (D)

N. menziesii (D,S)

N. solandri var. *cliffortioides* (D,S)

Pseudopanax linearis (D,S)

P. simplex (incl. *P. s.* var. *sinclairii*) (D,S)

P. sp. (unnamed; included in *P. colensoi*, as var. *fiordensis*, and as var. *tettnatus*, by Wardle 1968) (D,S)

- 5
- Cyathodes dealbata* (incl. *C. pumila*) (D,S)
C. empetrifolia (D,S)
C. sp. (*C. juniperina* agg.) (vs vs.
 >1 cm x >1 mm, up to 1.5 cm x
 2.5 mm) (D,S)
Dracophyllum longifolium var.
 (incl. *D. filifolium* and *D. oliveti*) (D,S)
D. palustre (D,S)
D. politum (incl. *D. densum*) (D,S)
D. pubescens var. *pubescens*
 (D,S)
D. uniflorum (D,S)
Epactis alpina (D,S)
E. pauciflora (D,S)
Exocarpus bidwillii (D,S)
Gaultheria antipoda (D,S)
G. rupestris (incl. *G. subcorymbosa*) (D,S)
G. sp. (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962) (D,S)
G. macrostigma X *G. sp.* (unnamed) (D)
Hebe canterburiensis (D,S)
H. salicifolia var. *salicifolia* (D)
H. sp. (*Leonohebe mooreae*) (D,S)
- Melicytus* sp.(unnamed) (S)
Myrsinæ divaricata s.s. (D,S)
Neomyrtus pedunculata (D)
Olearia arborescens (D)
O. colensoi var. *colensoi* (D,S)
O. ilicifolia (D)
O. arborescens X *O. ilicifolia* (D)
Pentachondra pumila (D,S)
Pimelea gnidioides var. (D,S)
P. longifolia var. (D,S)
Pittosporum rigidum var. (*P. crassicaule*) (D,S)
Pseudowintera traversii (D,S)
Solanum laciniatum (S)
- [DICOT LIANES]**
- Metrosideros fulgens* (D)
- [PSILOPSIDS AND LYCOPODS]**
- Tmesipteris tannensis* (D)
Lycopodium australe (D,S)
L. carnuum (D)
L. fastigiatum (D,S)
L. laterale (incl. *L. tamulosum*?)
 (D,S)
L. scariosum (D,S)
L. variuum (incl. *L. billardierei*
 and *L. novae-zelandicum*) (D,S)

[FERNS]

- Asplenium flaccidum* (D,S)
Blechnum discolor (D,S)
B. fluviatile agg. (D,S)
B. minus (D,S)
B. penna-marina (D)
B. procerum (D,S)
Leptospermum scoparium (D,S)
Leucopogon fasciculatus var. (D,S)

B. vulcanicum (D)
B. sp. (unnamed) (common lowland sp.; lower pinnae reduced in length) (D,S)
B. procerum × *B. sp. (unnamed)* (D)
Dicksonia squarrosa (D,S)
Gleichenia dicarpa (D,S)
Grammitis billardierei (D,S)
G. poeppigiana (S)
Histiopteris incisa (D,S)
Hymenophyllum armstrongii (D)
H. flabellatum (D,S)
H. lyallii (D)
H. malinii (D)
H. multifidum (D,S)
H. tarum (D,S)
H. rufescens (D)
H. villosum (D,S)
Hypolepis rufobarbata (D,S)
Leptopteris hymenophylloides (D,S)
L. superba (D,S)
Poa scaberrula (D,S)
Polystichum vestitum (D,S)
Schizaea australis (D,S)

Sticherus cunninghamii (D)
S. flabellatus (Given 1982) (S)
Trichomanes strictum (D,S)

ORCHIDS

Acanthostachys viridis (Townson 1907, as *Townsonia deflexa*) (D,S)
Apotostylis bifolia (D,S)
Caladenia catnea agg. (Townson 1907, as *C. minor*)
Calochilus paludosus (Townson 1907) (D)

Chiloglottis cornuta (D)
Gastrolechia cunninghamii (Townson 1907) (D)
Hyperanthus antarcticus (D,S)
Microtis unifolia (Townson 1907, as *M. portifolia*) (D)
Orthoceras novae-zealandiae (S)
Prasophyllum colensoi (D,S)
Pterostylis banksii (Townson 1907) (D)
P. itsoniana (D)
P. venosa (D,S)
Thelymitra cyanea (D,S)
T. pulchella (D,S)

GRASSES

Agrostis dyeri agg. (D)
A. personata (D,S)
Chionochloa australis (D,S)
C. juncea (D,S)
C. rubra ssp. *occulta* (D,S)
C. sp. (unnamed) (*C. flavescens* avct. S.I., non *C. flavescens* Zotov) (D,S)
C. rubra × *C. sp. (unnamed)* (D,S)

Cortaderia richardii (D,S)
Deyeuxia aucklandica (D,S)
Davenoides (incl. *D. a.* var. *brachyantha*) (D,S)
Hierochloe novae-zealandiae (D,S)

Microlaena avenacea (D,S)
M. colensoi ss.
M. thomsonii (D,S)
Poa colensoi (S)
Rytidosperma gracile (D,S)
R. nigricans (D,S)
R. setifolium (D,S)

SEDGES

- Baumea rubiginosa* (S)
B. taratifolia (D)
Carex acicularis (S)
C. cockayneana (Townson 1907) (D)
C. coriacea (D)
C. dallii (Petrie 1914, as *C. gibbsii*) (D)
C. libera (S)
Carpha alpina s.s. (S)
C. sp. (unnamed) (plants "taller and more slender" than in *C. alpina*, with "narrower lvs"; see Edgar 1970, p. 200) (D, S)
Gahnia procera (D, S)
G. rigida (D, S)
Isolepis aucklandica (D, S)
I. subtilissima (D)
Lepidosperma australe (D, S)
Oreobolus impar (Townson 1907, Cheeseman 1925, and Mason and Moar, 1955, as *O. pumilio*) (D, S)
O. pectinatus (D, S)
O. strictus (D, S)
Schoenus pauciflorus s.s. ("slender-form" of Edgar 1970, p. 194) (D, S)
Tetrapia capillaris (S)
Uncinia filiformis (D, S)
U. involuta (S)
U. nervosa s.s. (S)
U. rupestris (incl. *U. angustifolia*) (D)

RUSHES AND ALLIED PLANTS

- Centrolepis ciliata* (D)
Empodisma minus (D, S)
Grimardia setacea (D, S)
Juncus antarcticus (D, S)
J. planifolius (D, S)
J. pusillus (S)
Luzula colensoi (S)
L. sp. (unnamed; included in *L. picta*, as var. *limosa*, by Edgar 1966) (D)

MONOCOT HERBS OTHER THAN ORCHIDS, GRASSES, SEDGES, RUSHES

- Astelia lineatis* var. *lineatis* (D, S)
A. subulata (D, S)
A. sp. (unnamed; aff. *A. nervosa*) (lvs broad; not conspicuously white adaxially as in *A. nervosa*) (D, S)
Bulbinella modesta (S)
Dianella nigra (D, S)
Herpolirion novae-zelandiae (D, S)
Libertia pulchella (D)
Luzuriaga parviflora (D, S)
Phormium cookianum (D, S)

COMPOSITE HERBS

- Anaphalis* sp. (*Hedychium bellidifoloides* s.s.) (D)
Abrotanella lineatis ss (D, S)
Brachyglottis sp. (unnamed) (D, S)

Celmisia alpina s.s. (S)

C. dallii (Townson 1907, Chaceaman 1925) (D,S)

C. incana (incl. *C. discolor*) (S)

C. lateralis (incl. *C.l.* var. *villosa*) (D,S)

C. montoi ssp. (*C. dubia*) (D,S)

C. parva (D)

C. similis (D,S)

C. sp. (i) (unnamed; aff. *C. gracilenta*)
(plants tufted; lvs linear to linear-lanceolate; upper surface and margins thinly covered in tomentum; lower surface glabrous, except for midrib) (D)

C. sp. (ii) (unnamed; aff. *C. gracilenta*) (Forstera mackayi) (D,S)

(plants strongly rhizomatous) (Gentiana gracillifolia (incl. *G. vernicosa*) (S))

C. montoi x *C. similis* (D)

Craspedia sp. (unnamed) (S)

Gnaphalium audax (D)

Leptinella sp. (D)

Raoulia eximia (Petrie 1914) (S)

R. glabra (D)

R. grandiflora var. *grandiflora* (D,S)

R. tenuicaulis (incl. *R.t.* var. *dimorpha* and *R.t.* var. *pusilla*) (D)

DICOT HERBS OTHER THAN COMPOSITES

Actinotis novae-zelandiae (D,S)

Anisotome aromatic (D,S)

A. haastii (Townson 1907, as *Ligusticum haastii*) (D)

Centella uniflora (D)

Donatia novae-zelandiae (D,S)

Drosera arcturi (D,S)

D. binata (D)

D. spathulata (D,S)

D. stenopetala (D,S)

Epilobium brunnescens s.s. (D)

E. glabellum s.s. (D)

E. linnaeoides (Townson 1907) (D)

E. pernitens (D,S)

Euphtasia townsonii s.s. (D,S)

E. weltsteiniana (D)

Forstera mackayi (D,S)

Gentiana gracillifolia (incl. *G. vernicosa*) (S)

G. montana (incl. *G.m.* var. *stolonifera*, *G. bellidifolia* var. *pulchella*,

and *G. patula*) (Allan 1961, as *G. montana* var. *stolonifera* - type loc. Mt Frederick) (D,S)

G. spenceri (Townson 1907, Chaceaman 1925) (D,S)

G. townsonii s.s. (D,S)

G. sp. (i) (aff. *G. bellidifolia*) (S)

G. sp. (ii) (unnamed; aff. *G. montana*) (plants perennial,

forming small to large clumps; stems tall, robust, often decumbent at base; basal lvs densely rosulate, in adult plants usually
(cont. on p. 7)

placed some distance up the stem, i.e. well above ground level; blade usually apiculate; petiole broad, shorter than the blade; caulinne lf pairs(2)-4-10; infl umbelliform) (D,S)

Gonocarpus micranthus ssp.
micranthus (D)

Hydrocotyle microphylla (D)

Hypsela rivularis (D)

Jovellana repens (Townson 1907,
as *Calceolaria repens*; Cheeseman
1925) (D)

Kelleria dieffenbachiae (D,S)

K. laxa (D)

Liparophyllum gunnii (D,S)

Mitrasacme montana var.
halmsii (D,S)

M. novae-zelandiae (D,S)

Nertera ciliata (D)

N. depressa (incl. *N. cunninghamii*) (D)

Oreostylidium subulatum (S)

Oxalis lactea ssp. *lactea*
(Townson 1907, as *O. macrophylla*) (D)

O. macrocarpa ssp. *calycina* (D)

Phyllachne clavigera (D,S)

Plantago triandra ss.
(D)

Pratia angulata agg. (D)

Utricularia novae-zelandiae
(D,S)

No of species: 197.

8

DOUBTFUL AND ERRONEOUS RECORDS

<u>Record</u>	<u>Probable identity of plant</u>
<i>Agrostis parviflora</i> (Townson 1907, Cheeseman 1925)	<i>Agrostis</i> sp. (unnamed)
<i>Anisotome flexuosa</i> (Dawson 1961)	
<i>Astelia nervosa</i> (Mason and Moar, 1955, as <i>A. cochaynei</i>)	<i>Astelia</i> sp. (unnamed)
<i>Brachyglottis bellidioides</i> (Town- son 1907, and Mason and Moar, 1955, as <i>Senecio bellidioides</i>)	<i>Brachyglottis</i> sp. (unnamed)
<i>Calmisia graminifolia</i> (Mason and Moar 1955)	<i>Calmisia</i> sp.(i) or <i>C.</i> sp.(ii)
<i>C. laticifolia</i> (Townson 1907; and Mason and Moar, 1955)	<i>C. similis</i>
<i>C. sessiliflora</i> (Mason & Moar, 1955)	<i>C. similis</i>
<i>Chionochloa rigida</i> (Townson 1907, as <i>Danthonia raoulii</i>)	<i>Chionochloa juncea</i> ? or <i>C. rubra</i>
<i>C. teratifolia</i> (Mason & Moar, 1955, as <i>Danthonia rigida</i> var. <i>teratifolia</i>)	<i>Chionochloa juncea</i>
<i>Coprosma</i> aff. <i>C. crenulata</i> (Mason and Moar, 1955, as <i>C. aff. retusa</i>)	<i>Coprosma</i> sp. (i) (unnamed)
<i>Dracophyllum pronum</i> (Oliver 1929)	<i>Dracophyllum politum</i> [recorded by Oliver 1952, as <i>D. densum</i>]
<i>D. lessonianum</i> (Cochayne 1928, p. 281)	<i>D. longifolium</i>
<i>D. rosmatinifolium</i> (Townson 1907)	<i>D. politum</i>
<i>D. strictum</i> (Oliver, 1929 and 1952)	
<i>D. latifolium</i> (Cochayne 1928, pp. 181, 262)	<i>D. traversii</i>
<i>Epilobium pedunculare</i> (Mason and Moar, 1955)	<i>Epilobium brunnescens</i>
<i>Euphrasia revoluta</i> (Townson 1907)	<i>Euphrasia townsonii</i>
<i>Forsteria sedifolia</i> (Townson 1907)	<i>Forsteria mackayi</i>
<i>Hebe odora</i> (Townson 1907, as <i>Veronica buxifolia</i>)	<i>Hebe</i> sp. (<i>Leonohebe mooreae</i>)
<i>Pimelea traversii</i> (Cheeseman 1925)	<i>Pimelea gnidia</i>
<i>Pseudopanax colensoi</i> (Moar and Mason, 1955, as <i>Nothopanax colensoi</i>)	<i>Pseudopanax</i> sp. (unnamed)

<i>Raoulia mamillaris</i> (Townson 1907) Tetrachondra sp. (Moar and Mason, 1955)	<i>Raoulia eximia</i> (see Petrie 1914) <i>Mitrasacme montana</i> var. <i>holmsii</i>
--	---

REFERENCES

- Allan, H. H. 1961: "Flora of New Zealand." Government Printer, Wellington. Vol. 1. 1085 pp.
- Cheeseman, T. F. 1925: "Manual of the New Zealand Flora". Government Printer, Wellington. Ed. 2 (Ed. W. R. B. Oliver). 1163 pp.
- Cockayne, L. 1928: "The Vegetation of New Zealand". 2nd ed. Englemann, Leipzig. 456 pp.
- Dawson, J. W. 1961: A revision of the genus Anisotome (Umbelliferae). Univ. Calif. Publs Bot. 33: 1-98.
- Edgar, E. 1966: Huzula in New Zealand. N.Z.J. Bot. 4: 159-84.
- 1970: In "Flora of New Zealand." Government Printer, Wellington. Vol. 2. 354 pp.
- Franklin, D. A. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans. R. Soc. N.Z. Bot. 1: 155-73.
- Given, D. R. 1982: Records of Sticherus flabellatus (R. Br.) H. S. John (Pteridophyta-Gleicheniaceae) from South I., N.Z. N.Z.J. Bot. 20: 381-5.
- Mason, R.; Moar, N. T. 1955: Notes on the vegetation and the flora of Mt Augustus, Buller County. N.Z. J. Sci. Tech. 37A: 175-86.
- Oliver, W. R. B. 1929: A revision of the genus Dracophyllum. Trans. N.Z. Inst. 59: 678-714.
- 1935: The genus Ceprosma. Bishop Mus. Bull. 132. 207 pp.
- 1952: A revision of the genus Dracophyllum: supplement. Trans. Roy. Soc. N.Z. 80: 1-17.
- Petrie, D. 1914: Some additions to the flora of the Westport district. Trans. N.Z. Inst. 46: 30-1.

Townson, W. 1907: On the vegetation of the Westport district. Trans.
N.Z. Inst. 37: 380-433.

Wardle, P. 1968: The taxonomy and distribution of the stipitate
species of Pseudopanax in New Zealand. N.Z. J. Bot. 6: 226-36.
