

HIGHER PLANTS (LYCOPDS, FERNS, GYMNOSPERMS,  
FLOWERING PLANTS) OF MT BENSON<sup>1</sup>, LOCKETT RA., IRON HILL<sup>2</sup>, AND  
DIAMOND LAKES AREA, N.W. NELSON, 3500-5560 FT/1060-1690 M

A.P.Druce, Botany Division, D.S.I.R., May 1982

(Revised Sept 1982, March 1983, June 1983, July 1983

(Based on visits 1958, 1964, 1974, Jan 1982,

March 1982, Feb 1985)

(Additions by P.A.Williams 1985)


<sup>1</sup> Grid ref. S13 988545, alt. c. 5450 ft/1660m.

→ <sup>2</sup> Grid ref. S13 019515, alt. 5561 ft/1690m

Rocks: siltstone, argillite, sandstone, conglomerate;  
volcanics;  
limestone lenses; //  
alluvium (vicinity of Diamond L.)

River  
Waingaro

June 1984  
Nov 1984  
May 1985  
July 1985  
Apr. 1986  
Oct. 1987  
Dec. 1987  
June 1988  
Sept 1990  
June 1991  
Aug 1992  
Aug 1993)



INDIGENOUS SPECIES

(unc) - uncommon or local

GYMNOSPERM TREES AND SHRUBS*Halocarpus bidwillii* *Aristotelia fruticosa* var.*Lepidothamnus laxifolius**L.intermedius* x *L.laxifolius*  
(unc)*Phyllocladus alpinus* agg.*Podocarpus nivalis**P.hallii* x *P.nivalis* (unc)*Brachyglottis adamsii**B.bidwillii* var. *viridis**B.rotundifolia* var. (incl. *B.buchananii* & *B.cockaynei*) (unc)*Cassinia laevis**Coprosma atropurpurea**C.cheesemanii* ss.*C.ciliata* ssp. *ciliata**C.colensoi* ss.*C.depressa**C.feetissima**C.petiolaris* ssp. *petiolaris**C.propinqua* ssp. *propinqua*DICOT TREES*Dracophyllum traversii* (incl. *D.pyramidalis*)*Griselinia littoralis**Hohertia lyallii* var. (*H.glabrata*)  
(unc)*Nothofagus menziesii**N.solandri* var. *cliffortioides**Pittosporum patulum* (juv. only)  
(unc)*Pseudopanax linearis**P.sinuifolium* (incl. *P.s.var.sinclairii*) (unc)*P.sp.* (unnamed; included in *P.colensoi*, as var. *floraeensis*and as var. *ternatus*, by  
Wardle (1968) (unc)*C.pseudocuneata* ssp.*C.tigida* (Diamond L. - unc)*C.tugosa* ssp. *tugosa**C.setulata* (Lockett Ra. - unc)*C.sp.(i)* (unnamed; aff. *C.pseudocuneata*) (low-growing shrubs  
with small, thick, arcuate vs)*C.sp.(ii)* (unnamed) (*C.parviflora*  
var. *dumosa* sensu Allan 1961,  
non *C.p.* var. *dumosa* Cheeseman)DICOT SHRUBS*Archeria traversii* (incl. *A.t.* var.  
*australis*) (unc)

- C. sp. (iii) (unnamed; aff. *C. colensoi*)*  
 (low-growing shrubs with trailing branches) (unc)
- C. sp. (iv) (unnamed)* (prostrate shrubs; lvs plano-convex, linear; drupes orange) (Mt Benson and Iron Hill)
- C. sp. (v) (unnamed; included in *C. parviflora* by Oliver, 1935, and others)* (lf undersurface not minutely hairy as is usual in *C. parviflora*; drupes vs. unpigmented, occ. purplish red)
- Coriaria plumosa* (unc)
- Cyathodes dealbata* (incl. *C. pumila*)
- C. sempervirens*
- Dracophyllum longifolium* var. (incl. *D. filifolium* and *D. oliveri*)
- H. ciliolata*
- H. cockayneana* var.
- H. epactidea* var. *epactidea*  
 (Iron Hill - unc)
- H. haastii* var. (aff. var. *humilis*;  
 see Moore 1961, p. 940)
- H. macrantha* var. *brachyphylla*
- H. Odora* ss.
- H. tetragona* var. (*H. coarctata*)
- H. topiaria*
- H. sp.* (*Leonochebe masoniae*)  
 (incl. *L. m.* var. *rotundata*)
- Helichrysum intermedium* var.
- Leptospermum scoparium* (unc)
- D. pubescens* var. *pubescens*
- D. uniflorum*
- D. sp. (unnamed; aff. *D. kitkui*)* (low-growing shrubs; branches often rooting; infl *Mysine divaricata* ss. of few-flld racemes) (Mt Benson)
- M. nummularia*
- Epacris alpina*
- Exocarpus bidwillii*
- Gaultheria crassa*
- G. depressa* ss. (unc)
- > *G. sp. (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962)*
- Hebe albicans* var. *albicans*
- H. canterburniensis*
- G. rupestris* (incl. *G. subcordifolia*) (unc)
- Leucopogon fraseri* ss.
- L. suaveolens* (incl. *Cyathodes colensoi*)
- Olearia colensoi* var. *colensoi* (unc)
- O. lacunosa* var. *lacunosa*
- O. nummulariifolia* var. (lvs distinctly longer than broad, rather than suborbicular as in var. *nummulariifolia*)
- O. avicanniifolia* X *O. nummulariifolia* (unc)

*Pentachondra pumila*  
*Peraxilla tetrapetala* (unc)  
*Pimelea oreophila*  
*Pittosporum anomalum*  
*P. rigidum* var. (*P. crassicaule*)  
*Traversia baccharoides* (unc)

**[DICOT LIANES AND RELATED / TRAILING PLANTS]**

*Clematis forsteri*

*Muehlenbeckia axillaris*  
*M. australis* X *M. axillaris* (unc)  
*Rubus parvus* (unc)

*Cyathea colensoi*  
*Cystopteris tasmanica*  
*Grammitis billardierei*  
*G. givenii* (unc)  
*G. magellanica* ssp. *nothofageti*  
*G. patagonica* (unc)  
*G. poeppigiana*  
*Histiopteris incisa* (unc)  
*Hymenophyllum multifidum*  
*H. peltatum* (unc)  
*H. villosum*  
*Hypolepis millefolium*  
*Ophioglossum coriaceum* agg. (unc)  
*Polystichum cystostegia* (unc)  
*P. vestitum*

**[ORCHIDS]**

**[LYCOPODS]**  
*Lycopodium austalianum*  
*L. fastigiatum*  
*L. scariosum*  
*L. vatum* (incl. *L. billardieri* and  
*L. novae-zelandicum*) (unc)

*Adenochilus gracilis*  
*Aporostylis bifolia*  
*Caladenia lyallii*  
*Chiloglottis cornuta*  
*Corybas macranthus*  
*C. trilobus* s.s.  
*Microlis oligantha* (unc)  
*Prasophyllum colensoi*  
*Thelymitra formosa* (unc)  
*T. longifolia* (unc)

**[FERNS]**

*Asplenium richardii* (unc)  
*A. terrestris* ssp. *terrestris* (unc)  
*Blechnum fluitatile* agg.  
*B. penna-marina*  
*B. procerum*  
*B. sp. (unnamed)* (mountain sp.;  
 lower pinnae not reduced) ("The  
 Dunedin-Cass Variety" of Crookes 1963)

**[GRASSES]**

*Agrostis dyeri* s.s.  
*A. muelleriana*  
*A. personata*  
*A. sp.* (unnamed; aff. *A. dyeri*)  
 (panicle not contracted as in *A.*

- Chionochloa australis*  
*C. conspicua* ssp. *conspicua*  
*C. pallens* ssp. *pallens*
- C. rubra* ssp. *occulta*  
*C. sp. (unnamed)* (*C. flavescent*  
 avct. South Island, non *C. flav*)  
*Deschampsia novae-zelandiae*  
*D. tenella*  
*Deyeuxia aucklandica*  
*D. avenoides* (incl. *D. a. var. brachyantha*)  
*D. youngii* (L. Sylvester - unc)  
*Elymus* sp. (aff. *E. rectisetus*)  
 (plants glaucous)
- Festuca* sp. (i) (unnamed; aff. *F. rubra*)  
*F. sp. (ii)* (aff. *F. matthewsii*) (tufted, but not  
 tussock-forming, glaucous)
- Hierochloe novae-zelandiae*  
*H. recurvata* (unc)
- Koeleria cheesemanii*  
*K. sp.* (*K. novo-zelandica* aff.)  
 (plants slender, rhizomatous)
- Lachnagrostis lyallii* s.s. (incl. *L. filiformis* var. *semiglabra*)
- L. sp. (i)* (unnamed) (glumes up to 4 mm  
 long; awns short or absent) (unc)
- L. sp. (ii)* (*L. Sylvester*) (unc)
- Microlaena colensoi* s.s.  
*Poa breviglumis*  
*P. celsa*
- P. colensoi*  
*P. dipsacea* (unc)  
*P. imbecilla*  
*P. kirkii*  
*P. maidis*  
*P. novae-zelandiae* s.s.  
*P. sp. (unnamed; aff. *P. novae-zelandiae*)*
- Rytidosperma buchananii* (unc)  
*R. gracile*  
*R. nigricans*  
*R. setifolium*  
*Simplicia buchananii* (unc)  
*Trisetum youngii*  
*T. sp. (i)* (unnamed; included in *T. antarcticum*, as ssp. *tenellum*,  
 by Petrie 1912)  
*T. sp. (ii)* (unnamed) (plants slender;  
 lf sheath usually glabrous; lf  
 blade flat, up to 2.5 mm wide, margins  
 smooth vs. with a few hairs near  
 ligule)
- SEDGES**
- Carex acicularis*  
*C. breviculmis* (unc)  
*C. echinata*  
*C. flavidiformis*  
*C. gaudichaudiana*

*C. sp.(i)(unnamed; aff. C. coriacea)  
(lvs slender, bright green)*

*C. libera* (unc)

*C. pyrenaica* var. *cephalotes*

*C. sinclairii*

*C. wakatipu* s.s.

*C. sp.(ii)(unnamed; aff. C. testacea)* (unc)

*Carpha alpina* s.s.

*Eleocharis acuta*

*Gahnia procera*

→ *Oreobolus pectinatus*

*O. strictus*

*Schoenus pauciflorus* s.s. ("slender  
form" of Edgar 1970, p. 194)

{ *Isolepis aucklandica*

*I. caligenis* (unc)

*I. habra*

*I. subtilissima*

*Uncinia astonii*

*U. caespitosa*

*U. clarata*

*U. divaricata*

*U. drucei*

*U. filiformis*

*U. fuscovaginata* var. (plants glaucous)

*U. involuta*

*U. longifructus*

*U. nervosa* s.s. (unc)

*U. rubra*

*U. rupestris* (incl. *U. angustifolia*)

*U. silvestris* (incl. *U. affinis*)

*Gaimardia setacea*

*Juncus antarcticus*

*J. novae-zelandiae*

*J. pusillus* (unc)

*Luzula colensoi*

*L. migrata*

*L. picta* s.s.

*L. rufa* s.s.

*L. sp. (i)(unnamed; included in *L.  
picta*, as var. *limosa*, by Edgar  
1966)*

*L. sp. (ii)(unnamed; included in  
*L. Crinita*, as var. *petrieana*,  
by Edgar 1966)*

*L. colensoi* × *L. migrata*

*Marsippospermum gracile*

**MONOCOT HERBS OTHER THAN  
/ORCHIDS, GRASSES, SEDGES, RUSHES**

*Arthropodium candidum* (unc)

*Astelia graminea*

*A. lineatis* var. *lineatis*

*A. nervosa* s.s.

*A. nivicola* s.s. (unc)

*A. petriei*

*A. skottsbergii* (unc)

*A. sp. (i)(unnamed; included  
in *A. nivicola*, as var. *moticeae*,  
by Moore 1966)*

*A. sp. (ii)(unnamed; aff. *A. nervosa*)*

(lvs broad, not conspicuously white  
adaxially as in *A. nervosa*)

**RUSHES AND ALLIED PLANTS**

*Centrolepis ciliata*

*Empodisma minus*

*Bulbinella hookeri*

*Phormium cookianum*

*Potamogeton cheesemanii* (unc)

*P. suboblongus*

**COMPOSITE HERBS**

*Anaphalis* sp. (Helichrysum bellidoides

(S.S.)

(*Helichrysum bellidoides*, as var.

*crassa*, by

(Nordenstam 1978)

*Brachyscome radicata*

*B. sinclairii* s.s.

*Craspedia* sp. (i) (unnamed; included

in *C. lanata*, as var. *elongata*, by

Allan. 1961)

*Brachyglottis* sp. (unnamed; included

in *B. bellidoides*, as var. *crassa*, by

(Nordenstam 1978)

*C. sp. (ii)* (unnamed) (plants summer-

green, lvs green, both surfaces

densely clad in multicellular,

non-glandular hairs; lf margins

faintly to distinctly white with

cottony hairs; fls white)

*Calmisia alpina* s.s.

*C. bellidoides* (unc)

*C. gibbsii*

*C. hieracifolia* ssp.

*C. incana* (incl. *C. discolor*)

*C. laricifolia*

*C. lateralis* (incl. *C. l.* var. *villosa*)

*C. montei* ssp. (*C. coccinea* var. *lanceifolia*)

*C. rupestris* (unc)

*C. sessiliflora*

*C. similis*

*C. sinclairii* (incl. *C. allanii* + *C. diversifolia*)

*C. spectabilis* ssp.

*C. traversii*

*C. sp. (i)* (unnamed; aff. *C. gracilenta*)  
(plants strongly rhizomatous)

*C. sp. (ii)* (unnamed; aff. *C. gracilenta*)

(plants tufted; lvs linear to linear-lanceolate; lower surface densely covered in tomentum)

*C. sessiliflora* × *C. traversii* (unc)

*C. similis* × *C. spectabilis* (unc)

*C. s.* × *C. traversii* (unc)

*C. spectabilis* × *C. traversii* (unc)

*C. sp. (iii)* (unnamed) (plants summer-

green; lvs green, both surfaces

densely clad in short glandular

hairs <0.3 mm long; midrib with

or without long non-glandular hairs;

lf margins faintly white with

cottony hairs; fls white)

*C. sp. (iv)* (plants summer-green; lvs

green; fls white)

*Dolichoglossis scotzonericoides*

*Gnaphalium laterale*

*G. mackayi* s.s.

*G. paludosum* s.s.

*G. polylapis* (Ruby L. -unc)

*G. ruahinicum*

*G. traversii*

<i>Haastia sinclairii</i> s.s.	<i>A. polita</i> var.
<i>Helichrysum filicale</i>	<i>Anisotome aromatic</i>
<i>Lagenifera strangulata</i>	<i>A. deltaidea</i>
<i>Leptinella pyrathrifolia</i> agg.	<i>A. filifolia</i>
L. sp., (unnamed; included in L. <i>squalida</i> , as ssp. <i>mediana</i> , by Lloyd & Webb, 1987)	<i>A. haastii</i> (unc)
<i>Leucogenes grandiceps</i>	<i>A. pilifera</i>
<i>Microseris scapigera</i> var.	
<i>Pseudognaphalium</i> sp. (unnamed; aff. <i>P. luteoalbum</i> ) (mountain sp.; lvs mostly oblong-spathulate to narrowly oblong, rounded)	<i>Cardamine</i> sp.(i) ( <i>C. debilis</i> agg.) (cf. "Long Style" of Pritchard 1957)
<i>Raoulia apicinigra</i> s.s.	<i>C. sp.(ii)</i> (unnamed) (lvs mostly simple; blades ± reniform)
<i>R. eximia</i> var.	<i>C. sp.(iii)</i> ( <i>C. debilis</i> agg.)
<i>R. glabra</i>	<i>C. sp.(iv)</i> (unnamed) ("Scree Race" of Pritchard 1957)
<i>R. grandiflora</i> var. <i>grandiflora</i>	<i>C. sp.(v)</i> ( <i>C. debilis</i> agg.)
<i>R. tenuicaulis</i> (incl. <i>R. t.</i> var. <i>dimorpha</i> and <i>R. t.</i> var. <i>pusilla</i> )	<i>Cheesemania latisiliqua</i> (incl. <i>C. gibbsii</i> - type loc. Mt Lockett) (unc)
<i>Senecio waitavensis</i>	<i>Colobanthus acicularis</i> (Lockett Ra. and ridge NW of Iron Hill - unc)
<i>Taraxacum magellanicum</i>	<i>C. affinis</i>

#### DICOT HERBS OTHER THAN COMPOSITES

<i>Acaena anserinifolia</i>	<i>C. apetalus</i> (incl. <i>C. a.</i> var. <i>alpinus</i> )
<i>A. fissistipula</i>	<i>C. canaliculatus</i>
<i>A. inaequata</i>	<i>C. strictus</i>
<i>A. profundiflora</i>	<i>C. wallii</i> (incl. <i>C. masoniae</i> )
<i>A. saccaticupula</i> (unc)	<i>C. sp. (i)</i> (unnamed) (lvs and sepals strongly acicular; lvs unbordered; sepals 5, broader than lvs, without a thickened midrib) (ridge NW of Iron Hill - unc)
<i>Aciphylla anomala</i>	
<i>A. colensoi</i> var. <i>colensoi</i>	

- C. sp. (ii) (unnamed; aff. *C. buchananii*) (Iron Hill - unc)*
- Crassula sinclairii (unc)*
- Donatia novae-zelandiae*
- Drosera arcturi*
- D. spathulata*
- D. stenopetala*
- Epilobium atriplicifolium s.s.*
- E. brunneascens s.s.*
- E. chorifolium*
- E. cockayneum*
- E. glabellum s.s.*
- E. gracilipes*
- Ekomarovianum*
- E. innacordes*
- E. macropus*
- E. pernitens*
- E. porphyrium (Mt Benson - unc)*
- E. pubens (unc)*
- E. pyrenostachyum*
- E. rubromarginatum*
- E. tasmanicum (Lockett Ra. - unc)*
- E. sp. (i) (cf. *E. matthensis* & *E. vernicosum*)*
- E. sp. (ii) (unnamed; included in *E. brunneascens*, as ssp. *minutiflorum*, by Raven and Engelhorn, 1971)*
- E. sp. (iii) (unnamed; aff. *E. glabellum*)*
- Euphorbia cheesemanii (unc)*
- E. townsonii s.s.*
- E. sp. (i) (unnamed; aff. *E. laingii*)*
- E. sp. (ii) (unnamed; aff. *E. townsonii*)*
- E. sp. (iii) (unnamed; aff. *E. townsonii*)*
- E. sp. (iv) (unnamed; aff. *E. patriei*)*
- Forstera mackayi*
- F. sedifolia s.s.*
- F. tenella*
- Galium perpusillum s.s.*
- Gentiana bellidifolia var.*
- G. sp. (i) (unnamed; aff. *G. corymbifera*)*
- G. filipes*
- G. gracilifolia (incl. *G. vernicosa*)*
- G. montana (incl. *G. m. var. stolonifera*, *G. patula*, and *G. bellidifolia* var. *pulchella*)*
- G. tenuifolia (near Ruby L. - unc)*
- G. sp. (ii) (unnamed; aff. *G. bellidifolia*)*  
(plants forming small to large patches; lvs very glossy; fls large, creamy white)
- Geranium sessiliflorum var. *nz* (unc)*
- G. sp. (unnamed; aff. *G. microphyllum*)*
- Geum leiospermum*
- G. sp. (*G. parviflorum* agg.) (stock short; terminal leaflet about as long as broad; fls up to 13 mm diam.)*
- Gigliola decipiens*
- Gonocarpus aggregatus*
- G. micranthus ssp. *micranthus**
- Hydrocotyle microphylla*
- H. sulcata*
- H. sp. (unnamed; included in *H. novae-zelandiae*, as *H. novae-zelandiae*, var. *montana*, by Kirk 1899)*
- Kelleria dieffenbachii (unc)*
- K. laxa (unc)*
- K. multiflora*

*Lilaeopsis ruthiana* (unc)

*O. lactea* ssp. *lactea*

*Limosella lineata* agg. (unc)

*Lobelia roughii* (Lockhart Ra.-unc)

*Mentha cunninghamii*

*O. simpsonii*

*Montia fontana* s.s.

*Oxalis magellanica*

*Myosotis angustatae* (Iron Hill -unc)

*Parahebe cheesemanii* (unc)

?*M. macrantha* (unc)

*P. decora*

*M. tenericaulis* (unc)

*P. lyallii*

*M. sp. (i)* (*M. australis* agg.) (fls yellow)

*Phyllachne clavigata* (incl. *P. colensoi*)

*M. sp. (ii)* (*M. australis* agg.) (fls small, white)

*Plantago lanigera* s.l. (incl. *P. novae-zelandiae*)

*M. sp. (iii)* (unnamed; included in *M. pygmaea*, as *værdtveizi*, by Moore 1961)

*P. triandra* s.s.  
*P. uniflora*

*M. forsterti*

*Pratia angulata* agg.

*Myriophyllum pedunculatum* ssp. n.3

*P. mactodon* s.s.

*M. triphyllum*

?*Ranunculus altus*

*Neopaxia* sp. (incl. *Montia calycina*)

*R. cheesemanii* var. *cheesemanii*

?*Claytonia australasica* var. *biflora*)

*R. foliosus*

*Nertera balfouriana*

*R. glabrifolius*

*N. ciliata*

*R. insignis*

*Notothlaspi* sp. (unnamed) (*N. australis* auct., non *Thlaspi australe* Hook.f.)

*R. membranifolius*

*Oreomyrrhis colensoi* s.s. (incl.

*R. verticillatus*

*O. c. var. hispida* and *O. c. var. multifida*)

*Psychotria novae-zelandiae*

*O. sp.* (unnamed; aff. *O. ramosa*)

*Oreostylidium subulatum*

*Ottelia caespitosa* ssp. *caespitosa*

OSA

*Schizilema colensoi* (unc)

*S. haastii* s.s.

*S. nitens*

*Scleranthus uniflorus*

*Stackhousia minima*

*Stellaria decipiens* (incl. *S. parviflora*)

*S. gracilenta*

*Urtica incisa*

*Viola cunninghamii*

*V. filicaulis*

*V. lyallii*

*Wahlenbergia albomarginata* var. *albomarginata* (incl. *W. pygmaea* var. *laxa* & *W. brockiei*)

### Record not accepted

*Acaena glabra* (Mason 1949)

### ADVENTIVE SPECIES

#### GRASSES

*Agrostis capillaris*

*Anthoxanthum odoratum*

*Dactylis glomerata*

*Festuca nigrescens*

*Holcus lanatus*

*Poa annua*

*P. pratensis*

#### SEDGES

*Carex ovalis*

#### RUSHES

*Juncus articulatus*

*J. bufonius*

*J. conglomeratus*

*J. effusus*

*Senecio jacobaea*

*Taraxacum officinale* s.l.

#### DICOT HERBS OTHER THAN COMPOSITES

*Achillea millefolium*

*Ceratium fontanum* esp. *vulgare*

*Linum catharticum*

*Lotus pedunculatus*

*Prunella vulgaris*

*Rumex acetosella*

*R. pulcher*

*Trifolium repens*

*Veronica arvensis*

#### COMPOSITE HERBS

*Cirsium palustre*

*C. vulgare*

*Crepis capillaris*

*Hypochaeris radicata*

*Mycalis muralis*

No. of indigenous spp. : 401.

No. of adventive spp. : 28.

## REFERENCES

- Allan, H.H. 1961: "Flora of N.Z." Govt. Printer, Wellington. Vol. 1. 1085 pp.
- Crookes, M. 1963. "N.Z. Ferns". Ed. 6. Whitcombe & Tombs. 408 pp.
- Cheseman, T.F. 1906: "Manual of the N.Z. Flora". Govt. Printer, Wgtn. 1199 pp.  
1925: "Manual of the N.Z. Flora". Govt. Printer, Wgtn. Ed. 2. 1163 pp.
- Edgar, E. 1966: Luzula in N.Z. N.Z. Jl Bot. 4: 159-84.  
— 1970: Cyperaceae Sp. 167-285. In "Flora of NZ" Govt. Printer, Wgtn. Vol 2. 354 pp.
- Franklin, D. 1962: The Ericacea in N.Z. (Gaultheria and Pernetta).  
Trans. R. Soc. N.Z. Bot. 1: 155-73.
- Hackel, E. 1903: On new species of grasses in N.Z. Trans N.Z. Inst 35: 377-85.
- Kirk, T. 1899: "The Student's Flora of N.Z." Govt. Printer, (Wellington. 408 pp.
- Moore, L.B. 1961: In "Flora of New Zealand". Government Printer, Wgtn. Vol 1. 1085 pp.  
— 1966: Australasian asteliads (Liliaceae). N.Z. Jl Bot. 4: 201-40.
- Nordenstam, B. 1978: Opera Bot. 44: 1-84.
- Oliver, W.R.B. 1935: The genus Coptosma Bull. Bishop Mus. Monographs 132. 207 pp.
- Petrie, D. 1912: Description of new....species.... Trans. N.Z. Inst. 44: 177-87.
- Pritchard, G.G. 1957: Experimental taxonomic studies on species of Ceratamine Linn. in N.Z. Trans. R. Soc. N.Z. 85: 75-89.
- Raven, P.H.; Engelkorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. Jl Bot. 9: 345-50.
- Wardle, P. 1968: The taxonomy and distribution of the stipulate species of Paxifloraceae in N.Z. N.Z. Jl Bot. 6: 226-36.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the N.Z. region  
Rec. Dom. Mus. (Wellington) 5: 101-46.
- Lloyd, D.G.; Webb, C.J. 1987: N.Z. Jl Bot. 25: 99-105. Re reinstatement  
of heptmella at generic rank, and the status of the 'Cotuloid'  
(Asteraceae, Anthemidaceae). N.Z. Jl Bot. 25: 99-105.
- Mason, R. 1949: Station records of indigenous plants. Trans. R. Soc. N.Z. 77(5): 175-80.