

INDIGENOUS HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING PLANTS) OF MT DOMETT¹, MT CENTRE² AND LAKE AORERE³ AREA, DOMETT RA., N.W. NELSON, 2800-5325 FT (850-1620 M)

A.P. Druce, Botany Division, D.S.I.R., May 1982

(revised Sept 1982, June 1983, July 1986, Oct. 1987)

(Based on visit Jan. 1977)

July 1990
May 1991,

Aug. 1992


Aug. 1993

¹ Grid ref. S12 746563, alt. c. 5325 ft

² Grid ref. S12 781561, alt. c. 5000 ft

→ ³ Grid ref. S12 765570, alt. c. 3350 ft

Rocks: granite



(unc) - uncommon or local

GYMNOSPERM TREES AND SHRUBS

Halocarpus biformis
Lepidothamnus intermedius
L. laxifolius
Libocedrus bidwillii
Phyllocladus alpinus agg.

Podocarpus acutifolius
P. nivalis

Quintinia serrata (incl. *Q. acutifolia* and *Q. elliptica*)
Weinmannia racemosa var.
racemosa

DICOT SHRUBS

Archeria traversii (incl. *A.t. var. australis*)
Aristotelia fruticosa var.

MONOCOT TREES AND SHRUBS

Cordyline indivisa

DICOT TREES

Aristotelia serrata
Dracophyllum traversii (incl. *D. pyramidale*)
Elaeocarpus hookerianus
Fuchsia excorticata
Grevillea littoralis
Hoheria lyallii var. (*H. glabrata*)
Metrosideros umbellata
Nothofagus fusca
N. menziesii
Pseudopanax linearis
P. simplex (incl. *P.s. var. sinclairii*)
P. sp. (unnamed; included in *P. colensoi*, as var. *fiordensis* and as var. *tornatus*, by Wardle 1968)

Brachyglottis adamsii
B. bidwillii var. *nitidis*
Brotundifolia var. (incl. *B. buchananii* & *B. cockaynei*)
B. bidwillii X *B. rotundifolia* (unc)
Cassinia auvilliersii
Coprosma cheesemanii s.s.
C. ciliata ssp. *ciliata*
C. colensoi s.s.
C. crenulata
C. depressa
C. foetidissima
C. perpusilla ssp. *perpusilla*
C. pseudocuneata ssp.
C. rhamnoides (incl. *C. polymorpha*)
C. rugosa ssp. *rugosa*
C. serrulata

- C.sp.(i)*(unnamed; aff. *C.pseudocuneata*)(low-growing shrubs with small, thick, arcuate vs)
C.sp.(ii)(unnamed; aff. *C.colensoi*)(low-growing shrubs with trailing branches)
C.sp.(iii)(unnamed) (*C.parviflora* var. *dumosa* sensu Allan 1961, non *C.p.* var. *dumosa* Cheeseman)
C.sp.(iv)(unnamed; included in *C.parviflora* by Oliver, 1935, and others)(If undersurface not minutely hairy as is usual in *C.parviflora*; diopes vs. unpigmented, occ. purplish red)
- Coriaria angustissima*
C.satmentosa
Cyathodes dealbata (incl. *C.pumila*)
C.ampellifolia
-Dracophyllum pubescens var. *pubescens*
D.longifolium var. (incl. *D.filifolium* & *D.oliveti*)
D.politum
D.uniflorum var. *uniflorum*
Epacris alpina
Exocarpus bidwillii
Gaultheria crassa
G.depressa s.s.
- G.rupestris* (incl. *G.subcorymbosa*)
G.sp.(unnamed; included in *G.depressa*, as var. *novae-zelandiae*, by Franklin 1962)
G.crassa X *G.rupestris* (unc)
G.rupestris X *G.sp.*(unnamed)(unc)
Hebe canterbutiensis
H.ciliolata
H.salicifolia var. *salicifolia*
H.topiaria
H.tetragona var. (*H.coarctata*)
H.sp.(*Leonohebe mooreae*)
Leptospermum scoparium
Leucopogon fraseri s.s.
Melicytus sp.(unnamed)
- Myrsinie divaricata* s.s.
M.nummularia
Neomyrtus pedunculata
Olearia arborescens
O.colensoi var. *colensoi*
O.ilicifolia
O.lacunosa var. *lacunosa*
O.arborescens X *O.ilicifolia*
O.capillaris X *O.ilicifolia*
O.ilicifolia X *O.lacunosa* (und)
Pentachondra pumila
Pimelea oreophila
Pittosporum rigidum var. (*P.crassicaule*)

Pseudopanax anomalus

Pseudowintera colorata

DICOT LIANES AND RELATED/

TRAILING PLANTS

Muehlenbeckia australis

M. axillaris

M. australis × *M. axillaris*

Rubus cissoides var. (leaflets broader
than in var. *cissoides*)

R. parvus

Cyathea colensoi

Grammitis billardierei

G. givenii (unc)

G. magellanica ssp. *nothofageti*

G. paeppigiana

Histiopteris incisa

Hymenophyllum armstrongii

H. demissum

H. multifidum

H. ratum

H. rufescens

H. villosum

Hypolepis millefolium

H. rufobarbata

Leptopteris superba

Ophioglossum coriaceum ass.

Phymatosorus pustulatus

Polystichum cystostegia

P. vestitum

Pteridium esculentum

Schizaea australis (unc)

LYCOPODS

Lycopodium australianum

L. fastigiatum

L. laterale (incl. *L. ramulosum*?)

L. scariosum

L. vatum (incl. *L. billardierei* and

L. novae-zelandicum)

L. volubile

FERNS

Asplenium terrestris ssp. *terrestris* (unc)

Blechnum discolor

B. fluitatile ass.

B. penna-martini

B. procatum

B. sp. (unnamed) (mountain
sp.; lower pinnae not reduced
in length) ("The Dunedin-Cass
Variety" of Crookes 1963)

ORCHIDS

Adenochilus gracilis

Aporostylis bifolia

Caladenia lyallii

Corybas trilobus s.s.

Liparanthus antarcticus

Microtis unifolia

Prasophyllum colensoi

Pterostylis hissoniana (unc)

P. venosa

Thelymitra cyanea

T. hatchii

T. longifolia

GRASSES

Agrostis oressbia

A. personata

A. sp. (unnamed; aff. *A. dyeri*)

Chinochloa australis

C. cheesemanii

C. conspicua ssp. *conspicua*

C. pallens ssp. *pallens*

Lachnagrostis lyallii ss. (incl.)

L. filiformis var. *semiglabra*)

Microlaena avenacea

M. colensoi

Poa breviglumis

P. celsa

P. cita ssp. *cita*

P. colensoi

P. Kirkii

P. maia

P. novae-zelandiae ss.

C. rubra ssp. *occulta*

C. sp. (unnamed) (*C. flavesans*

avct. South Island, non *C. flaves-*
ans (Löts.)

C. conspicua × *C. pallens* (unc)

C. c. × *C. rubra* (unc)

C. pallens × *C. rubra*

C. rubra × *C. sp.* (unnamed) (unc)

Deschampsia novae-zelandiae

D. tanella

Deyeuxia aucklandica

Festuca sp. (unnamed) (tufted but
not tussock-forming, glaucous)

Hierochloë novae-zelandiae

H. recurvata

H. redolens

Rytidosperma gracile

R. nigricans

R. setifolium

R. viride (unc)

Trisetum youngii

T. sp. (unnamed) (plant slender)

If sheath glabrous or nearly so;
If blade flat, up to 2.5 mm wide,
margins smooth, vs. with a few
hairs near ligule)

SEDGES

Carex acicularis

C. coerulea

C. dallii

C. dissita ss.

C. echinata var. (*C. stellulata*
var. *australis*)

C. pyrenaica var. *cephalotes*
C. sinclairii
C. trachycarpa
C. wakatipu s.s.
C. sp. (Section *Acutae*)
Carex alpina s.s.
Eleocharis acuta
Gahnia procera
Oreobolus impar
O. pectinatus
O. strictus
Schoenus pauciflorus s.s.
Isolepis aucklandica
I. caligenis
I. habra
I. subtilissima
Uncinia clavata
U. divaricata
U. filiformis plants
U. fuscovaginata var. *(flavous)*
U. involuta
U. longifluctus
U. rupestris (incl. *U. angustifolia*)
U. uncinata
U. vitidis
U. sp. (unnamed) (*U. affinis* sensu Hamlin 1959, non *U. riparia* var. *affinis* Col. ex Clarke)

RUSHES AND ALLIED PLANTS

Centrolepis ciliata
Empodisma minus
Gaimardia setacea
Juncus antarcticus
J. novae-zelandiae
J. pusillus
Luzula colensoi
L. migrata
L. picta s.s.
L. subclavata
L. sp. (unnamed; included in *L. picta*, as var. *limosa*, by Edgar 1966)

MONOCOT HERBS OTHER THAN ORCHIDS, GRASSES, SEDGES, RUSHES

Astelia graminea
A. linearis
A. nervosa s.s.
A. nivicola s.s.
A. petriei
A. skottsbergii
A. subulata (Mt Centre - unc)
A. sp. (unnamed; aff. *A. nervosa*)
 (lvs broad, not conspicuously white adaxially as in *A. nervosa*)
Harpolithion novae-zelandiae

Libertia peregrinans

L. pulchella

Luzula parviflora

Phormium cookianum

Potamogeton suboblongus

COMPOSITE HERBS

Abrortanella caespitosa

A. linearis s.s.

A. pusilla

Anaphalis sp. (*Helichrysum bellidioides* ss) *G. paludosum* s.s.

Brachyglottis sp. (unnamed) (*B. bellidioides* var. *crassa*)

Brachyscome radicata

Calmisia alpina s.s.

C. argentea (Mt Centre) (unc)

C. armstrongii

C. dallii

C. gibbsii

C. haastii ssp. *haastii*

Cincana (incl. *C. discolor*)

C. laticifolia

C. montoi ssp. (*C. coriacea* var. *lancifolia*)

C. parva

C. petriei

C. sessiliflora

C. spectabilis ssp.

C. travetsii

C. armstrongii x *C. spectabilis* (unc)

Leptinella pyrothrifolia agg.

L. sp. (unnamed; included in
L. squamida, as ssp. *mediana*,
by Lloyd and Webb, 1987)

Craspedia sp. (unnamed) (summer-
green; lvs elliptic-spathulate, green
both surfaces clad in hispid hairs; lf
margins faintly to distinctly
white with cottony hairs; fls
white)

Dolichoglottis scorzoneraoides

Gnaphalium laterale (?)

G. limosum s.s.

G. paludosum s.s.

Brachyglottis sp. (unnamed) (*B. bellidioides* var. *crassa*)

Pseudognaphalium sp. (unnamed; aff.
P. luteoalbum) (lvs mostly linear-
spathulate to linear, acute)

Lagenifera cuneata

L. pumila

L. strangulata

Leucogenes grandiceps

Microseris scapigera var.

Raoulia glabra

R. grandiflora var. *grandiflora*

R. tanicaulis (incl. *R. t.* var.

dimorpha and *R. t.* var. *pusilla*)

Senecio wairauensis

Taraxacum magellanicum

DICOT HERBS OTHER THAN COMPOSITE

Acaena ansetinifolia

A. profundeincisa

<i>Aciphylla colensoi</i> var. <i>colensoi</i>	<i>Diosera arcturi</i>
<i>A. hookeri</i>	<i>D. binata</i>
<i>A. polita</i> var.	<i>D. spathulata</i>
<i>A. colensoi</i> × <i>A. hookeri</i> (unc)	<i>D. stenopetala</i>
<i>A. c. × A. polita</i> (unc)	<i>Epilobium alsinoides</i> s.s.
<i>Anisotome aromatica</i>	<i>E. brunnescens</i> s.s.
<i>A. deltoides</i>	<i>E. chlorifolium</i>
<i>A. haastii</i>	<i>E. cockayneanum</i>
<i>A. sp. (i)</i> (unnamed; included in <i>A. imbricata</i> , as var. <i>prostrata</i> , by Dawson 1961)	<i>E. glabellum</i> s.s.
<i>A. sp. (ii)</i> (unnamed) (turf-forming in seepages; infls short, up to 5 cm)	<i>E. komarovianum</i>
<i>Actinotis novae-zelandiae</i>	<i>E. linnaeoides</i>
<i>Cardamine</i> sp. (i) (<i>C. debilis</i> agg.) ("Narrow Petal" of Pritchard 1957)	<i>E. heteroides</i>
<i>C. sp. (ii)</i> (<i>C. debilis</i> agg.) (cf. "Long Style" of Pritchard 1957)	<i>E. pernitens</i>
<i>C. sp. (iii)</i> (<i>C. debilis</i> agg.) ("Glossy Leaf" of Pritchard 1957)	<i>E. tasmanicum</i> (unc)
<i>C. sp. (iv)</i> (unnamed) ("Scree Race" of Pritchard 1957)	<i>E. sp. (unnamed; included in <i>E. brunnescens</i>, as ssp. <i>minutiflorum</i>, by Raven and Engelhorn 1971)</i>
<i>Cheesemania latissiliqua</i> (incl. <i>C. gibbsii</i>) (unc)	<i>Euphrasia townsonii</i> s.s.
<i>Colobanthus affinis</i>	<i>E. sp. (unnamed; aff. <i>E. petriei</i>)</i>
<i>Donatia novae-zelandiae</i>	<i>Forstera mackayi</i>
	<i>F. sedifolia</i> s.s.
	<i>F. tenella</i>
	<i>Galium perpusillum</i> s.s.
	<i>Gentiana bellidifolia</i>
	<i>G. gracilifolia</i> (incl. <i>G. vernicosa</i>)
	<i>G. montana</i> (incl. <i>G. m. var. stolonifera</i> , <i>G. patula</i> , & <i>G. bellidifolia</i> var. <i>pulchella</i>)
	<i>G. spenceti</i>
	<i>G. sp. (unnamed; aff. <i>G. bellidifolia</i>)</i> (plants forming small to large patches; stems decumbent; fls large, creamy white)

<i>Geranium</i> sp. (aff. <i>G. microphyllum</i>)	<i>Nettaria balfouriana</i>
<i>Geum uniflorum</i>	<i>N. ciliata</i>
<i>G. sp. (i)</i> (<i>G. parviflorum</i> agg.) (stock short; terminal leaflet about as long as broad; fls up to 13 mm diam.)	<i>N. depressa</i> (incl. <i>N. cunninghamii</i>)
<i>G. sp. (ii)</i> (<i>G. parviflorum</i> agg.) (stock creeping and branching; terminal leaflet slightly longer than broad; fls up to 8 mm diam.)	<i>N. sp.</i> (unnamed; aff. <i>N. dichondrifolia</i>)
<i>Gingidia montana</i> (unc)	<i>Oreomythis colensoi</i> s.s. (incl. <i>O.c. var. hispida</i> and <i>O.c. var. multifida</i>)
<i>Gonocarpus aggregatus</i>	<i>Oreostylidium subulatum</i>
<i>Gunnera dentata</i> var. <i>dentata</i>	<i>Oursia caespitosa</i> ssp. <i>caespitosa</i>
<i>Hydrocotyle dissecta</i>	<i>O. lactea</i> ssp. <i>lactea</i>
<i>H. microphylla</i>	<i>O. macrocarpa</i> ssp. <i>calycina</i>
<i>H. sulcata</i>	
<i>H. sp.</i> (unnamed; included in <i>H. novae-zelandiae</i> , as var. <i>montana</i> , by Kirk 1899)	<i>O. simpsonii</i>
<i>Kelletia dieffenbachii</i>	
<i>K. laxa</i>	
<i>K. multi-flora</i>	
<i>Lilaeopsis ruthiana</i>	<i>Oxalis magellanica</i>
<i>Liparophyllum gunnii</i>	<i>Parahelbe lyallii</i>
<i>Mitrasacme novae-zelandiae</i>	<i>Phyllachne clavigera</i>
<i>Montia fontana</i> s.s.	<i>Plantago lanigera</i> s.l.
<i>Myosotis forstenii</i>	<i>P. raoulii</i>
<i>M. lyallii</i> (incl. <i>M. lyallii</i> var. <i>townsonii</i> and <i>M. elatii</i>) (unc)	<i>P. triandra</i> s.s.
<i>M. sp.</i> (unnamed; included in <i>M. pygmaea</i> , as var. <i>drucei</i> , by Moore 1961)	<i>P. uniflora</i>
	<i>Pratia angulata</i> ass.
	<i>Psychotrichia novae-zelandiae</i>
	<i>Ranunculus altus</i>
	<i>R. foliosus</i>
<i>Myriophyllum triphyllum</i>	
<i>Neopaxia</i> sp. (incl. <i>Montia calycina</i> and <i>Claytonia australasica</i> var. <i>biflora</i>)	

R. insignis
R. membranifolius
R. reflexus
R. verticillatus

Schizolema haastii s.s.
Stellaria decipiens (incl. S. parviflora)
Urtica incisa
Viola cunninghamii
V. filicaulis
V. lyallii
Wahlenbergia albomarginata
var. albomarginata (incl. W. pygmaea var. laxa and W. brockiei)

No. of species: 352.

REFERENCES

- Allan, H.H. 1961: "Flora of N.Z." Govt. Printers, Wellington. Vol. 1. 1085 pp.
Crookes, M. 1963: "N.Z. Ferns". Ed. 6. Whitcombe & Tombs. 408 pp
- Dawson, J.W. 1961: Univ. Calif. Public Bot. 33: 1-98
Edgar, E. 1966: Luzula in N.Z. N.Z. Jl Bot. 4: 159-84.
Franklin, D. 1962: Trans. R. Soc. N.Z. Bot. 1: 155-73.
Hamlin, B.G. 1959: Dom. Mus. Bull. 19. 106 pp.
Lloyd, D.G.; Webb, C.J. 1987: NZ Jl Bot. 25: 99-105.
Moore, L.B. 1961: In "Flora of N.Z.". Govt. Printers, Wellington. Vol. 1. 1085 pp.
Oliver, W.R.B. 1935: The genus Coprosma. Bull Bishop Mus Honolulu 132. 207 pp.
Pritchard, G.G. 1957: Trans. R. Soc. N.Z. 85: 75-89.
Raven, P.H.; Engelhorn, T. 1971: NZ Jl Bot. 9: 345-50.
Wardle, P. 1968: N.Z. Jl Bot. 6: 226-36.
Zofov, V.D. 1965: Rec. Dom. Mus. Wellington 5: 101-46.