

HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING PLANTS) OF SOUTHERN HOPE RANGE^t, N W NELSON,
1500-4080 FT / 450-1244 M

A.P. Druce, Botany Division, D.S.I.R., May 1982
(revised Sept. 1982, June 1983, June 1984, April 1986, Dec. 1987,
June 1988, Oct. 1988, Aug. 1990, June 1991, Aug. 1992)

Revised and rewritten March 1993

Based on visits Nov. 1976, Nov. 1978, Feb. 1986, Dec. 1987,
Feb. 1993


(Members of 1993 field party: Tony Druce, Peter Williams,
Shannel Courtney, Kerry Ford, Denise Cutler, Arnold Heine,
Roland Heine)

^t Highest point: Mt Hope, 1244 m, grid ref. NZMS 260

Rocks: granite

NZS 770 5:0.

(map point)



(unc) - uncommon or local
 (N) - seen only at north end of range, at foot of Lookout Ra.

INDIGENOUS PLANTS

GYMNOSPERM TREES AND SHRUBS

- Dactylium cupressinum*
- Halocarpus bidwillii*
- H. biformis*
- H. bidwillii* X *H. biformis*
- Lagarostrobos colensoi* (unc)
- Lepidothamnus intermedius*
- L. laxifolius*
- L. intermedius* X *L. laxifolius*
- Libocedrus bidwillii*
- Phyllocladus* sp. (unnamed; aff. *P. alpinus*) ("lowland" ecotype of Wardle 1969)
- Podocarpus hallii*
- P. nivalis*
- P. hallii* X *P. nivalis* (unc)

C. foetidissima

- C. micocarpa*
- C. petpusilla* ssp. *perpusilla*
- C. propinqua* ssp. *propinqua* (unc)
- C. pseudocuneata* ssp. (lvs narrower than in ssp. *pseudocuneata*)
- C. thamnoides* (incl. *C. polymorpha*)
- C. sp.(i)* (unnamed) (*C. parviflora* var. *dumosa* sensu Allan 1961, non *C. p. var. dumosa* Cheesem.)
- C. sp.(ii)* (unnamed; aff. *C. pseudocuneata*) (low-growing shrubs with small, thick, arcuate lvs)
- C. sp.(iii)* (unnamed; included in *C. parviflora* by Oliver, 1935, and others) (lf undersurface not minutely hairy as is usual in *C. parviflora*; drupes usually unpigmented, occasionally purplish red)

Cyathodes dealbata (incl. *C. pumila*)

- C. empetrifolia*
- C. sp.* (*C. juniperina* agg.) (lvs small, vs. < 1 cm X < 1 mm)
- Dracophyllum longifolium* var. (incl. *D. filifolium* and *D. oliveri*)
- D. palustre*
- D. pronum* (N)
- D. traversii* (incl. *D. pyramidale*) (unc)
- D. uniflorum*
- D. longifolium* X *D. palustre* (unc)

DICOT TREES AND SHRUBS

- Archeria traversii* (incl. *A. t. var. australis*) (unc)
- Aristotelia fruticosa* var. (unc)
- Brachyglottis bidwillii* var. *viridis* (unc) (N)
- B. rotundifolia* var. (incl. *B. buchananii* and *B. cockaynei*) (unc)
- Catpodetus serratus*
- Cassinia varillletsii*
- Coprosma cheesemanii* s.s.
- C. colensoi* s.s.
- C. crenulata* (unc, except at north end of range at foot of Lookout Ra.)

<i>Elaeocarpus hookerianus</i>	<i>Nothofagus fusca</i>
<i>Epacris alpina</i>	<i>N. solandri</i> var. <i>cliffortioides</i>
<i>Exocarpus bidwillii</i>	<i>N. solandri</i> var. <i>solandri</i>
<i>Fuchsia excorticata</i>	<i>N. truncata</i>
<i>Gaultheria antipoda</i>	<i>N. fusca</i> X <i>N. solandri</i>
<i>G. crassa</i> (unc)	<i>Olearia arborescens</i> (unc)
<i>G. depressa</i> s.s.	<i>O. lacunosa</i> var. <i>lacunosa</i> (unc)
<i>G. macrostigma</i> (unc)	<i>O. virgata</i> var. <i>implicita</i> (incl. <i>O.</i> <i>virgata</i> var. <i>divaricata</i> and <i>O.</i> <i>virgata</i> var. <i>laxiflora</i>) (unc)
<i>G. sp.</i> (unnamed; included in <i>G. depressa</i> , as var. <i>nova-</i> <i>zealandiae</i> , by Franklin (1962))	<i>Pentachondra pumila</i>
<i>G. crassa</i> X <i>G. macrostigma</i> (unc)	<i>Pimelea oreophila</i> (unc)
<i>G. crassa</i> X <i>G. sp.</i> (unnamed) (unc)	<i>Pittosporum anomalum</i>
<i>G. macrostigma</i> X <i>G. sp.</i> (un- named) (unc)	<i>P. divaricatum</i>
<i>Griselinia littoralis</i>	<i>P. rigidum</i> var. (<i>P. crassicaule</i>)
<i>Hebe canterburiana</i>	<i>Pseudopanax anomalous</i>
<i>H. gracillima</i> (unc)	<i>P. colensoi</i> s.s.
<i>H. salicifolia</i> var. <i>salicifolia</i> (unc)	<i>P. crassifolius</i>
<i>H. tetragona</i> var. (<i>H. coactata</i>)	<i>P. linearis</i>
<i>H. topiaria</i>	<i>P. simplex</i> (incl. <i>P. simplex</i> var. <i>sinclairii</i>)
<i>H. sp.</i> (<i>Leionchebea masoniae</i>)	<i>P. sp.</i> (unnamed; included in <i>P.</i> <i>colensoi</i> , as var. <i>tornatus</i> , by Wardle (1968))
<i>Kunzea ericoides</i> agg. (unc)	<i>P. anomalous</i> X <i>P. simplex</i> (unc)
<i>Leptospermum scoparium</i> (from Hope scarp northwards)	<i>Pseudowintera colorata</i>
<i>Leucopogon fasciculatus</i> var. (juv. lvs not much broader than adult lvs)	<i>Weinmannia racemosa</i> var. <i>racemosa</i>
<i>L. fraseri</i> s.s.	[DICOT LIANES AND RELATED]
<i>L. suaveolens</i> (incl. <i>Cyathodes</i> <i>colensoi</i>)	[TRAILING PLANTS]
<i>Metrosideros umbellata</i>	<i>Muehlenbeckia axillaris</i> (unc)
<i>Myrsine divaricata</i> s.s.	<i>Rubus cissoides</i> var. ("form c" of Allan 1961; leaflets broader than in var. <i>cissoides</i>)
<i>M. nummularia</i>	
<i>Neomyrtus pedunculata</i>	

Lycopods

Lycopodium australianum
L.fastigiatum
L.scariosum
L.varium
L.volubile

H.flabellatum
H.malingii
H.minimum(unc)
H.multifidum
H.peltatum(unc)
H.ratum
H.villosum

FERNS

Asplenium flaccidum
Blachnum chambersii
B.colensoi
B.discolor
B.fluviatile agg.
B.minus(unc)
B.penna-marina
B.procerum
B.vulcanicum(unc)
B.sp(i)(unnamed)(lowland sp.; lower pinnae reduced in length)
B.sp.(ii)(unnamed)(mountain sp.; lower pinnae not reduced in length) ("The Dunedin-Cass Variety" of Crookes 1963)

Cyathea colensoi
Gleichenia dicarpa
Grammitis billardierei
G.givenii(unc)
G.magellanica ssp. nothofagi (unc)
G.poepigiana (unc)
Histopteris incisa(unc)
Hymenophyllum armstrongii (unc)
H.bivalve
H.demissum

Hypolepis millefolium(unc)
Polystichum vestitum(unc)
Schizaea australis(unc)
Sticherus cunninghamii (unc)

ORCHIDS

Adenochilus gracilis
Aporostylis bifolia
Caladenia carnea agg.
C.lyallii(unc)
Chiloglottis cornuta
Corybas cryptanthus(unc)
C.fivularis s.s.(unc)
C.trilobus s.s.
Gastrodia cunninghamii (unc)
Hyperanthus antarcticus(unc)
Prasophyllum colensoi
Pterostylis irsoniana(unc)
P.venosa(unc)
Thelymitra cyanea
T.decora(unc)
T.formosa(unc)
T.hatchii
T.longifolia

GRASSES

Agrostis dyeri s.s.(unc)
A.personata
A.sp.(unnamed; aff.A.dyeri)(panicle

spreading, not contracted as in <i>A. dycti</i>)	<i>C. sinclairii</i>
<i>Chionochloa australis</i>	<i>C. sp.(i)(unnamed; aff. C. dissita)</i> (plants smaller than in <i>C. dissita</i>)
<i>C.conspicua</i> ssp. <i>conspicua</i> (unc)	(unc)
<i>C.tubra</i> ssp. <i>occulta</i>	
<i>C.australis</i> X <i>C.tubra</i> (unc)	<i>C.sp(i)(unnamed; aff. C.conspicua)</i> (lvs slender, bright green)
<i>C.tubra</i> X <i>C.sp.</i> (unnamed; <i>C.</i> <i>flavescens</i> avct. South Island, non <i>C.flavescens</i> Zотов) (unc)	
<i>Deschampsia tenella</i> (unc)	<i>Carpha alpina</i> s.s.
<i>Deyeuxia aucklandica</i>	<i>Gahnia procera</i>
<i>D.avenoides</i> (incl. <i>D.a.</i> var. <i>brachyantha</i>)	<i>Isolepis aucklandica</i>
<i>Hierochloe recurvata</i> (unc)	<i>I.caligenis</i> (unc)
<i>H.rodolensi</i> (unc)	<i>I.habra</i>
<i>H.novae-zelandiae</i>	<i>I.subtilissima</i>
<i>Lachnagrostis</i> sp. (unnamed; included in <i>L.filiformis</i> , as var. <i>semiglabra</i> , by Zотов 1965)	<i>Oreobolus pectinatus</i>
<i>Microlaena avenacea</i>	<i>O.strictus</i>
<i>M.colensoi</i> (unc) (N)	<i>Schoenus pauciflorus</i> s.s. (= "slender form" of Edgat 1970, p. 194) (unc)
<i>Poa breviglumis</i> (unc)	<i>Uncinia divaricata</i>
<i>P.colensoi</i>	<i>U.filiformis</i> (unc)
<i>Rytidosperma gracile</i>	<i>U.fuscovaginata</i> var. (plants glaucous)
<i>R.nigricans</i>	<i>U.involuta</i>
<i>R.setifolium</i> (unc)	<i>U.nervosa</i> s.s.
<i>R.sp.(undetermined)</i> (unc)	<i>U.rupestris</i> (incl. <i>U.angularifolia</i>)
	<i>U.uncinata</i>
	<i>U.zotovii</i>
	<i>U.sp.(unnamed; aff. U.nervosa)</i> (plants more slender than in <i>U.nervosa</i> ; see Edgat 1970, p. 231)

SEDGES

- Carex aciculatis*
C.breviculinis (unc)
C.cockayneana (unc)
C.echinata var. (*C.stellulata*
var. *australis*)
C.gaudichaudiana
C.libera

RUSHES AND ALLIED PLANTS

- Centrolepis ciliata*
Empodisma minus
Gaimardia setacea

- Juncus antarcticus*
J. novae-zelandiae
J. pusillus
Luzula colensoi(unc)
L. picta s.s.
L. subclavata
L. sp. (unnamed; included in *L. picta*, as var. *limosa*, by Edgar 1966)

- C. Similis*
C. spectabilis ssp.
C. similis x *C. spectabilis* (unc)
Craspedia sp. (plants summer-green, in carpetgrass and bogs; lvs green, recurved; laminae folded, narrowing abruptly to long slender petiole; both surfaces (cont. on p.9))
Gnaphalium laterale (unc)
G. mackayi s.s.
G. paludosum s.s.
Helichrysum filicaule (unc)
Lagenifera cuneata
L. strangulata
Leptinella sp. (unnamed; included in *L. squalida*, as ssp *mediana*, by Lloyd and Webb, 1987) (unc)

**MONOCOT HERBS OTHER THAN
ORCHIDS, GRASSES, SEDGES, RUSHES/
AND ALLIED PLANTS**

- Astelia linearis* var. *linearis*
A. nervosa s.s.
Libertia pulchella (unc)
Luzuriaga parviflora (unc)
Phormium cookianum

- Raoulia glabra*
R. grandiflora var. *grandiflora*
R. Subsericea (unc)
Senecio minimus (unc)
S. wairauensis (unc)

COMPOSITE HERBS

- Abrotanella caespitosa*
Anaphalis sp. (*Helichrysum bellidioides* s.s.)
Brachyglottis sp. (unnamed; included in *B. bellidioides*, as var. *crassa*, by Nordenstam 1978)
Calmsia alpina s.s.
C. dallii (unc, near Mt Hope)
C. hieracifolia ssp.
C. incana (incl. *C. discolor*)
C. lateralis (incl. *C.l.* var. *villosa*) (unc, except at north end of range at foot of Lookout Ra.)
C. monroi ssp. (*C. semicordata* var. *semicordata*)
C. petrici (unc)

- DICOT HERBS OTHER THAN COMPOSITES**
- Acaena anethinifolia* (unc)
A. inermis (unc)
A. profunda (incisa) (unc)
Anisotome aromatica
Calobanthus apetalus (incl. *C. apetalus* var. *alpinus*) (unc)
Coprosma talbotiae (unc)
Donatia novae-zelandiae
Dioseta arctuti
D. spathulata
D. stenopetala (unc)

- Epilobium alsinoides* s.s. (unc)
E. brunnescens s.s. (unc)
E. cochlearneanum
E. parnitens
E. sp. (unnamed; included in *E. brunnescens*, as ssp. *minutiflorum*, by Raven and Engdorn, 1971) (unc)
Euphrasia townsonii
Forstera mackayi
F. tenella
Gentiana montana (incl. *G. in. var. stolonifera*, *G. patula*, and *G. bellidifolia* var. *pulchella*)
G. spenceri
Geranium sp. (unnamed; aff. *G. microphyllum*)
Gonocarpus aggregatus
G. micranthus ssp. *micranthus*
G. montanus (unc)
Hydrocotyle microphylla (unc)
H. moschata (unc)
Hypericum japonicum agg. (unc)
Kelleria diaphanbachii (unc)
K. laxa
K. tessellata
Liparophyllum gunnii
Montia fontana s.s.
Nepaxia sp. (incl. *Mentia calycina* and *Claytonia australasica* var. *biflora*) (unc)
Nertera balfouriana
N. ciliata
N. depressa (incl. *N. cunninghamii*)
N. sp. (unnamed; aff. *N. dichotoma*)
- diifolia) (hairs present on both surfaces of lf, straight, not hooked as in *N. dichondrifolia*)
Oreomythis colensoi s.s. (incl. *O.c. var. hispida* and *O.c. var. multifida*)
Orestylidium subulatum
Outisia lactea ssp. *lactea*
O. caespitosa var. *gracilis* (unc)
O. sp. (unnamed) (plants creeping, prostrate, forming extensive patches on damp banks beside streams in beech forest; lf lamina upto 1x1 cm)
Oxalis magellanica
Plantago uniflora
Ptatis angulata agg.
Psychrophila novae-zealandiae (unc)
Schizolema edule (unc)
Smitens (unc)
Stackhousia minima
Stellaria decipiens (incl. *S. parviflora*)
Utricularia novae-zelandiae
Viola cunninghamii
V. filicaulis
Wahlenbergia albomarginata var.

ADVENTIVE PLANTS

GYMNOSPERM TREES

**Abies* sp. (juv.) (unc)

Pinus contorta (juv.) (unc)

**Pseudotsuga menziesii* (juv.) (unc)

* all plants seen were removed

GRASSES

Agrostis capillaris
Anthoxanthum odoratum (unc)
Holcus lanatus (unc)
Festuca nigrescens (unc)
Poa trivialis (unc)

RUSHES

Juncus articulatus (unc)
J. conglomeratus
J. effusus

COMPOSITE HERBS

Cirsium vulgare (unc)
Hieracium caespitosum (unc)
H. pilosella (unc)
Hypochaeris radicata
Mycelis muralis (unc)
Taraxacum officinale s.l. (unc)

DICOT HERBS OTHER THAN COMPOSITES

Cerastium fontanum ssp. *vulgare* (unc)
Epilobium obscurum (unc)
Rumex acetosella (unc)
Sagina procumbens (unc)
Trifolium repens (unc)

No. of indigenous species: 280.

No. of adventive species : 22.

REFERENCES

- Allan, H.H. 1961: *Flora of N.Z.* Vol. I.
 Crookes, M. 1963: *N.Z. Ferns*.
 Edgar, E. 1966: *N.Z. Jl Bot.* 4: 159-84.
 ———— 1970: In *Flora of N.Z.* Vol. 2.
 Franklin, D.A. 1962: *Trans. R. Soc. N.Z. Bot.* 1: 155-73.
 Lloyd, D.G.; Webb, C.J. 1987: *N.Z. Jl Bot.* 25: 99-105.
 Oliver, W.R.B. 1935: The genus *Cephaelis*: *Bishop Mus. Bull.* 132.
 Raven, P.H.; Engelhorn, T. 1971: *N.Z. Jl Bot.* 9: 345-50.
 Wardle, P. 1968: *N.Z. Jl Bot.* 6: 226-36.
 ———— 1969: *N.Z. Jl. Bot.* 7: 76-95.
 Zotov, V.D. 1965: *Rec. Dom. Mus. Wellington* 5: 101-46.

(cont. from p. 7 - *Craspedia* sp.)

clad in long multicellular, non-glandular hairs, with or without cottony tips, over short glandular hairs; margins faintly white with cottony hairs; fls white)

