

HIGHER PLANTS (GYMNOSEPERMS, LYCOPODS, FERNS, FLOWERING PLANTS) OF MT BALDY<sup>1</sup> AND MT SODOM<sup>2</sup>, SOUTHERN ARTHUR RANGE, NORTH-WEST NELSON  
4200-5150 FT (1280-1565 M)

→ <sup>1</sup> Grid ref. S19 968173, alt. c. 5050 ft (1535 m)  
<sup>2</sup> Grid ref. S19 990200, alt. c. 5150 ft (1565 m)


A.P. Druce, Botany Division, D.S.I.R., April 1982

(Revised Sept 1982, June 1983, June 1984,

Based on visits March 1982 and  
Feb. 1990

Oct. 1986,  
June 1988,  
Nov. 1988,  
June 1990,  
June 1991,  
Aug. 1992,  
Aug. 1993)

Rocks: predominantly indurated  
sandy limestone  
(Summit Limestone)



B - seen on Mt Baldy

S - seen on Mt Sodom

unc - uncommon

### INDIGENOUS PLANTS

#### GYMNOSPERM TREES AND SHRUBS

*Lepidothamnus laxifolius* (unc B, S) <sup>?</sup> *C. propinqua* ssp. *propinqua* (B, S)  
*Phyllocladus alpinus* agg. (unc B)

*Podocarpus nivalis* (B, S)

#### DICOT TREES

*Dracophyllum traversii* (incl. *D. pyramidalis*) (B, S)

*Hohertia lyallii* var. (*H. glabratata*) (B)

*Nothofagus menziesii* (B, S)

*Pseudopanax simplex* (incl. *P. variifolius* var. *sinclairii*) (unc S)

*P. sp.* (unnamed; included

in *P. colensoi*, as var. *fiordeensis* and as var. *ternatus*, by Wardle 1968) (unc B, unc S)

#### DICOT SHRUBS

*Archetia traversii* (incl. *A. t. var. australis*) (unc S)

*Anstotelia fruticosa* var. (B, S)

*Brachyglottis bidwillii* var. *viridis* (unc B, unc S)

*B. grayi* (incl. *B. laxifolia*) (unc B)

*Cassinia vauvilliersii* (unc S)

*Coprosma atropurpurea* (B, S)

*C. cheesemanii* ss. (B, S)

*C. ciliata* ssp. *ciliata* (B, S)

*C. depressa* (B, S)

*C. propinqua* ssp. *propinqua* (B, S)

*C. parpusilla* ssp. *parpusilla* (B, S)

*C. pseudocuneata* ssp. (unc S)

*C. rugosa* ssp. *rugosa* (unc B, unc S)

*C. setulata* (unc B, unc S)

*C. sp.(i)* (unnamed) (*C. parviflora*

var. *dumosa* sensu Allan 1961; non

*C. p. var. dumosa* Cheeseman)

(unc B, unc S)

*C. sp.(ii)* (unnamed; aff. *C. pseudocuneata*) (low-growing shrubs with small, thick, arcuate lvs) (unc S)

*C. sp.(iii)* (unnamed) (prostrate shrubs lvs plano-convex, linear; drapes orange) (B, S)

*C. sp.(iv)* (unnamed; aff. *C. colensoi*) (low-growing shrubs with trailing branches) (unc S)

*C. plumosa* (unc S)

*C. sarmentosa* (unc S)

*Cyathodes dealbata* (incl. *C. pumila*) (unc B, S)

*Dracophyllum longifolium* var. (incl. *D. filifolium* & *D. oliveti*) (B, S)

*D. uniflorum* (B, S)

*Gaultheria crassa* (unc S)

*G. depressa* ss. (unc B, unc S)

*G. rupestris* (incl. *G. subcorymbosa*)

(unc S)

\* 2 forms present: one with few hairs,  
the other with many hairs on the lvs

G. sp. (unnamed; included in G.  
depressa, as var. novae-zelandiae,  
by Franklin 1962) (B,S)

Hebe canterbutiensis (B,S)

H. haastii var. (aff. var. humilis;  
see Moore 1961, p. 940) (B,S)

H. macrantha var. brachyphylla (B,S)

H. tetragona var. (H. coarctata) (unc S)

H. topiaria (B,S)

H. vernicosa (S)

H. sp. (aff. H. glaucophylla) (B,S)

Helichrysum intermedium var. (B,S)

H. i. X Anaphalis sp. (H. bellidifoloides  
s.s.) (unc S)

Leucopogon fraseri s.s. (B,S)

L. suaveolens (incl. Cyathodes colensoi)  
(B,S)

Malicytus sp. (i) (unnamed) (B,S)

M. sp. (ii) (unnamed; aff.

M. alpina; prostrate

hermaphrodite shrubs; branches

up to 40 cm long, often rooting; lvs linear-oblong; fr white) (unc B) sp.; lower pinnae not reduced in length) (B,S)

Myrsine divaricata s.s. (B,S)

M. nummularia (unc B,S)

Olearia lacunosa var. lacunosa  
(B,S)

O. nummulariifolia var. (lvs distinctly  
longer than broad, rather than  
suborbicular as in var. nummul-  
ariifolia) (B, unc S)

Pentachondra pumila (B,S)

Pimelea oreophila (B,S)\*

P. prostrata (S)

Pittosporum anomalum (unc B, unc S)

Traversia baccharoides (B)

## DICOT LIANES AND RELATED

### TRAILING PLANTS

Clematis forsteri (unc B)

Muehlenbeckia axillaris (B,S)

M.a. x M. australis (B)

### LYCOPODS

Lycopodium austalianum (unc B,  
unc S)

L. fastigiatum (B,S)

L. scariosum (unc S)

### FERNS

Asplenium lyallii (unc B)

A. sp. (unnamed; aff. A. trichomanes) (unc B)

Blechnum fluitatile agg. (S)

B. pennula-martini (B,S)

B. procetum (unc S)

B. sp. (unnamed) (mountain

lvs linear-oblong; fr white) (unc B)

sp.; lower pinnae not reduced

in length) (B,S)

Cystopteris tasmanica (B,S)

Grammitis magellanica ssp.

nittofageti (unc S)

G. patagonica (B,S)

G. poeppigiana (B,S)

Hymenophyllum sp. (cf. H. villosum) (S)

Hypolepis millefolium (B,S)

Leptopteris superba (unc S)

Ophioglossum cordaceum agg. (unc B,  
unc S)

+ ("The Dunedin-Cass Variety"  
of Crookes 1963)

*Polystichum cystostegia* (unc B)

*P. vestitum* (B,S)

*Hierochloe novae-zelandiae* (unc B,S)

*Lachnagrostis* sp. (unnamed; included in *L. filiformis*, as var *semiglabra*, by Zotov 1965) (B,S)

*Microlaena colensoi* s.s. (unc B,S)

*Poa breviglumis* (B,S)

*P. celosa* (unc B, unc S)

*P. cita* ssp. *cita* (B)

*P. colensoi* (B,S)

*P. imbecilla* (unc B)

*P. kitkii* (unc S)

*P. maia* (S)

*P. sp.* (*P. novae-zelandiae* or

*P. subvestita*) (B,S)

### GRASSES

{ *Elymus canysii* (B)

{ *E. sp.* (aff. *E. tectorius*) (plants glaucous; awns 4-6 cm long) (B,S)

*Agrostis dyeri* s.s. (B,S)

*A. personata* (unc B,S)

*A. sp.* (unnamed; aff. *A. dyeri*)

(panicle not contracted as in *A. dyeri*) (B,S)

*Rytidosperma gracile* (unc S)

*R. nigricans* (unc B,S)

*R. setifolium* (B,S)

*Chionochloa australis* (B,S)

*C. conspicua* ssp. *conspicua* (B,S)

*C. pallens* ssp. *pallens* (B,S)

*Trisetum youngii* (unc B, unc S)

*T. sp.* (i) (unnamed) (plants slender; lf sheath vs. glabrous) (B,S)

*C. sp.* (unnamed) (*C. flavesans* auct.)

South Island, non *C. flavesans*

Zotov (B,S)

*T. sp.* (ii) (unnamed; included in

*T. antarcticum*, as ssp. *tenellum* by Petrie 1912) (unc B,S)

*Deschampsia tenella* (unc B,S)

*Festuca* sp. (aff. *matthewsii*) (tufted, but not tussock-forming, glaucous) (B,S)

*T. sp.* (iii) (unnamed; aff. *T. antarcticum*)

(unc S)

## SEDGES

- Carex acicularis* (B,S)  
*C. comans* (B)  
*C. enysii* (unc B)  
*C. libera* (unc S)  
*C. pyrenaica* var. *cephalotes* (unc B)  
*C. wakatipu* s.s. (B,S)  
*C. sp.(i)* (unnamed; aff. *C. testacea*)  
 (lvs vs. reddish; utricle margins  
not scabrid) (unc B)  
*C. sp.(ii)* (unnamed; aff. *C. wakatipu*)  
 (vs. smaller than *C. wakatipu*;  
utricle dark brown to black) (B,S)  
*C. sp.(iii)* (unnamed) (lvs green,  
stiff, erect; stigmas 3) (B,S)  
*C. sp.(iv)* (unnamed) (lvs vs. recurved;  
stigmas 3) (unc B)  
*C. sp.(v)* (unnamed; aff. *C. coriacea*) (unc B)  
*Carphe alpina* s.s. (B,S)  
*Oreobolus impar* (S)  
*O. pectinatus* (B,S)  
*O. strictus* (unc B, unc S)  
*Schoenus pauciflorus* s.s. ("slender  
form" of Edgar 1970, p. 194) (B,S)  
*Uncinia caespitosa* var. (S)  
*U. divaricata* (B,S)  
*U. filiformis* (S)  
*U. fuscovaginata* var. (plants  
glaucous) (B,S)  
*U. involuta* (unc B,S)  
*U. nervosa* s.s. (unc B, unc S)  
*U. uncinata* (unc S)  
*U. tvpestris* (incl. *U. angustifolia*) (unc S)  
*U. sp.* (unnamed) (*U. affinis*  
sensu Hamlin 1959, non *U.*  
*riparia* var. *affinis* Col. ex  
Clarke) (B)

## RUSHES AND ALLIED PLANTS

- Juncus novae-zelandiae* (unc S)  
*Luzula migrata* (unc B,S)  
*L. picta* s.s. (unc S)  
*L. sp.* (unnamed; included in *L. picta*,  
as var. *limosa*, by Edgar 1966)  
(unc B,S)

## MONOCOT HERBS OTHER THAN

### ORCHIDS, GRASSES, SEDGES,

- Athroroidium candidum* (unc S)  
*Astelia graminea* (B,S)  
*A. lineatis* var. *lineatis* (unc B,S)  
*A. nervosa* s.s. (B,S)  
*A. nivicola* s.s. (unc B, unc S)  
*A. petriei* (B,S)  
*A. skottsbergii* (S)  
*A. sp.* (unnamed; included in *A.*  
*nivicola*, as var. *moticeae*, by  
Moore 1966) (unc S)  
*Bulbinella hookeri* (B)  
*Libertia peregrinans* (unc S)  
*Phormium cookianum* (B,S)

## COMPOSITE HERBS

- Abrotanella linearis* s.s. (B,S)  
*Anaphalis* sp. (*Helichrysum bell.* s.s.) (B,S)  
*Brachyglottis* sp. (B,S)  
*Brachyscome radicata* (unc S)  
*B. sinclairii* s.s. (B,S)  
*Celmisia dallii* (unc B,S)  
*C. incana* (incl. *C. discolor*) (unc B,S)  
*C. laticifolia* (unc B,S)  
*C. montoi* ssp. (B,S)  
*C. petriei* (B,S)

*C. spectabilis* ssp. (B,S)

*C. traversii* (unc B)

*Craspedia* sp.(i) (unnamed; included in *C. lanata*, as var. *elongata*, by Allan 1961) (B,S)

*C. sp.*(ii) (summer-green; lvs elliptic-spathulate, green, both surfaces clad in non-glandular, multicellular hairs; margins faintly to distinctly white with cottony hairs; fls white) (B)

*C. sp.*(iii) (summer-green; lvs elliptic-spathulate, green, both surfaces densely clad in short glandular hairs < 0.3 mm long; midrib with or without long non-glandular hairs; margins ± faintly white with cottony hairs; fls white) (B,S)

*Gnaphalium audax* (S)

*G. mackayi* ss. (unc B, unc S)

*G. ruahinicum* (B,S)

*Lagenifera strangulata* (S)

*Haptinella* sp. (unnamed; included in *L. squalida*, as var. *mediana*, by Lloyd and Webb (1987)) (B,S)

*Microrsetis scapigera* var. (B,S)

*Pseudognaphalium* sp. (aff. *P. utcoalbum*) (mountain sp.; lvs mostly oblong-spath. to nw oblong, rounded) (unc B, unc S)

*Raoulia glabra* (B,S)

*R. grandiflora* var. *grandiflora* (S)

*Senecio glaucophyllus* ssp. *glaucophyllus* (B,S)

*S. waitavensis* (B,S)

*Taraxacum magellanicum* (B,S)

### DICOT HERBS OTHER THAN COMPOSITES

*Acaena anserinifolia* (B,unc S)

*A. profundeincisa* (B,S)

*Aciphylla ferox* (B,S)

*Anemone tanuicaulis* (B)

*Anisotome aromatica* (B,S)

*A. deltoidea* (S)

*A. filifolia* (unc B,unc S)

*A. haastii* (unc S)

*A. pilifera* (B,S)

*Cardamine* sp.(i) (*C. debilis* agg.) (plants slender, much-branched; lvs small; fls small) (B)

*C. sp.*(ii) (unnamed) (lvs mostly simple; blade ± reniform) (B,S)

*C. sp.*(iii) (*C. debilis* agg.) (cf. "Long Style" of Pritchard 1957) (S)

	E.sp.(ii)(unnamed; aff. <i>E.petrici</i> )	7
<i>Cheesemannia latisiliqua</i> (B,S)	(incl. <i>C.gibbsii</i> ) (corolla pure white)(B) <i>Forstera mackayi</i> (S)	
<i>Colobanthus affinis</i> (B,unc S)	<i>F.tanella</i> (B,S)	
<i>C.canaliculatus</i> (B,S)	<i>Galium perpusillum</i> s.s.(B,S)	
<i>C.wallii</i> (incl. <i>C.masoniae</i> )(unc B,unc S)	<i>Gentiana bellidifolia</i> (B,S)	
<i>C.sp.</i> (unnamed)(lvs and sepals strongly acicular; lvs unbordered; sepals 5, broader than lvs, without a thickened midrib)(unc B,unc S)	<i>G.filipes</i> (unc B,S) <i>G.gracilifolia</i> (incl <i>G.vernicosa</i> )(B,S)	
<i>C.canaliculatus</i> × <i>C.wallii</i> (unc S)	<i>G.sp.</i> (unnamed; aff. <i>G.corym-</i> <i>bifera</i> )(unc B,S)	
<i>Kallstroemia dieffenbachii</i> (unc B,unc S)	<i>Geranium</i> sp.(aff. <i>G.microph.</i> )(B,S)	
<i>K.laxa</i> (unc B,S)	<i>Geum leiospermum</i> (unc B)	
<i>K.multiflora</i> (unc B,S)	<i>G.sp.</i> ( <i>G.parviflorum</i> agg.)(stock short; terminal leaflet about as long as broad; fls up to 13 mm diam.)(B,S)	
<i>Epilobium chlorifolium</i> (B,S)	<i>Gingidia decipiens</i> (B,S)	
<i>E.cockayneum</i> (B,S)	<i>G.montana</i> (unc B)	
<i>E.elegans</i> (unc S)	<i>Hydrocotyle microphylla</i> (B)	
<i>E.glabellum</i> s.s.(B,S)	<i>H.moschata</i> (unc B)	
<i>E.gracilipes</i> (B,S)	<i>H.sp.</i> (unnamed; included in <i>H.novae-zealandiae</i> , as var. <i>montana</i> , by Kirk 1899)(B,S)	
<i>E.margaretae</i> (unc B)	<i>Hypericum japonicum</i> agg.(unc S)	
<i>E.petnitens</i> (unc S)	<i>Mentha cunninghamii</i> (B)	
<i>E.pubens</i> (unc B)	<i>Myosotis brockiei</i> (unc B)	
<i>E.vernicosum</i> (B,S)	<i>M.macrantha</i> (B,S)	
<i>E.sp.</i> (unnamed; included in <i>E.</i> <i>brunnescens</i> , as ssp. <i>minuti-</i> <i>florum</i> , by Raven and Engel- horn, 1971)(B,S)	<i>M.sp.(i)</i> ( <i>M.australis</i> agg.)(fls small, white)(unc S)	
<i>Euphrasia cheesemanii</i> (B)	<i>M.sp.(ii)</i> (unnamed; included in <i>M.pygmaea</i> , as var. <i>drucei</i> , by Moore 1961)(B,S)	
<i>E.townsonii</i> s.s.(unc B,S)	<i>Nertera ciliata</i> (B,S)	
<i>E.zelandica</i> agg. (corolla 4-5 mm wide)(unc B, unc S)	<i>N.sp.</i> (unnamed; aff. <i>N.dichondrifolia</i> )(hrs present on both surfaces of lf, straight, not hooked)(S)	
<i>E.sp.(i)</i> (unnamed; aff. <i>E.laingii</i> ) (unc B,S)		

<i>Notothlaspi</i> sp. ( <i>N. australe</i> avct., non <i>Thlaspi australe</i> Hook.f.) (B,S)	
<i>Oreomyrrhis colensoi</i> s.s. (incl. <i>Scleranthus brockiei</i> (unc B,S))	
O.c. var. <i>hispida</i> and O.c. var. <i>multifida</i> (B,S)	<i>S. uniflorus</i> (unc B)
<i>O. sp.</i> (unnamed; aff. <i>O. tamosa</i> ) (B,S)	<i>Stellaria decipiens</i> (incl. <i>Sparviflora</i> )
<i>Oreopanaxthera alpina</i> (B)	<i>Urtica incisa</i> (B,S) (unc B,S)
<i>Oulisia lactea</i> ssp. <i>lactea</i> (B,S)	<i>Viola cunninghamii</i> (B,S)
	<i>V. lyallii</i> (B)
<i>Oxalis exilis</i> (unc B)	<i>Wahlenbergia albomarginata</i>
<i>O. magellanica</i> (B,S)	var. <i>albomarginata</i> (incl. <i>W.</i>
<i>Parahellebore linifolia</i> ssp. <i>linifolia</i> (B)	<i>pygmaea</i> var. <i>taxa</i> and <i>W.</i>
<i>P. lyallii</i> (B)	<i>brockiei</i> ) (B,S)
<i>Phyllachne clavigera</i> (incl. <i>P. colensoi</i> ) (B,S)	
<i>Plantago lancea</i> s.l. (incl. <i>P. n.z.</i> ) (unc B,S)	<u>ADVENTIVE PLANTS</u>
<i>P. raoulii</i> (B,S)	(see p. 9)

*Pratia angulata* agg. (B,S)

#### REFERENCES

*Psychrophila novae-zealandiae* (unc B,S)

*Ranunculus altus* (unc B)

*R. foliosus* (unc S)

Allan, H.H. 1961: "Flora of N.Z." Vol. 1.

Crookes, M. 1963 "N.Z. Ferns"

Domin, K. 1908: Engl. Bot. Jb 40: 573-85.

Edgar, E. 1966: N.Z. Jl Bot. 4: 157-84.

— 1970: "Flora of N.Z." Vol. 2.

Franklin, D. 1962: T.R. Soc. N.Z. Bot. 1: 157-73

Hamlin, B.G. 1959: Dom. Mus. Bull. 19.

Kirk, T. 1899: "The Students' Flora of N.Z."

Moore, L.B. 1961: "Flora of N.Z." Vol. 1.

— 1966: N.Z. Jl Bot. 4: 201-40

Petrie, D. 1912: T.N.Z. Inst. 44: 179-87.

Pritchard, G.G. 1957: T.R. Soc. N.Z. 85: 75-89.

Raven, P.H.; Engelman, T. 1971: N.Z. Jl Bot. 9: 345-50

Zotov, V.D. 1965: Rec. Dom. Mus. Wellington 5: 101-46.

Lloyd, D.G.; Webb, C.J. 1987: N.Z. Jl Bot. 29: 99-105.

*R. insignis* (B,S)

*R. membranifolius* (unc B, unc S)

*R. mirus* (B,S)

*R. verticillatus* (unc B,S)

*Schizolema* sp. (unnamed; included in *S. haastii*, as ssp. *cyanopetalum*, by Domin 1908) (B,S)

## ADVENTIVE PLANTS

### GRASSES

- Agrostis capillaris* (B, unc S)
- Dactylis glomerata* (B)
- Poa annua* (B, unc S)
- P. pratensis* (B)

*Mycelis muralis* (B, S)

*Senecio jacobaea* (unc B)

*Taraxacum officinale* s.l. (unc B)

### SEDGES

- Carex ovalis* (B)

### DICOT HERBS OTHER THAN COMPOSITES

*Cerastium fontanum* ssp. *vulgare* (B, S)

*Digitalis purpurea* (B)

*Linum catharticum* (S)

*Sagina procumbens* (B, unc S)

*Stellaria media* (B)

*Trifolium repens* (B, S)

*Veronica arvensis* (B)

### COMPOSITE HERBS

- Cirsium arvense* (B)
- C. vulgare* (B)
- Hieracium pilosella* (unc B, unc S)
- H. sp.* (unc B, unc S)
- Hypochaeris radicata* (B, S)

No. of adventive species: 22.

(20 on Baldy; 11 on Sodom)

No. of indigenous species: 267.

(212 on Baldy - 42 confined there;

225 on Sodom - 55 confined there;

170 seen on both mountains)

