

INDIGENOUS HIGHER PLANTS (LYCOPODS, QUILLWORTS, FERNS,
GYMNOSPERMS, FLOWERING PLANTS) OF MTS COBB¹, PROSPECT², RAN-
OLF³ AND MYTTON^{4†}, PEEL RA., N.W.NELSON, 3500-5650 FT/1060-1720M

¹ Grid ref. S13 921558, alt. c. 5625 ft/1710m

² Grid ref. S13 940536, alt. c. 5550 ft/1685m

→ ³ Grid ref. S13 939520, alt. c. 5650 ft/1720m


⁴ Grid ref. S13 990493, alt. c. 5000ft/1520 m

† excluding limestone area (for which see LIST 211)

A.P. Druce, Botany Division, D.S.I.R., and B.V. Sneddon,
Botany Dept, Victoria University of Wellington, April 1982
(Revised Sept. 1982, June 1983, June 1984, Nov. 1984, June 1986,
Aug. 1986, Oct. 1986, June 1988, Sept. 1990, June 1991, Aug. 1992, Aug. 1993)

(Based on visits Jan. 1980, Nov. 1980, March 1982,
Dec. 1985, April 1986)

Rocks: mainly schist, phyllite, greywacke



(unc) - uncommon or local

GYMNOSPERM TREES AND SHRUBS

Halocarpus bidwillii

H. biformis

Lepidothamnus laxifolius

Phyllocladus alpinus agg.

Brachyglottis adamsii

B. bidwillii var. *vitidis*

B. rotundifolia var. (incl. *B. buchananii* & *B. cockaynei*) (unc)

Cassinia vauvilliersii

Podocarpus nivalis

Coprosma atropurpurea (unc)

C. cheesemanii ss.

C. ciliata ssp. *ciliata*

C. colensoi ss.

C. crenulata (unc)

C. depressa

C. foetidissima

C. petrusilla ssp. *petrusilla*

C. pseudocuneata ssp.

C. setulata (unc)

C. sp. (i) (unnamed; aff. *C. pseudocuneata*) (low-growing shrubs with small, thick, arcuate lvs)

C. sp. (ii) (unnamed) (*C. parviflora* var. *dumosa* sensu Allan 1961, non *C. p. var. dumosa* Cheeseman)

C. sp. (iii) (unnamed; included in *C. parviflora* by Oliver, 1935, and others) (lf undersurface not minutely hairy as in *C. parviflora*; drupes vs. unpigmented, occ. purplish red)

DICOT SHRUBS

Archeria traversii (incl. *A. t.* var. *australis*)

Aristotelia fruticosa var.

<i>C. sp. (iv) (unnamed; aff. <i>C. colensoi</i>)</i>	<i>Leucopogon fasciculatus</i>
(low-growing shrubs with trailing branches) (unc)	<i>L. fraseri</i> ss.
<i>C. cheesemanii</i> × <i>C. ciliata</i> (unc)	<i>L. suaveolens</i> (incl. <i>Cyathodes colensoi</i>)
<i>Cyathodes dealbata</i> (incl. <i>C. pumila</i>)	<i>Malicytus</i> sp. (i) (unnamed)
<i>C. petrifolia</i>	<i>M. sp. (ii) (unnamed; aff. <i>M. alpinus</i>)</i> (prostrate hermaphrodite shrubs; branches up to 40 cm long, often rooting; lvs linear-oblong; fr white) (unc)
<i>C. sp. (C. juniperina agg.)</i> (lvs small, us. < 1 cm x < 1 mm)	
<i>Dracophyllum longifolium</i> var. (incl. <i>D. filifolium</i> and <i>D. oliveri</i>)	
<i>D. pubescens</i> var. <i>pubescens</i>	<i>Myrsinodivaricata</i> ss
<i>D. uniflorum</i>	<i>M. nummularia</i>
<i>Gaultheria crassa</i>	<i>Olearia colensoi</i> var. <i>colensoi</i> (unc)
<i>G. depressa</i> ss.	<i>O. lacunosa</i> var. <i>lacunosa</i>
<i>G. sp. (unnamed; included in <i>G. depressa</i>, as var. <i>novae-zelandiae</i>, by Franklin 1962)</i>	<i>O. nummulariifolia</i> var. (lvs distinctly longer than broad, rather than suborbicular as in var. <i>nummulariifolia</i>) (unc)
<i>G. crassa</i> × <i>G. nubicola</i> (unc)	
<i>G. macrostigma</i> × <i>G. sp. (unnamed)</i> (unc)	
<i>Hebe albicans</i> var. <i>albicans</i> (unc)	<i>Pentachondra pumila</i>
<i>H. canterburiensis</i>	<i>Pimelea oreophila</i>
<i>H. ciliolata</i>	<i>Pittosporum anomalum</i> (unc)
<i>H. cochayniana</i> var.	<i>Pittosporum rigidum</i> var. (<i>P. crassicaule</i>)
<i>H. haastii</i> var. (aff. var. <i>humilis</i> ; see Moore 1961, p. 94c)	<i>Traversia baccharoides</i> (unc)
<i>H. mactantha</i> var. <i>brachyphylla</i>	
<i>H. odora</i> ss	
<i>H. tefragona</i> var. (<i>H. coarctata</i>)	
<i>H. topiaria</i>	
<i>H. sp. (Leonotisbebe masoniae*)</i>	
<i>Helichrysum intermedium</i> var. (unc)	

DICOT LIANES AND RELATED/

TRAILING PLANTS

Muehlenbeckia axillaris
M. a. × M. australis (unc)

* (incl. *L.m. var. rotundata*)

Lycopods and Quillworts

Lycopodium australianum
L. fastigiatum
L. scariosum
Isoetes alpinus

Ferns

Asplenium flabellifolium agg. (unc)
A. flaccidum
Blechnum fluviatile agg.
B. pennata-matina
B. procerum
B. sp. (unnamed) (mountain sp.;
 lower pinnae not reduced in
 length) ("The Dunedin-Cass
 Variety" of Crookes (1963))
Cystopteris tasmanica (unc)
Grammitis billardierei
G. paeppigiana
Histicopteris incisa (unc)
Hymenophyllum multifidum
H. villosum
Hypolepis millefolium
Ophioglossum coriaceum agg. (unc)
Polystichum cystostegia (unc)
P. vestitum
Schizaea australis

Orchids

Aporostylis bifolia
Caladenia lyallii
Chiloglottis cornuta

Corybas trilobus ss.

Prasophyllum colensoi

GRASSES

A. dyeri ss.

A. muelleriana

A. personata

A. sp. (unnamed; aff. A. dyeri)
 (panicle not contracted as in
A. dyeri)

Chionochloa australis

C. cheesemanii (unc)

C. conspicua ssp. *conspicua*

C. pallens ssp. *pallens*

C. rubra ssp. *occulta*

C. sp. (unnamed) (*C. flavesans*
 avct. South Island, non *C.*
flavesans Zotov)

C. australis × *C. pallens* (unc)

Deschampsia novae-zelandiae

D. tenella

Deyrollea aucklandica

Festuca sp. (aff. *F. matthensis*) (tufted,
 but not tussock-forming, glaucous)
 (unc)

Hierochloe novae-zelandiae

H. recurvata (unc)

Koeleria cheesemanii (unc)
Lachnagrostis lyallii ss. (incl. *L.
 filiformis* var. *somiglabra*)
L. sp. (i) (L. Henderson)
L. sp. ii (unnamed) (awn short or absent)
Microlaena colensoi s.s.
Poa breviglumis
P. celsa
P. colensoi
P. dipsacea (unc)
P. kirkii
P. maia
P. novae-zelandiae
P. subvestita

SEDGES

Carex acicularis
C. echinata var. (*C. stellulata* var.
australis)
C. flaviformis (unc)
C. gaudichaudiana (unc)
C. libera (unc)
C. pyrenaica var. *cephalotes*
C. sinclairii
C. wakatipu s.s. (unc)
Carpha alpina s.s.
Eleocharis acuta (unc)
Gahnia procera
Isolepis aucklandica
I. habra
Oreobolus pectinatus
O. strictus (unc)
Schoenus pauciflorus s.s. (= "slender
 form" of Edgar 1970, p. 194)

Rytidosperma gracile
R. nigrans (unc)
R. setifolium
Simplicia buchananii (unc)
Trisetum youngii
T. sp. (i) (unnamed) (plants slender;
 lf sheath glabrous or nearly so)
T. sp. (ii) (unnamed; included in *T.
 antarcticum*, as *ssp. tenellum*,
 by Petrie 1912)

Uncinia astonii
U. caespitosa
U. clavata
U. divaricata
U. drucei (unc)
U. filiformis
U. fuscovaginata var. (glaucous)
U. longifructus (unc)
U. nervosa s.s. (unc)
U. silvestris (incl. *U. affinis*)

U.sp. (unnamed) (U. affinis

*Hamlin 1959, non U. riparia var.
affinis Col. ex Clarke)*

A.sp. (unnamed; aff. A. nervosa)

*(Ivs broad; not conspicuously white
adaxially as in A. nervosa)*

Bulbinella hookeri (unc)

Libertia pulchella

Luzuriaga parviflora

Phormium cookianum

Potamogeton cheesemanii

P.suboblongus(unc)

RUSHES AND ALLIED PLANTS

Centrolepis ciliata

Gaimardia setacea

Juncus antarcticus(unc)

J.novae-zelandiae

J.pusillus(unc)

Luzula colensoi

L.migrata

L.subclavata(unc)

L.sp.(unnamed; included in L.

*ciliata, as var. petricana, by Edgar
1966)*

L.coldensoi × L.migrata (unc)

Marsippospermum gracile

COMPOSITE HERBS

Abrotanella caespitosa(unc)

A.linearis s.s.

A.pusilla(unc)

*Anaphalis sp. (*Helichrysum bellidoides* s.s.)*

*Brachyglottis sp. (unnamed; in-
cluded in *B.bellidoides*, as var.
crassa, by Nordenstam 1978)*

Brachyscome radicata

Calmisia alpina(unc)

C.bellidoides(unc)

C.gibbsii

C.hieracifolia ssp.

*C.incana (incl. *C.discolor*)*

C.laticifolia

*C.lateralis (incl. *C.l.* var. *villosa*)
(unc)*

*C.montoi ssp. (*C.coriacea*
var. *lancifolia*)*

C.tupastris

C.sessiliflora

C.similis(unc)

*C.sinclairii (incl. *C.allani* & *C.dutrietzii*)*

MONOCOT HERBS OTHER THAN ORCHIDS

GRASSES, SEDGES, RUSHES

Arthropodium candidum(unc)

Astelia graminea

A.linearis var. linearis

A.nervosa s.s.

A.nivicola s.s.

A.petrici

A.skottsbergii

C. spectabilis ssp.

C. traversii

C. sp. (unnamed; aff. C. gracilanta)

(plants strongly rhizomatous) (unc)

C. similis × *C. spectabilis* (unc)

Craspedia sp.(i)(unnamed)(plants summer-green; lvs linear-lanceolate, up to 12 X 0.6cm, vs. less, green to white, thinly to densely clad in cottony hairs; scapes up to 25cm, vs. <15cm fls white) (L. Henderson)

C. sp. (ii) (unnamed) (plants summer-green; lvs green, both surfaces densely clad in multicellular, non-glandular hairs, margins ± white with cottony hairs; fls white)

C. sp. (iii) (plants summer-green; lvs green, both surfaces sparsely clad in hairs, or almost glabrous; fls white)

C. sp. (iv)

L. sp. (unnamed; included in *L. squalida*, as *spp. mediana*, by Lloyd and Webb, 1987)

Leucogenes grandiceps

Microseris scapigera var

Raoulia eximia var. (*R. rubra*)

R. glabra (unc)

R. grandiflora var. *grandiflora*

Senecio wairauensis

Taraxacum magellanicum

DICOT HERBS OTHER THAN COMPOSITES

Acaena fissistipula (unc)

A. profundeincisa

Aciphylla anomala

A. colensoi var. *colensoi*

A. fetox

A. polita var

A. sp. (unnamed; aff. A. glaucescens) (unc)

Anisotome aromatica

A. doltoidea

A. haastii

A. pilifera

A. sp. (i) (unnamed; included in

A. imbricata, as var. *prostrata*, by Dawson 1961) (unc)

A. sp. (ii) (unnamed) (turf-forming in seepages; infls up to 5cm tall)

Cardamine sp.(i) (*C. debilis* agg.)

(cf. "Long Style" of Pitchard 1957)

C. sp. (ii) (unnamed) (lvs mostly simple; blade ± reniform) (unc)

Haastia sinclairii s.s.

Lagenifera pinnatifida (unc)

L. strangulata

Leptinella pyrathrifolia agg.

<i>C. sp. (iii) (aff. <i>C. corymbosa</i>)</i> (L. Henderson)	hoch, 1971)
<i>C. sp. (iv) (unnamed) ("Scree Race")</i> of Pritchard (1957)	<i>E. sp. (ii) (unnamed; aff. <i>E. globellum</i>)</i> <i>Euphtasia revoluta</i>
<i>C. sp. (v) (<i>C. dabilis</i> agg.) (cf. "Tussock Race" of Pritchard 1957)</i>	<i>E. townsonii</i> ss.
<i>Cheesemanii latisiliqua</i> (incl. <i>C. gibbsii</i>) (unc)	<i>E. sp. (unnamed; aff. <i>E. laingii</i>)</i>
<i>Colobanthus aciculatus</i> (unc)	<i>Forstera mackayi</i>
<i>C. affinis</i> s.s. (unc)	<i>F. sedifolia</i>
<i>C. apetalus</i> (incl. <i>C. a.</i> var. <i>alpinus</i>) (unc)	<i>F. tenella</i> (unc)
<i>C. canaliculatus</i> (unc)	<i>Galium perpusillum</i> s.s.
<i>C. wallii</i> (incl. <i>C. masoniae</i>) (unc)	<i>Gentiana bellidifolia</i>
<i>C. sp. (unnamed; aff. <i>C. buchananii</i>)</i> (Mts Cobb, Prospect, Ranolf) (unc)	<i>G. gracilifolia</i> (incl. <i>G. vernicosa</i>)
<i>C. sinclairii</i> (unc)	<i>G. montana</i> (incl. <i>G. m.</i> var. <i>stolon- ifera</i> , <i>G. patula</i> , and <i>G. ballidi- folia</i> var. <i>pulchella</i>)
<i>Crassula novae-zelandiae</i>	<i>G. sp. (i) (unnamed)</i>
<i>Drosera arcturi</i>	<i>G. sp. (ii) (unnamed; aff. <i>G. corymbifera</i>)</i>
<i>D. spathulata</i> (unc)	<i>Geranium</i> sp. (aff. <i>G. microphyllum</i>)
<i>Epilobium brunnescens</i> ss. (unc)	<i>Geum leiospetrum</i> (unc)
<i>E. chlorifolium</i>	<i>G. uniflorum</i>
<i>E. cochayneanum</i>	<i>G. sp. (G. parviflorum agg.)</i> (Stock short; terminal leaflet about as long as broad; fls up to 13 mm diam.)
<i>E. komarovianum</i> (unc)	<i>Gingidia decipiens</i>
<i>E. macropus</i> (unc)	<i>Gonocarpus aggregatus</i> (unc)
<i>E. margaretae</i>	<i>G. micranthus</i> ssp. <i>micranthus</i> (unc)
<i>E. pernitens</i>	<i>Giratiola nana</i> (unc)
<i>E. porphyrium</i> (unc)	<i>Hydrocotyle sulcata</i>
<i>E. tubromarginatum</i> (unc)	<i>H. sp. (unnamed;</i>
<i>E. tasmanicum</i> (unc)	included in <i>H. novae-zelandiae</i> ,
<i>E. sp. (i) (unnamed; included in <i>E. brunnescens</i>, as ssp. <i>minuti- florum</i>, by Raven and Engel-</i>	as var. <i>montana</i> , by Kirk (1899)

Hypела sp. (unnamed) (minute herb;

fls 3mm diam.; corolla lobes narrow-ly (lanceolate) (L Henderson)

O.sessilifolia ssp. *sessilifolia*

O.simpsonii

Kelleria dieffenbachii

K.laxa (unc)

K.multiflora

Lilaeopsis ruthiana (unc)

Limosella lineata (unc)

Oxalis magellanica

Mitrasacme novae-zelandiae (unc)

Parahelbe cheesemanii (unc)

Montia fontana ss.

P.decora (unc)

Neopaxia sp. (incl. *Montia calycina* ?)

P.lyallii (unc)

Claytonia austroasica var. *biflora*)

Phyllachne clavigera (incl. *P.celensis*)

Myosotis lyallii (incl. *M.l.* var. *Townsonii* and *M.elderi*) (unc)

Plantago lanigera s.l. (incl. *P.h.z.*)

M.macrantha (unc)

P.raoulii

M.sp.(i) (unnamed; included in *M.pygmaea*, as var. *drucei*, by Moore (1961)) (unc)

P.triandra s.s. (L. Henderson)

M.sp.(ii) (*M.australis* agg.) (fls small, white)

P.uniflora

Myriophyllum pedunculatum ssp. *n.z.*

Pratia angulata agg.

M.triphyllum (unc)

P.mactodon s.s. (unc)

Nertera balfouriana

Psychrophila novae-zelandiae

N.ciliata

Ranunculus altus

Notothlaspi sp. (*N.austrole* avct.)

R.foliosus

Oreomyrrhis colensoi s.s. (incl. *O.c.* var. *hispida* and *O.c.* var. *multifida*)

R.glabrifolius (unc)

O.sp. (unnamed; aff. *O.ramosa*) (unc)

R.insignis

Oreostylidium subulatum (unc)

R.vetticillatus

Otisia caespitosa ssp. *caespit.*

R.sp. (unnamed; aff. *R.foliosus*)

O.lactea ssp. *lactea*

(L. Henderson)

Schizolema haastii s.s.
S. nitens

10

Scleranthus uniflorus (unc)
Stellaria decipiens (incl. S. parvi-
flora)
S. gracilenta
Urtica incisa
Viola cunninghamii
V. lyallii (unc)
Wahlenbergia albomarginata
var. albomarginata (incl. W. pyg-
maea var. laxa & W. brockiei)

No. of species: 340.

REFERENCES

11

- Allan, M.H. 1961: "Flora of New Zealand". Government Printer, Wellington. Vol. 1. 1085 pp.
- Crookes, M. 1963. "New Zealand Ferns". Ed. 6. Whitcombe & Tombs. 408 pp.
- Dawson, J.W. 1961: A revision of the genus Anisotome. Univ. Calif. Publics Bot. 33: 1-98.
- Edgar, E. 1966: Luzula in New Zealand. N.Z. Jl Bot. 4: 159-84.
— 1970: Cyperaceae. Pp. 167-285. In "Flora of New Zealand". Government Printer, Wellington. Vol. 2. 354 pp.
- Franklin, D. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans. R. Soc. N.Z. Bot. 1: 155-73
- Nordenstam, B. 1978: Opéra bot. 44: 1-84.
- Hamlin, B.G. 1959: A revision of the genus Ulinaria. Dom. Mus. Bull. 19. 106 pp.
- Kirk, T. 1899: "The Student's Flora of New Zealand". Govt. Printer, Wellington. 408 pp.
- Lloyd, D.G.; Webb, C.J. 1987: The reinstatement Heptinella at generic rank, & the status of the "Cotyledæ" (Asteraceæ, Anthemideæ). N.Z. Jl Bot. 25: 99-105.
- Moore, L.B. 1961: In "Flora of New Zealand". Government Printer, Wellington. Vol. 1. 1085 pp.
- Oliver, W.R.B. 1935: The genus Coprosma. Bull. Bishop Mus. Honolulu 132. 207 pp.
- Pritchard, G.G. 1957: Experimental taxonomic studies on species of Cardamine Linn. in New Zealand. Trans. R. Soc. N.Z. 85: 75-89.
- Raven, P.H.; Engelhorn, T. 1971: New taxa and new combinations in Australian Epilobium (Onagraceæ). N.Z. Jl Bot. 9: 345-50.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.
- Wardle, P. 1968: The taxonomy and distribution of the stipitate species of Pseudopanax in New Zealand. N.Z. Jl Bot. 6: 226-36.

