


INDIGENOUS HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING PLANTS) OF HERENGA SCENIC RESERVE†, 17.5 KM S.S.W. OF EAST CAPE, S.L. TO 300 FT (90 M)

†Gnd. ref. N 72 830380

A.P. Druce, Wellington, May 1983 (revised July 1985, Aug. 1990, Feb. 1994)
Based on visit Jan. 1983



GYMNOSPERM TREES

Dacrycarpus dactydioides

MONOCOT TREES

Cordyline australis

DICOT TREES

Alectryon excelsus var. *excelsus*.
Aristotelia serrata
Beilschmiedia tawa
Carpodetus serratus
Corynocarpus laevigatus.
Dysoxylum spectabile

Flacocarpus dentatus (incl. *E.d.* var. *obovatus*)
Fuchsia excorticata
Hedycarya arborea
Hoheria populnea var. *lanccolata*
Knightia excelsa
Macropiper excelsum var. *excelsum*
Melicope ternata
Melicytus tamiflorus ssp. *tamiflorus*
Metrosideros excelsa

- Myrsine australis
- Nestegis lanceolata
- Oleaia tani var. colorata
- Streblus heterophyllus
- Pennantia corymbosa
- Pittosporum tenuifolium ssp. tenuifolium
- Pseudopanax arboreus var. arboreus
- P. crassifolius
- Schefflera digitata
- Vitex lucens
- Weinmannia racemosa

DICOT SHRUBS

- Brachyglottis repanda s.s.
- Carmichaelia cunninghamii var. (incl. C. flagelliformis and C. hookeri)
- Cassinia leptophylla var. leptophylla
- Coprosma areolata
- C. rhamnoides ssp. rhamnoides (incl. C. polymorpha)
- C. robusta
- Coriaria arborea
- Geniostoma rupestre var. ligustrifolium
- Hebe stricta var. stricta
- H. sp. (Veronica arborea)
- Kunzea ericoides agg.
- Leptospermum scoparium

MONOCOT LIANES

- Freycinetia baueriana ssp. banksii
- Ripogonum scandens

DICOT LIANES

- Calystegia fuguniorum
- Clematis cunninghamii
- Metrosideros colensoi (incl. M.c. var. pendens)
- M. perforata
- Rubus cissoides var. cissoides

LYCOPODS

- Lycopodium volubile

FERNS

- Adiantum cunninghamii
- Asplenium flaccidum
- A. gracillimum
- A. oblongifolium
- Blechnum chambersii
- B. filiforme
- B. sp. (unnamed) (common lowland sp. lower pinnae reduced in length)
- Cyathea dealbata
- C. medullaris
- Doodia media ssp. australis
- Histiopteris incisa
- Hymenophyllum demissum
- Lastreopsis glabella
- L. hispida
- Leptopteris hymenophylloides
- Pellaea rotundifolia
- Phymatosorus pustulatus
- P. scandens
- Pneumatopteris pennigera
- Polystichum richardii
- Pteridium esculentum
- Pteris pendula
- Dicksonia squarrosa
- Pyrosia eleagnifolia

GRASSES

Cortaderia fulvida
Dichelachne chinita
Microlaena avenacea
M. stipoides
Oplismenus imbecillus

SEDGES

Carex comans
C. flagellifera
C. geminata s.s.
C. lambertiana
C. lessoniana
C. pumila
C. sp. (unnamed; aff. *C. geminata*)
 (much larger than *C. geminata*;
 lvs up to 2.5 cm wide)
Cyperus ustulatus
Gahnia lacera
G. pauciflora
Isolepis inundata
I. nodosa

RUSHES

Juncus australis
J. gregiflorus

**MONOCOT HERBS OTHER THAN
GRASSES, SEDGES, RUSHES**

Astelia solandri
Collospermum hastatum
Dianella nigra
Typha orientalis

COMPOSITE HERBS

Leptinella squalida s.s.
Gnaphalium gymnocephalum
Lagenifera pumila
Pseudognaphalium sp. (unnamed; aff. *P. luteoalbum*) (lvs mostly linear-spathulate to linear, acute)

Senecio rufiglandulosus var.
rufiglandulosus

DICOT HERBS OTHER THAN COMPOSITE

Acaena anserinifolia
Cardamine sp. (*C. debilis* agg.)
 ("Narrow Petal" of Pritchard
 1957)
Epilobium nummulatifolium
E. rotundifolium
Gunnera monoica

Hydrocotyle elongata
H. novae-zeelandiae s.s.
Lobelia anceps
Nertera depressa (incl. *N.*
cunninghamii)
Oxalis exilis
Peperomia tetraphylla
P. urvilleana
Ranunculus reflexus

No. of species: 113.

REFERENCE

Pritchard, G.G. 1957: Experimental taxonomic studies on species of *Cardamine* Linn. in N.Z. Trans. R. Soc. N.Z. 85: 75-89.

