

INDIGENOUS HIGHER PLANTS (PSILOPSIDS, LYCOPODS, FERNS,  
GYMNOSPERMS, FLOWERING PLANTS) OF UPPER TAWARAU G.,  
WEST OF TE KUITI, 250-275 M

Grid ref. N 82 475830

A.P.Druce, Wellington; and C.C.Ogle, N.Z.  
Wildlife Service, Wellington

April 1984 (revised June 1984, June 1985,

Based on visits Nov. 1980 (C.C.O.) July 1985,  
and Jan. 1984 (A.P.D. and C.C.O.) Apt. 1986,  
June 1991

GYMNOSPERM TREES

- Dacrycarpus dacrydioides*
- Dacrydium cupressinum*
- Phyllocladus trichomanoides*
- Podocarpus hallii*
- Prumnopitys ferruginea*
- P. taxifolia*

- Fuchsia excorticata*
- Griselinia littoralis*
- Hedera populnea* var.  
*lanceolata*
- Knightia excelsa*
- Laphamia bullata*
- Melicytus lanceolatus*  
(incl. M.l. var. *latior*)

- Myrsine australis*
- M. salicina*

- Nestegis lanceolata*
- N. montana*

- Olearia tani* var. *colorata*
- Pennantia corymbosa*
- Pittosporum tenuifolium* s.l.  
(incl. P.t. ssp. *colensoi*)
- Plagianthus regius* var. *regius*
- Pseudopanax arboreus* var. *arbo.*
- P. crassifolius*
- P. laetus*

MONOCOT TREES AND SHRUBS

- Cordyline banksii*
- C. indivisa*

DICOT TREES

- Aristotelia serrata*
- Baileya media tawa* (juv. only)
- Carpodetus serratus*
- Elaeocarpus dentatus* (incl.  
*E.d.* var. *obovatus*)
- E. hookerianus*

+ Additions by P.J.Bellingham and E.K.Cameron, Dec. 1984



*Quintinia serrata* (incl. *Q. acutifolia* and *Q. elliptica*)

*Schefflera digitata*

*Weinmannia racemosa* var.  
*racemosa*

#### DICOT SHRUBS

*Alseuosmia macrophylla*

*Brachyglottis repanda* s.s.

*Coprosma grandifolia*

*C. lucida* s.s.

*C. propinqua* ssp. *propinqua* (incl.  
*C.p. var. latiseula*)

*C. hammondii* (incl. *C. polymorpha*)

*C. robusta*

*C. rotundifolia*

*C. tenuicaulis*

*C. tanifolia*

*C. propinqua* X *C. robusta*

*Coriaria arborea*

*Fuchsia excorticata* X *F. petiolaris*

*Gaultheria antipoda*

*Hebe stricta* var. *stricta*

*Kunzea ericoides* agg.

*Leptospermum scoparium* var.

*Leucopogon fasciculatus* var.

*Melicope simplex*

*Neomyrtus pedunculata*

*Olearia townsonii* (incl. *O. furfuracea* var. *rubicunda* and *O. townsonii*)

*Q. virgata* var. (lvs smaller  
than in var. *virgata*)

*Pseudopanax anomalous*

*Pseudowintera colorata*

*Rhabdothamnus solandri*

#### DICOT LIANES

*Clematis foetida*

*C. forsteri*

*C. paniculata*

*Metrosideros colensoi* (incl.  
*M.c. var. pendens*)

*M. perforata*

*Muehlenbeckia australis*

*Parsonsia capsularis* var.

*Rubus australis*

*R. cissoides*

#### PSILOPSIDS

*Tmesipteris elongata* (incl. *T.e.*  
ssp. *robusta*)

#### LYCOPODS

*Lycopodium deuterodensum*

*L. variabile* (incl. *L. billardierei* & *L. n.z.*)

*L. volubile*

#### FERNS

*Adiantum cunninghamii*

*Asplenium bulbiferum*

*A. flaccidum*

*A. gracillimum*

*A. polyodon*

*Blechnum chambersii*

*B. colensoi*

- 3
- |                                                                                                                    |                                                |
|--------------------------------------------------------------------------------------------------------------------|------------------------------------------------|
| <i>B. discolor</i>                                                                                                 | <i>Hypolepis ambigua</i>                       |
| <i>B. fluviatile</i> agg.                                                                                          | <i>H. tufo-barbata</i>                         |
| <i>B. membranaceum</i>                                                                                             | <i>Leptopteris hymenophylloides</i>            |
| <i>B. penna-martini</i>                                                                                            | <i>L. superba</i>                              |
| <i>B. vulcanicum</i>                                                                                               | <i>L. hymenophylloides</i> x <i>L. superba</i> |
| <i>B. sp. (i) (unnamed)</i> (common lowland<br>sp.; lower pinnae reduced in<br>length)                             | <i>Paesia scaberrula</i>                       |
| <i>B. sp. (ii) (unnamed)</i> (cliff sp.;<br>lower pinnae not reduced) ("The<br>Green Bay Species" of Crookes 1963) | <i>Phymatosorus diversifolius</i>              |
| <i>Botrychium bifolium</i>                                                                                         | <i>Pneumatopteris pennigera</i>                |
| <i>Ctenopteris heterophylla</i>                                                                                    | <i>Polystichum richardii</i>                   |
| <i>Cyathea smithii</i>                                                                                             | <i>P. silvaticum</i>                           |
| <i>Dicksonia fibrosa</i>                                                                                           | <i>P. vestitum</i>                             |
| <i>D. squarrosa</i>                                                                                                | <i>Pteridium esculentum</i>                    |
| <i>Diplazium austale</i>                                                                                           | <i>Sticherus cunninghamii</i>                  |
| <i>Grammitis billardierei</i>                                                                                      | <i>Trichomanes teniforme</i>                   |
| <i>G. pseudociliata</i>                                                                                            | <i>T. venosum</i>                              |
| <i>Histiopteris incisa</i>                                                                                         | <b>[ORCHIDS]</b>                               |
| <i>Hymenophyllum armstrongii</i>                                                                                   | <i>Caladenia</i> sp.                           |
| <i>H. demissum</i>                                                                                                 | <i>Chiloglottis cornuta</i>                    |
| <i>H. dilatatum</i>                                                                                                | <i>Corybas oblongus</i>                        |
| <i>H. flabellatum</i>                                                                                              | <i>C. rivularis</i> ss.                        |
| <i>H. flexuosum</i>                                                                                                | <i>C. trilobus</i> s.s.                        |
| <i>H. multifidum</i>                                                                                               | <i>Eatinia autumnalis</i>                      |
| <i>H. pulcherrimum</i>                                                                                             | <i>E. mucronata</i> ss.                        |
| <i>H. ratum</i>                                                                                                    | <i>Gastroleia cunninghamii</i>                 |
| <i>H. revolutum</i>                                                                                                | <i>Orthoceras novae-zealandiae</i>             |
| <i>H. sanguinolentum</i> agg.                                                                                      | <i>Pterostylis banksii</i>                     |
|                                                                                                                    | <i>P. graminea</i>                             |
|                                                                                                                    | <i>Thelymitra longifolia</i>                   |

## GRASSES

*Cortaderia fulvida*  
*Elymus hardyioides*  
*Hierochloe redolens*  
*Microlaena avenacea*  
*M. stipoides*  
*Poa anceps* ssp. *anceps*  
*Rytidosperma gracile*

## SEDGES

*Carex diffusa* s.s.  
*C. forsteri*  
*C. geminata* s.s.  
*C. lessoniana*  
*C. secta* s.s.  
*C. solandri*  
*C. vitigera*  
*Bahnia pauciflora*  
*G. setifolia*  
*Isolepis pottsii*  
*I. reticularis*  
*Morelotia affinis*  
*Schoenus maschalinus*  
*Uncinia banksii*  
*U. distans*  
*U. ferruginea*  
*U. gracilenta*  
*U. uncinata*  
*U. zotovii*

## RUSHES

*Juncus gregiflorus*  
*J. sarophorus*  
*Luzula picta* s.s.  
*L. subclavata*  
*L. picta* x *L. subclavata*

## MONOCOT HERBS OTHER THAN [ORCHIDS, GRASSES, SEDGES, RUSHES]

*Athropodium candidum*  
*Astelia fragrans*  
*A. grandis*  
*Collospermum microspermum*  
*Dianella nigra*  
*Libertia grandiflora*  
*L. pulchella*  
*Potamogeton suboblongus*

## COMPOSITE HERBS

*Anaphalis kerriensis*  
*Craspedia minor* var. *minor*  
(incl. *C. major*)  
*Gnaphalium gymnocephalum*

*G. limosum* s.s.  
*Lagenifera pumila*  
*Senecio minimus*  
*S. rufiglandulosus* var.

## DICOT HERBS OTHER THAN [COMPOSITES]

*Acaena ansetinifolia*  
*Cardamine sp(i)(C. debilis agf.)*  
("Long Style" of Pitchard 1957)

C. sp.(ii) ("C. dabilis agg.") ("Glossy leaf" of Pritchard 1957)

Elatostema rugosum

Epilobium nerteroides

Epeduncularz s.s.

E. pubens

E. rotundifolium

Gunnera monoica

Haloragis erecta ssp. erecta

Hydrocotyle dissecta

H. elongata

H. heteromeira

H. microphylla

H. moschata

Mycosotis sp (cf. M. forsteri)

Nertera depressa (incl. N. cunninghamii)

N. sp. (unnamed; aff. N. dichondrifolia) (hairs present on both surfaces of lf, straight, not hooked as in N. dichondrifolia)

Oxalis exilis

O. magellanica agg.

Parahabea diffusa ssp. diffusa

Plantago taoulie var. (lvs narrow-lv lanceolate)

Pratia angulata agg.

Ranunculus reflexus

Schizeilema trifoliolatum  
Stellaria decipiens (incl. S. parviflora)

Urtica incisa

Viola filicaulis

No. of species: 201.