

(265)

HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING
PLANTS) OF LEAD HILLS, N.W. NELSON, 3300-5281 FT (1000-1600M)



A. P. Druce, Wellington, May 1985

(Revised July 1985, June 1986, July 1990, May 1991, Aug. 1992)

Highest point: Lead Hills, 5281 ft, grid ref. S8 967767 Aug. 1993

(Based on visit Feb. 1985)

Rocks: granite and granitic deposits (moraine)



INDIGENOUS SPECIES

(unc) - uncommon or local

GYMNOSPERM TREES AND SHRUBS

- Halocarpus bidwillii*
- H. biformis*
- Lepidothamnus intermedius*
- L. laxifolius*
- L. intermedius* x *L. laxifolius*
- Libocedrus bidwillii*
- Phyllocladus alpinus* agg.
- Podocarpus hallii* (unc)
- P. nivalis*
- P. hallii* x *P. nivalis* (unc)

DICOT TREES

- Dracophyllum traversii* (incl. *D. pyramidalis*)
- Griselinia littoralis*
- Metrosideros umbellata*
- Nothofagus solandri* var. *cliffortoides*
- Pseudopanax linearis*
- P. simplex* (incl. *P. s. var. sinclairii*) (unc)
- P. sp. (unnamed; included in *P. colensoi*, as var. *fildensis* and as var. *ternatus*, by Wardle 1968)*
- Weinmannia racemosa* var. *racemosa* (unc)

DICOT SHRUBS

- Archeria traversii* (incl. *A.t. var. australis*)
- Aristotelia fruticosa* var.
- Brachyglottis adamii*
- B. bidwillii* var. *vividis*
- B. rotundifolia* var. (incl. *B. buchananii* and *B. cockaynei*)

Cassinia laevis

- Coprosma cheesemanii* s.s.
- C. ciliata* ssp. *ciliata*
- C. colensoi* s.s.
- C. crenulata*
- C. foetidissima* (unc)
- C. pseudocuneata* ssp.
- C. setulata* (unc)
- C. sp. (i)* (unnamed; aff. *C. pseudocuneata*) (low-growing shrubs with small, thick, arcuate lvs)
- C. sp. (ii)* (unnamed; aff. *C. colensoi*) (low-growing shrubs with trailing branches)

C. sp. (iii) (unnamed) (*C. patriflora* var. *dumosa* sensu Allan 1961, non *C. p. var. dumosa* Cheeseman)

Cyathodes dealbata (incl. *C. pumila*)

C. campetifolia

C. sp. (C. juniperina agg.)

Dracophyllum longifolium var. (incl. *D. filifolium* & *D. oliveri*)

D. palustre

D. pubescens var. *pubescens*

D. uniflorum var. *uniflorum*

*D. sp. (unnamed; aff. *D. Kirkii*)* (low-growing shrubs; branches often root ing; infl of few-fl'd racemes)

Epaactis alpina

Exocarpus bidwillii

Gaultheria antipoda

G. crassa

*G. sp. (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962)*

Hebe canterburiana
H. ciliolata
H. tetragona var. (*H. coarctata*)
H. topiaria
H. sp.(i) (*Leonohebe masoniae*)
H. sp.(ii) (*Leonohebe mooreae*)
Leptospermum scoparium var.
Melicytus sp. (unnamed)
Myrsine divaricata s.s.
M. nummularia
Olearia colens var. *colensoi*
O. lacunosa var. *lacunosa*
O. nummularifolia var. (lvs distinctly
longer than broad, rather than suborb-
icular as in var. *nummularifolia*)
Pentachondra pumila
Peraxilla tetrapetala (unc)
Pittosporum anomalum
P. rigidum var. (*P. crassicaule*)
Pseudopanax anomalus
Pseudowintera traversii

LYCOPODS

Lycopodium austrianum
L. fastigiatum
L. laterale (incl. *L. lamulosum*)
L. scariosum
L. varium (incl. *L. billardierei* and *L.*
novae-zelandiae) (unc)

FERNS

Blechnum penna-martini
B. pratense
B. sp. (unnamed) (mountain sp.;
lower pinnae not reduced in
length) ("The Dunedin-Cass Variety"
of Crookes and Dobbie, 1963)
Glaucium dicarpa
Grammitis billardierei
G. givenii (unc)
G. magellanica ssp. *nothofageti*
G. poeppigiana

Histiopteris incisa (unc)
Hymenophyllum multifidum
H. villosum
Ophioglossum coriaceum agg.
Polystichum vestitum
Schizaea australis

ORCHIDS

Aporostylis bifolia
Plasophyllum colensoi
Thelymitra cyanea (unc)

GRASSES

Agrostis dyeri s.s.
A. sp. (unnamed; aff. *A. dyeri*)
 (panicle spreading, not contracted
as in *A. dyeri*)
Chiionochloa australis
C. conspicua ssp. *conspicua*
C. pallens ssp. *pallens*
C. rubra ssp. *occulta*
Deyeuxia aucklandica
D. avenoides (incl. *D. a.* var. *brachyantha*)

Festuca sp. (unnamed; aff. *F. rubra*)
Hierochloe novae-zelandiae
Lachnagrostis lyallii s.s.
 (incl. *L. filiformis* var.
semiglabra)

Mictolaena colensoi
Poa colensoi s.s.
Rytidosperma gracile
R. neogriechs
R. setifolium

SEDGES

Carex aciculatis
C. echinata var. (*C. stellulata*
 var. *australis*)

* (incl. *L. m.* var. *rotundata*)

C. gaudichaudiana
C. libera
Carpha alpina s.s.
Eleocharis acuta (unc)
Gahnia procera
Isolepis aucklandica
Oreobolus impar
O. pectinatus
O. strictus
Schoenus pauciflorus s.s. (= "slender form" of Edgar 1970, p. 194)
Uncinia involuta
U. nervosa s.s.
U. rupestris (incl. *U. angustifolia*)
U. zotovii

RUSHES AND ALLIED PLANTS

Centrolepis ciliata
Empodisma minus
Gaimardia setacea
Juncus antarcticus
J. novae-zelandiae
Luzula colensoi
L. migiata
L. tufa ss. (unc)

MONOCOT HERBS OTHER THAN [ORCHIDS, GRASSES, SEDGES, RUSHES]

Astelia graminea
A. linearis var. *linearis*
A. skottsbergii
A. sp. (unnamed; aff. *A. nervosa*)
 (lvs broad, not conspicuously white adaxially as in *A. nervosa*)
Heteroplitron novae-zelandiae
Luzuriaga parviflora
Polygonum cookianum

COMPOSITE HERBS

Abrotanella caespitosa
Anaphalis sp. (*Helichrysum bellidioides* s.s.)
Brachyglossis sp. (unnamed; included

in *B. bellidioides*, as var. *crassa*, by Nordnastam 1978)
Celmisia alpina s.s.
C. dallii (unc)
C. hieracifolia ssp.
C. incana (incl. *C. discolor*)
C. laticifolia
C. lateralis (incl. *C. l.* var. *villosa*)
C. monticola ssp. (*C. coriacea* var. *lancifolia*)
C. parva
C. petriei
C. similis
C. spectabilis ssp.
C. sp. (unnamed; aff. *C. gracilenta*)
 (lvs linear to linear-lanceolate; upper surface and margins ± thinly covered in tomentum; lower surface glabrous, exc. for midrib)
C. similis x *C. spectabilis* (unc)

Craspedia sp. (plants summer-green; lvs green, both surfaces clad in very short hairs or almost glabrous; fls white)

Gnaphalium mackayi ss.
Leptinella sp. (unnamed)*
Microseris scapigera var.
Raoulia eximia var.
R. grandiflora var. *grandiflora*
Senecio wairauensis

DICOT HERBS OTHER THAN COMPOSITES

Aciphylla polita var. (plants more robust & segments coarser than in var. *polita*)
Actinotis novae-zelandiae
Anisotome aromatica
A. deltoidea
A. haastii
A. pilifera

* (included in *L. squalida*, as ssp. *mediana*, by Lloyd & Webb, 1987)

- Chionohebe pulvinaris*
 A.sp. (unnamed; included in *A. imbricata*) *Oxalis simpsonii*
 (as var. *prostrata*, by Dawson 1961) *Oxalis magellanica*
Colobanthus apetalus (incl. *C. a.*)
 var. *alpinus*
Coprosma talbotii
Donatia novae-zelandiae
Dioscorea arcturi
D. spathulata
D. stenopetala
Epilobium heteroides
E. pernitens
Euphrasia townsonii s.s.
 E.sp. (unnamed; aff. *E. laingii*)
Forstera mackayi
 F.sp. (unnamed) (lvs closely placed, not
 recurved, persisting well down stem
 and becoming thin and translucent;
 costa not apparent)
Gentiana gracilifolia (incl. *G. verni-*
 cosci)
G. montana (incl. *G. m.* var. *stolonifera*,
 G. patula, *G. bellidifolia* var. *pulchella*)
G. sp. (unnamed; aff. *G. bellidifolia*) (plants
 forming small to large patches; stems
 decumbent; fls large, creamy white)
Geranium sp. (aff. *G. microphyllum*)
Gonocarpus aggregatus
G. micranthus ssp. *micranthus* (unc)
Hydrocotyle microphylla
Kelleria laxa (unc)
K. tessellata

- Phyllachne clavigera* (incl. *P. colensoi*)
Plantago lanigera s.l.
P. uniflora
Pratia angulata agg.
Ranunculus verticillatus (unc)
Schizolejeum haastii ss.
Viola cunninghamii

ADVENTIVE SPECIES

GRASSES

Holcus lanatus

RUSHES

Juncus articulatus

J. sp. (cf. *J. effusus* + *J. conglomeratus*)

COMPOSITE HERBS

Hypochoeris radicata

DICOT HERBS OTHER THAN COMPOSITES

Ceratium fontanum ssp. *vulgare*

Epilobium obscurum

E.O. × E. heteroides (unc)

Sagina procumbens

No. of indigenous spp.: 200.

No. of adventive spp.: 7.

- Liparophyllum gunnii*
Montia fontana s.s.
Neopaxia sp. (incl. *Montia calycina*
 and *Claytonia austrolasica* var. *biflora*)
Nertera ciliata
N. scapanioides
Oreomytthis colensoi s.s. (incl. *O.c.* var.
 hispida and *O.c.* var. *multifida*)
Oreostylidium subulatum (unc)

REFERENCES

Allan, H.H. 1961: "Flora of New Zealand." Government Printer, Wellington.
Vol. 1. 1085 pp.

Crookes, M.; Dobbie, H.B. 1963: "New Zealand Flora". Ed. 6. Whitcombe and Tombs, 408 pp.

Dawson, J.W. 1961: A revision of the genus Anisotome (Umbelliferae). Univ. Calif. Publics Bot. 33: 1-98.

Edgar, E. 1970: In "Flora of New Zealand". Government Printer, Wellington. Vol 2. 354 pp.

Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans R. Soc. NZ. Bot. 1: 155-73.

Nordenstam, B. 1978: Opera Bot. 44: 1-84.

Wardle, P. 1968: The taxonomy and distribution of the stipitate species of Ricinocarpos in New Zealand. N.Z. Jl Bot. 6: 226-36.

Zetou, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.

Lloyd, D.G.; Webb, C.J. 1987: The reinstatement of Leptinella at generic rank, and the status of the "Cotuleae" (Asteraceae, Anthemideae). N.Z. Jl Bot. 25: 99-105.