

HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS,  
FLOWERING PLANTS) OF GLASGOW RA., N.W. NELSON.

450-1475 M / 1500-4862 FT

(Highest point: 4862 ft, grid ref. S25 513932)

A.P.Druce, Wellington


P.A.Williams, Botany Division, D.S.I.R., Nelson

W.D.Burke, Nelson

June 1986 (revised July 1986, Nov. 1986, June 1988,

Oct. 1988,  
July 1990,  
June 1991,  
July 1992)

(Based on visits Feb. 1986, Dec. 1986)



## INDIGENOUS PLANTS

(unc) - uncommon or local

### GYMNOSPERM TREES AND SHRUBS

*Dacrydium cupressinum*

*Halocarpus biformis*

*Lepidothamnus intermedius* (unc) *colensoi*, as var. *frödensis*

*L. laxifolius*

*Libocedrus bidwillii* (unc)

*Phyllocladus alpinus* agg.

*Podocarpus hallii*

*P. nivalis* (unc)

*Prumnopitys ferruginea*

*Pseudopanax linearis*

*P. simplex* (incl. *P. s. var. sinclairii*)

*P. sp.* (unnamed; included in *P.*)

*colensoi*, as var. *frödensis*

and as var. *tenuatus*, by

Wardle 1968)

*Quintinia serrata* (incl. *Q. acutifolia* and *Q. elliptica*)

*Weinmannia racemosa* var. *racemosa*

### MONOCOT TREES AND SHRUBS

*Cordyline indivisa* (unc)

### DICOT TREES

*Carpodetus serratus*

*Dracophyllum traversii* (incl. *D. pyramidale*)

*Elaeocarpus hookerianus*

*Fuchsia excorticata* (unc)

*Griselinia littoralis*

*Hohoria lyallii* var. (*H. glabrata*) (juv. only) (unc)

*Metrosideros umbellata*

*Myrsine salicina*

*Nothofagus fusca*

*N. menziesii*

*N. solandri* var. *cliffortioides*

*N. truncata*

### DICOT SHRUBS

*Archeria traversii* (incl. *A. t. var. australis*)

*Aristotelia fruticosa* var. (unc)

*Brachyglottis bidwillii* var.

*B. rotundifolia* var. (incl. *B. buchananii* & *B. cockaynei*)

*Cassinia vauvilliersii* (unc)

*Coprosma cheesemanii* ss.

*C. ciliata* ssp. *ciliata*

*C. colensoi* ss.

*C. crenulata*

*C. depressa*

*C. foetidissima*

*C. grandifolia*

*C. perpusilla* ssp. *perpusilla*

<i>C. propinqua</i> ssp. <i>propinqua</i> (unc)	<i>Gaultheria crassa</i> × <i>G.</i> sp. (unnamed) (unc)
<i>C. pseudocuneata</i> ssp. ( <i>Ivs nattan-</i> <i>et than in</i> ssp. <i>pseudocuneata</i> )	<i>G. macrostigma</i> × <i>G.</i> sp. (unnamed)
<i>C. hammoides</i> (incl. <i>C. poly-</i> <i>morpha</i> )	<i>Hebe canterburiana</i> <i>H. ciliolata</i>
<i>C. setulata</i>	<i>H. cockayneana</i> (unc) <i>H. gracillima</i> (unc)
<i>C. sp. (i)</i> (unnamed) ( <i>C. parviflora</i> var. <i>dumosa</i> Allan 1961, non <i>C.</i> p. var. <i>dumosa</i> Cheeseman)	<i>H. macrantha</i> (Cockayne 1928, pp. 315-6) <i>H. salicifolia</i> var. <i>salicifolia</i> (unc)
<i>C. sp. (ii)</i> (unnamed; aff. <i>C. pseudo-</i> <i>cuneata</i> ) (low-growing shrubs with small, thick, arcuate <i>Ivs</i> )	<i>H. sp.</i> ( <i>Leonotis</i> <i>masoniae</i> ) (incl. <i>L. m.</i> var. <i>rotundata</i> )
<i>C. cheesemanii</i> × <i>C. ciliata</i> (unc)	<i>Leptospermum scoparium</i>
<i>Cyathodes dealbata</i> (incl. <i>C. pumila</i> )	<i>Leucopogon fasciculatus</i> var.
<i>C. empetrifolia</i>	<i>Melicytus alpinus</i> s.s.
<i>C. sp.</i> ( <i>C. juniperina</i> agg.) ( <i>Ivs</i> vs. > 1 cm x > 1 mm, up to 1.5 cm x 2.5 mm)	<i>Metrosideros parkinsonii</i> <i>Myrsine divaricata</i> s.s.
<i>Dracophyllum longifolium</i> var. (incl. <i>D. filifolium</i> and <i>D. oliveri</i> )	<i>M. nummularia</i> <i>Neomyrtus pedunculata</i>
<i>D. politum</i> (incl. <i>D. densom</i> )	<i>Olearia avicenniifolia</i> (unc)
<i>D. pubescens</i> var. <i>pubescens</i>	<i>O. colensoi</i> var. <i>colensoi</i>
<i>D. uniflorum</i> var. <i>uniflorum</i>	<i>O. lacunosa</i> var. <i>lacunosa</i>
<i>D. politum</i> × <i>D. pubescens</i> (unc)	<i>Pentachondra pumila</i>
<i>Gaultheria crassa</i>	<i>Pittosporum rigidum</i> var. ( <i>P.</i> <i>classicaule</i> )
<i>G. depressa</i> s.s.	<i>Pseudopanax anomalus</i>
<i>G. nubicola</i> (unc)	<i>Pseudowintera colorata</i>
<i>G. sp.</i> (unnamed; included in <i>G.</i> <i>depressa</i> , as var. <i>novae-</i> <i>zealandiae</i> , by Franklin 1962)	

## DICOT LIANES AND RELATED

### TRAINING PLANTS

*Metrosideros fulgens*  
*Muehlenbeckia axillaris* (unc)  
*Rubus cissoides* var. (leaflets  
broader than in var. *cissoides*)

## PSILOPSIDS AND LYCOPODS

*Tmesipteris tannensis*  
*Lycopodium australianum*  
*L. fastigiatum*  
*L. scariosum*  
*L. variuum* (incl. *L. billardierei*  
and *L. novae-zelandicum*)  
*L. volubile*

## FERNS

*Asplenium flaccidum*  
*Blechnum discolor*  
*B. fluviatile* agg.  
*B. penna-marina*  
*B. procerum*  
*B. sp.(i)* (unnamed) (common lowland  
sp.; lower pinnae reduced in length)  
*B. sp. (ii)* (unnamed) (mount-  
ain sp.; lower pinnae not reduced)  
("The Dunedin-Cass Variety" of  
Crookes 1963)  
*B. procerum* X *B. sp. (ii)* (unc)

*Ctenopteris heterophylla*  
*Cyathea colensoi*  
*Dicksonia squarrosa*  
*Gleichenia dicarpa*

*Grammitis billardierei*  
*G. givenii* (unc)  
*G. magellanica* spp. *nothofageti*  
*G. poeppigiana*  
*Histeropteris incisa*  
*Hymenophyllum armstrongii*  
*H. bivalve*  
*H. demissum*  
*H. dilatatum*  
*H. ferrugineum*  
*H. flabellatum*  
*H. lyallii* (unc)  
*H. minimum* (unc)  
*H. multifidum*  
*H. peltatum* (unc)  
*H. revolutum*  
*H. rufescens*  
*H. villosum*  
*Hypolepis distans* (unc)  
*H. millefolium*  
*H. rufobracteata* (unc)  
*Leptopteris superba*  
*Paesia scabrella*  
*Polystichum cystostegia* (unc)  
*P. vestitum*  
*Pteridium esculentum*  
*Schizaea australis* (unc)  
*Trichomanes endlicherianum*  
*T. elongatum*  
*T. tentiforme*

## ORCHIDS

*Adenochilus gracilis* (unc)  
*Aporostylis bifolia*  
*Chiloglottis cornuta*  
*Corybas acuminatus* (unc)  
*C. macranthus* (unc)

*C. thilobus* s.s.  
*Cyrtostylis reniformis* s.s.  
*Dendrobium cunninghamii*  
*Earia autumnalis*  
*E. mucronata* s.s.  
*Lyperanthus antarcticus*  
*Prasophyllum colensoi*  
*Pterostylis ifsoniana* (unc)  
*P. venosa*  
*Thelymitra cyanea*  
*T. hatchii* (unc)  
*T. longifolia* (unc)

**GRASSES** *T. pulchella* (unc)

*Agrostis dyeri* s.s.

*A. personata*

*A. sp.* (unnamed; aff. *A. dyeri*)  
(panicle not contracted as in *A. dyeri*)

*Chionochloa australis*

*C. conspicua* ssp. *conspicua*

*C. pallens* ssp. *pallens*

*C. rubra* ssp. *occulta*

*C. sp.* (unnamed) (*C. flavesans*  
auct. South Island, non *C. flavesans* Zотов).

*C. australis* × *C. pallens* (unc)

*C. pallens* × *C. rubra*

*Deschampsia tenella*

*Deyeuxia aucklandica*

*Hierochloe novae-zelandiae*

*Microlaena avenacea*

*M. colensoi* s.s.

*Poa breviglumis*  
*Poa celsa*  
*P. colensoi*

*P. maia*

*P. novae-zelandiae* s.s.

*P. sp.* (unnamed; aff. *P. novae-zelandiae*) (lvs much narrower  
than in *P. novae-zelandiae*) (unc)

*Rytidosperma gracile*

*R. nigricans*

*R. setifolium*

*Trisetum* sp. (unnamed) (plants  
slender; lf sheath glabrous or  
nearly so; lf blade flat, up to  
2.5 mm wide, margins smooth,  
vs. with a few hairs near ligule)  
(unc)

**SEDGES**

*Carex aciculatis*

*C. cockayneana* (unc)

*C. dallic* (unc)

*C. dissita* s.s.

*C. gaudichaudiana* (unc)

*C. libera*

*C. pyrenaica* var. *cephalotes*  
(unc)

*C. sp. (I)* (unnamed; aff. *C. coriacea*)

*C. sp. (II)* (unnamed; aff. *C. dissita*)

## *Carpha alpina* s.s.

C. sp. (unnamed) (plants "taller and more slender" than in *C. alpina*, with "narrower lvs"; see Edgar 1970, p. 200) (unc)

*Grahnia proceria*

*Isolepis aucklandica*

*I. caligenis*

*I. habra*

*Oreobolus impar*

*O. pectinatus*

*O. strictus*

*Schoenus pauciflorus* s.s.<sup>†</sup>

*Uncinia astonii*

*U. clavata*

*U. divaricata*

*U. filiformis*

*U. fuscovaginata* var. (plants glaucous)

*U. gracilenta*

*U. involuta*

*U. longifluctos*

*U. nervosa* s.s.

*U. rupestris* (incl. *U. angustifolia*)

*U. uncinata*

*U. viridis*

*U. zotovii*

*U. sp.* (unnamed; aff. *U. nervosa*)

(more slender than *U. nervosa*; see Edgar 1970, p. 231)

## RUSHES AND ALLIED PLANTS

*Centrolepis ciliata*

*C. pallida* (unc)

*Empodium minus*

*Grimardia setacea*

*Juncus antarcticus*

*J. novae-zelandiae*

*J. pusillus*

*Luzula colensoi*

*L. migrata*

*L. subclavata* (unc)

*L. sp.* (unnamed; included in *L. picta*, as var. *limosa*, by Edgar 1966)

*L. colensoi* × *L. migrata* (unc)

## MONOCOT HERBS OTHER THAN

### ORCHIDS, GRASSES, SEDGES, RUSHES

*Astelia linearis* (incl. *A. l.*

var. *novae-zelandiae*)

*A. nivicola* s.s.

*A. petriei*

*A. skottsbergii*

*A. sp.* (i) (unnamed; included in

*A. nivicola*, as var. *moiceae*, by Moote 1966)

*A. sp.* (ii) (unnamed; aff. *A. nervosa*)

(lvs broad, not conspicuously white adaxially as in *A. nervosa*)

*Dianella nigra*

*Herpolitha novae-zelandiae*

*Libertia peragrinans* (unc)

*L. pulchella*

*Luzuriaga parviflora*

*Phormium cookianum*

† ("slender form" of Edgar 1970, p. 194)

## COMPOSITE HERBS

*Abrotanella caespitosa*

*A. linearis* s.s.

*A. pusilla*

*Anaphalis* sp. (*Helichrysum bellidioides* s.s.)

*Brachyglottis* sp. (unnamed; included in *B. bellidioides*, as var. *crassa*, by Nordenstam 1978)

*Celmisia alpina* s.s.

*C. argentea* (unc)

*C. armstrongii*

*C. dallii*

*C. gibbsii*

*C. incana* (incl. *C. discolor*)

*C. laricifolia*

*C. lateralis* (incl. *C. l.* var. *villosa*)  
(unc)

*C. monroi* ssp. (*C. coriacea* var. *lancifolia*)

*C. sessiliflora*

*C.* sp. (unnamed; aff. *C. gracilenta*)

(tufted; lvs linear to linear-lanceolate, up to 45cm x 1-2cm; uppersurface and margins thinly covered in tomentum; lower surface densely covered)

*Calpina* x *C.* sp. (undetermined) (unc)

*Craspedia* sp. (unnamed) (plants

summer-green; lvs green; both surfaces clad in multicellular hairs; margins faintly to distinctly white with cottony hairs; fls white)

*Dolichoglottis scorzoneraoides*

*Gnaphalium avdax*

*G. involucratum* (unc)

*G. laterale* (unc)

*G. limosum* s.s.

*G. sphaericum* (unc)

*Lagenifera concata* (unc)

*L. pumila* (unc)

*L. strangulata*

*Leucogenes grandiceps* (unc)

*Raoulia eximia* var.

*R. glabra* (unc)

*R. grandiflora* var. *grandiflora*

*Senecio minimus*

*S. wairauensis*

## DICOT HERBS OTHER THAN COMPOSITES

*Acaena ansetinifolia*

*A. profundeincisa* (unc)

*Aciphylla anomala*

*A. colensoi* var. (*A. indurata*)

*A. hookeri*

*A. polita* var.

*A. colensoi* x *A. hookeri*

*Actinotis novaezelandiae*

- Anisotome aromatica*
- A. haastii*
- A. sp.(i)* (unnamed; included in *A. imbricata*, as var. *prostrata*, by Dawson 1961)
- A. sp.(ii)* (unnamed) (mat-forming; infls up to 5 cm tall)
- Cardamine sp.(i)* (*C. debilis* agg.)  
(cf. "Long Style" of Pritchard 1957)
- C. sp.(ii)* (*C. debilis* agg.) ("Glossy Leaf" of Pritchard 1957?)
- Colobanthus affinis* (unc)
- Donatia novae-zelandiae*
- Drosera arcturi*
- D. spathulata*
- D. stenopetala*
- Elatine gratioloides* (unc)
- Epilobium brunnescens* s.s.
- E. cockayneum*
- E. parnitens*
- E. sp.* (unnamed; included in *E. brunnescens*, as ssp. *minutiflorum*, by Raven & Engelhorn, 1971)
- Euphrasia revoluta* (unc)
- E. townsonii* s.s.
- Forstera mackayi*
- F. tenella*
- F. sp.* (unnamed; aff. *F. sedifolia*)
- F. sp.* (unnamed) (not seen in field; specimens from Glasgow Ra. in CHR)
- Gentiana spenceri*
- G. townsonii*
- G. sp.(i)* (unnamed; included in *G. bellidifolia*, as var. *australis*, by Cheeseman 1925)
- G. sp.(ii)* (unnamed; aff. *G. montana*)  
(Stems tall, robust, often decumbent at base; basal lvs densely rosulate, in adult plants vs. placed well above ground level; infl umbelliform)
- Geranium sp.* (unnamed; aff. *G. microphyllum*)
- Geum uniflorum*
- Gonocarpus aggregatus*
- Gunnera monoica*
- Hydrocotyle heteromeria* (unc)
- H. microphylla* (unc)
- Kelleria dieffenbachii*
- K. laxa* (unc)
- K. multiflorus* (unc)
- Mitrasacme novae-zelandiae*
- Myosotis forsteri* (unc)
- M. lyallii* (incl. *M. l.* var. *townsonii* and *M. elderi*) (unc)
- Myriophyllum pedunculatum*  
ssp *novae-zelandiae* (unc)
- Nasturtia balfouriana*
- N. ciliata*
- N. dapressa* (incl. *N. cunninghamii*)
- N. scapanioides* (unc)

*N. sp.* (unnamed; aff. *N. dichondrifolia*) (hairs present on both surfaces of lf, straight, not hooked as in *N. dichondrifolia*)

*Oreomyrtus colensoi* s.s. (incl. *O.c.* var. *hispida* and *O.c.* var. *multifida*)

*Oreostylidium subulatum*

*Ourisia lactea* ssp. *lactea*

*O. simpsonii*

*Oxalis magellanaica*

*Parathebe lyallii* (unc)

*Phyllachne colensoi* var.

*Plantago lanigera* s.l. (incl. *P.n.z.*)

*P.uniflora*

*Pratia angulata* agg.

*Psychrophila novae-zelandiae*

*Ranunculus insignis* (unc)

*R.membranifolius*

*Schizoclema haastii* s.s.

*S. nitens* (unc)

*Stellaria decipiens* (incl. *S.parviflora*)

*Urtica incisa* (unc)

*Viola cunninghamii*

*V.filicaulis*

#### RECORDS NOT ACCEPTED

*Brachygottis bidwillii* × *Olearia arborescens*(?) (Cochayne 1928, p. 316, as *Senecio bidwillii* × *O. arborescens*)

? *Hebe odora* (Cochayne 1928, p. 316, as *H.buxifolia*)

*H.vernicosa* (Cochayne 1928, p. 316)

*H.odora* × *H.macrantha* (Cochayne 1928, 316, as *H.buxifolia* × *H.macrantha*)

*H.odora* × *H.vernicosa* (Cochayne 1928, p. 316, as *H.buxifolia* × *H.vernicosa*)

*Gentiana montana* (Cochayne 1928, p. 316)

*Ourisia macrophylla* (Cochayne 1928, p. 316)

*Hebe carnosula* (Townson 1907, as *Veronica carnosula*)

#### ADVENTIVE PLANTS

*Agrostis capillaris* (unc)

*Mycelis muralis* (unc)

No. of indigenous spp.: 329.

No. of adventive spp.: 2.

10

## REFERENCES

- Allan, H.H. 1961: "Flora of New Zealand." Govt. Printer, Wgt'n. Vol. 1. 1085 pp.
- Cheeseman, T.F. 1925: "Manual of the N.Z. Flora." Govt. Printer, Wgt'n. 1163 pp.
- Crookes, M. 1963: "New Zealand Ferns." Ed. 6. Whitcombe & Tombs. 408 pp.
- Dawson, J.W. 1961: A revision of the genus Anisotome (Umbelliferae). Univ. Calif. Public Bot. 33: 1-98.
- Edgar, E. 1966: Luzula in New Zealand. N.Z. Jl Bot. 4: 159-84.
- 1970: In "Flora of New Zealand." Govt. Printer, Wgt'n. Vol. 2. 354 pp.
- Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans R. Soc. N.Z. Bot. 1: 155-73.
- Cochrane, L. 1928: "The Vegetation of New Zealand". Engelmann, Leipzig.
- Moore, L.B. 1961: In "Flora of New Zealand." Government Printer, Wellington. Vol. 1. 1085 pp.
- Moore, L.B. 1966: Australasian asteliads (Liliaceae). N.Z. Jl Bot. 4: 201-40.
- Pritchard, G.G. 1957: Experimental taxonomic studies on species of Cardamine Linn. in New Zealand. Trans. R. Soc. N.Z. 85: 75-89.
- Raven, P.H.; Engelhorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. Jl Bot. 9: 345-50.
- Townson, W. 1907: On the vegetation of the Westport District. Trans. N.Z. Inst. 39: 380-433.
- Wardle, P. 1968: The taxonomy and distribution of the stipitate species of Pseudopanax in New Zealand. N.Z. Jl Bot. 6: 226-36.
- Nordenstam, B. 1978: Opera Bot. 44: 1-84.