

HIGHER PLANTS (PSILOPSIDS, LYCOPODS, FERNS, GYMNOSPERMS,  
FLOWERING PLANTS) OF EDITOR HILL<sup>1</sup> AND LOOKOUT PK<sup>2</sup>,  
MARLBOROUGH, FROM OPOURI SADDLE TO SUMMITS  
(480-1032 M)

<sup>1</sup> Grid ref. NZMS 260 027 675084

→ <sup>2</sup> Grid ref. " " P27 713073


<sup>3</sup> Grid ref. " " P27 723059

A.P.Druce, Wellington, and W.D.Burke, Nelson,  
Nov. 1985

Additional records by G.M.Rogers, Wellington, and A.P.D.,  
1986; and by A.P.D., 1989 and 1990)  
(Based on visits Nov. 1985, April 1986, March 1989,  
April 1990)

(Revised June 1990, July 1991, Oct. 1992)

Rocks: graywacke and argillite  
(with small areas of volc. argillite,  
pillow lavas, and conglomerate on  
Editor Hill)



(unc) - uncommon

(s) - seen only near Saddle

### INDIGENOUS PLANTS

#### GYMNOSPERM TREES AND SHRUBS

- Dacrydium cupressinum* (s)
- Libocedrus bidwillii*
- Phyllocladus alpinus* agg.
- Podocarpus hallii*
- P. nivalis* (unc)
- P. acutifolius* × *P. nivalis* (unc)
- P. hallii* × *P. nivalis* (unc)
- Prumnopitys ferruginea* (unc)

#### MONOCOT TREES AND SHRUBS

- Cordyline banksii* (s)

#### DICOT TREES

- Aristotelia serrata* (s)
- Carpodetus serratus*
- Elaeocarpus dentatus* (incl.  
*E. d. var. obovatus*) (s)
- Griselinia littoralis*

- Hedycarya arborea* (juv.) (s)
- Metrosideros umbellata*
- Myrsine australis*
- Nothofagus fusca*
- N. menziesii*
- N. solandri* var. *cliffortioides*
- N. truncata*
- Olearia rani* var. *colorata*
- Pseudopanax colensoi* ss.  
(adult plants uncommon,  
epiphytic only)
- P. crassifolius*
- P. edgerleyi* (unc)
- P. simplex* (incl. *P. s. var. sinclairii*) (juv.) (unc)
- Weinmannia racemosa* val-  
*racemosa*

**DICOT SHRUBS:**

*Brachyglottis rotundifolia* var.  
(*B. elaeagnifolia*)

*Cassinia leptophylla* var. *lept.* (unc)

*C. vauvilliersii* (unc)

*Coprosma colensoi* s.s.

*C. foetidissima*

*C. grandifolia* (s)

*C. lucida* s.s. (unc)

*C. microcarpa*

*C. pseudocuneata* ssp.

*C. rhinoides* (incl. *C. polymorpha*)

*C. robusta* (s)

*C. sp.(i)* (unnamed; included in

*C. parviflora* by Oliver, 1935,

and others) (If undersurface

not minutely hairy as is usual

in *C. parviflora*; drupes vs. un-

pigmented, occ. purplish red)

*C. sp.(ii)* (unnamed; aff. *C. colensoi*) *Ripogonum scandens* (s)  
(low-growing shrubs with trailing  
branches) (unc)

*depressa*, as var. *novae-zelandiae*, by Franklin (1962)

*Hebe rigidula* var.

*H. vernicosa*

*Leptospermum scoparium*  
(unc)

*Leucopogon fasciculatus* var.

*Myrsine divaricata* s.s.

*Pentachondra pumila*

*Peraxilla colensoi* (unc)

*Pimelea longifolia* var.  
*longifolia*

*Pittosporum rigidum* var. (*P.*  
*crassicaule*)

*Pseudopanax anomalus*

*Pseudowintera axillaris* (s)

*P. colorata*

**MONOCOT LIANES**

*C. sp.(iii)* (unnamed) (*C. parviflora* *Clematis paniculata* (s)  
var. *dumosa* sensu Allan 1961, *Metrosideros diffusa*  
non *C. p.* var. *dumosa* Cheeseman (unc)) *Muehlenbeckia australis*

*Cyathodes* sp. (*C. juniperina* *Rubus cissoides* var. (leaf-  
agg.) (lvs vs. <1cm x <1mm) lfts broader than in var.

*Dracophyllum longifolium* var. *cissoides* (unc)

*Gaultheria antipoda*

*G. sp.* (unnamed; included in *G.*

**DICOT LIANES**

## PSILOPSIDS AND LYCOPODS

*Tmesipteris elongata* (incl. *T.e. ssp. robusta*) (unc)  
*T. tannensis*  
*Lycopodium australe* (unc)  
*L. fastigiatum* (unc)  
*L. scariosum* (unc)  
*L. variuum* (incl. *L. billardierei* and *L. novae-zelandicum*) (unc)  
*L. volubile*

*Histiopteris incisa*

*Hymenophyllum armstrongii* (unc)  
*H. bivalve*

*H. demissum*

*H. dilatatum*

*H. flabellatum*

*H. malingii* (unc)

*H. multifidum*

*H. latum*

*H. revolutum*

*H. sanguinolentum* agg.

*H. villosum*

*Hypolepis rufobarbata* (unc)

*Leptopteris hymenophylloides*

*Poa scaberula*

*Phymatosorus pustulatus* (unc)

*Polystichum vestitum* (unc)

*Pteridium esculentum* (S)

*Ryrosia eleagnifolia*

*Rumohra adiantiformis*

*Sticherus cunninghamii* (unc)

*Trichomanes teniforme*

## FERNS

*Asplenium flaccidum*  
*Blechnum chambersii* (unc)  
*B. discolor*  
*B. fluviatile* agg.  
*B. minus* (unc)  
*B. penna-marina* (unc)  
*B. procerum*  
*B. sp. (unnamed)* (mountain sp.; lower pinnae not reduced in length; "The Dunedin-Cass Variety" of Crookes 1963)  
*Ctenopteris heterophylla*  
*Cyathea colensoi*  
*C. dealbata*  
*C. medullaris* (S)  
*C. smithii*  
*Dicksonia squarrosa* (unc)  
*Grammitis billardierei*  
*G. givenii* (unc)  
*G. magellanica* ssp. *nothofageti*

*Adenochilus gracilis* (unc)

*Aporostylis bifolia* (unc)

*Chiloglottis cornuta*

*Cotybas trilobus* ss.

*Dendrobium cunninghamii*

*Earina autumnalis*

*E. mucronata* ss.

*Pterostylis montana* ss. (S)

*P. venosa* (unc)

*Thelymitra longifolia* (unc)

## GRASSES

*Agrostis personata* (unc)  
*A.* sp. (unnamed; aff. *A. dyeri*)  
(panicle spreading, not contracted  
as in *A. dyeri*)

*Chionochloa cheesemanii*  
*C.* sp. (undetermined)

## RUSHES

*Juncus antarcticus* (unc)  
*J. novae-zelandiae* (unc)  
*J. pallidus* (unc) (s)  
*Luzula* sp. (unnamed; included  
in *L. picta*, as var. *limosa*, by  
Edgar 1966) (unc)

## MONOCOT HERBS OTHER THAN

### ORCHIDS, GRASSES, SEDGES, RUSHES

*Microlaena avenacea* (unc)  
*Rytidosperma gracile*

## SEDGES

*Carex dissita* s.s.  
*Gahnia pauciflora* (s)  
*G. procera* (unc)  
*G. setifolia* (s)  
*Isolepis habra* (unc)  
*Oreobolus pectinatus* (unc)  
*Schoenus maschalinus* (unc)  
*S. pauciflorus* s.s. (unc)  
*Uncinia astonii* (unc)  
*U. clavata* (unc)  
*U. filiformis*  
*U. fuscovaginata* var. (glaucous) (unc)  
*U. gracilenta* (unc)  
*U. nervosa* s.s. (unc)  
*U. rupestris* (incl. *U. angustifolia*)  
*U. silvestris* (incl. *U. affinis*)  
*U. uncinata*  
*U. zotovii*

*Astelia solandri*

*A.* sp. (i) (unnamed; aff. *A. nervosa*)  
(lvs broader than in *A. nervosa*,  
not conspicuously white ad-  
axially as in *A. nervosa*)

*A.* sp. (ii) (unnamed; aff. *A. nervosa*)  
(lvs narrower than in *A. nervosa*)

*Dianella nigra* (s)

*Libertia grandiflora* (s)

*L. pulchella*

*Luzuriaga parviflora*

*Phormium cookianum*

## COMPOSITE HERBS

*Anaphalis* sp. (*Helichrysum bellidioides* s.s.) (unc)  
*Brachyglottis* sp. (unnamed)  
*Calmisia gracilenta* var.  
*C. hieracifolia* var.  
*C.* sp. (unnamed; aff. *C. gracilenta*)  
(plants strongly rhizomatous) (unc)  
*Gnaphalium delicatum* (unc)  
*Helichrysum filicaule* (unc)

Lagenifera pumila (unc)  
Raoulia glabra (unc)  
Senecio minimus (unc)

Hydrocotyle moschata (unc)  
Nettlera depressa (incl. N. cunn-  
inghamii)  
Viola filicaulis

### DICOT HERBS OTHER THAN COMPOSITES

#### COMPOSITES

Acaena austro-nigra (unc)  
Epilobium pernitens (unc)  
E. sp. (unnamed; included in E.  
brunneum, as ssp. minutum,  
florum, by Raven and Engel-  
horn, 1971) (unc)  
Gonocarpus micranthus ssp.  
micranthus (unc)  
G. montanus (unc).

No. of indigenous spp.: 169.

### ADVENTIVE PLANTS

#### COMPOSITE HERBS

Hypochoeris radicata (unc)  
Leontodon taraxacoides (unc)

#### GRASSES

Agrostis capillaris (unc)

### DICOT HERBS OTHER THAN COMPOSITES

Sagina procumbens (unc)

No. of adventive spp.: 4.

### REFERENCES

- Franklin, D.A. 1962: Trans Roy. Soc. N.Z. Bot. 1: 155-73.  
Allan, H.H. 1961: "Flora of N.Z." Govt. Printer, Wellington. Vol. 1. 1085 pp.  
Crookes, M. 1963: "N.Z. Flora". Ed. 6. Whitcombe & Tombs. 408 pp.  
Edgar, E. 1966. Luzula in N.Z. N.Z. J. Bot. 4: 159-84.  
Oliver, W.R.B. 1935: The genus Coprosma. Bishop Mus. Bull. 132. 207 pp.  
Raven, P.H.; Engelhorn, T. 1971: New taxa and new combinations in Australasian  
Epilobium (Onagraceae). N.Z. J. Bot. 9: 345-50.