

(289)


INDIGENOUS VASCULAR PLANTS (LYCOPODS, FERNS, GYMNO-
SPERMS, FLOWERING PLANTS) OF MT STORMY[†], STORMY
RIDGE, NORTH-WEST NELSON, 500-1084 M

A.P. Druce, Wellington, April 1987

(revised Oct. 1987, Oct. 1988, July 1990, July 1992)

(Based on visit January 1987)

[†] Grid ref. L27 A56912



(unc) - uncommon

GYMNOSPERM TREES AND SHRUBS

Dacrydium cupressinum (juv.)
Halocarpus biformis
Lepidothamnus laxifolius
L.intermedius × *L.laxifolius*
Phyllocladus sp. (aff. *P.alpinus*)
Podocarpus hallii

Archeria traversii (incl. A.t. var.
australis)

Brachyglottis rotundifolia var.
(incl. *B.buchananii* and *B.cock-
aynei*)

Coprosma colensoi ss.

C.ctenulata

C.foetidissima

C.pseudocuneata ssp.

C.thamnoides (incl. *C.polymorpha*)

C.sp. (unnamed; aff. *C.coldensoi*)

(low-growing shrub with
trailing branches)

Cyathodes dealbata (incl. *C.pumila*)

C.empetrifolia

C.sp. (*C.juniperina* ass) (vs vs.
7cm x 7mm, up to 1.5cm x 2.5mm)

Dracophyllum longifolium var.
(incl. *D.filifolium* & *D.oliveti*)

D.palustre

D.politum (incl. *D.densum*)

Pseudopanax crassifolius (unc) *Gaultheria antipoda*

P.linearis

G.rupestris (incl. *G.subcordym-*

P.simplex (incl. *P.s.var.sinclairii*) *bosa*)

P.sp. (unnamed; included in *G.sp.* (unnamed; included in

P.colensoi, as var. *fiordensis* *G.depressa*, as var. *nova-
zelandiae*, by Franklin 1962)

and as *P.ternatus*, by

Wardle 1968)

G.sp. (unnamed) X *G.*

macrostigma

Hebe sp. (*Leonotis* *mooreae*)

Weinmannia racemosa var.
racemosa

DICOT SHRUBS

Alseuosmia pusilla (unc)

Leptospermum scoparium var.
Leucopogon fasciculatus var.
Myrsine divaricata s.s.
Olearia colensoi var. *colensoi*
Pentachondra pumila
Pittosporum rigidum var. (*P. crassifolium*)
Pseudowintera colorata

DICOT LIANES

Metrosideros fulgens

PSILOPSIDS AND LYCOPODS

Tmesipteris tannensis
Lycopodium fastigiatum
L. scariosum

FERNS

Asplenium flaccidum
Blechnum discolor
B. fluviatile agg.
B. proctatum

Cyathea colensoi
C. smithii
Dicksonia squarrosa
Grimmitis billardierei
Histiopteris incisa
Hymenophyllum armstrongii
H. bivalve
H. demissum
H. dilatatum
H. flabellatum
H. lyallii
H. multifidum
H. ratum
H. revolutum
H. rufescens

Lindsaea trichomanoides

Phymatosorus pustulatus
Trichomanes reniforme

ORCHIDS

Aporostylis bifolia
Eatina autumnalis
Hyperanthus antarcticus
Prasophyllum colensoi
Pterostylis banksii
P. venosa
Thelymitra sp.

GRASSES

Chionochloa australis
C. rubra ssp. *occulta*
Deyauxia aucklandica
Hierochloe novae-zelandiae
Microlaena avenacea
M. thomsonii
Rytidosperma nigricans

SEDGES

Carex aciculata
Carpha alpina s.s.

Isolepis aucklandica
Oreobolus impar
O. strictus
Schoenus pauciflorus ss. (= "slender form" of Edgar 1970, p. 194)
Uncinia astonii
U. filiformis
U. involuta
U. rupestris (incl. *U. angustifolia*)
U. uncinata
U. zotovii

*U.sp.(unnamed; aff. *U. nervosa*)*
*(more slender than *U. nervosa*;*
see Edgar 1970, p. 23)

RUSHES

Empodium minus
Luzula sp. (undetermined)

MONOCOT HERBS OTHER THAN

ORCHIDS, GRASSES, SEDGES, RUSHES

Astelia linearis var. *linearis*
A. trinervia
A. sp.(i)(unnamed; included in
**A. nivicola*, as var. *nivicolae*,*
by Moore (1966)
*A. sp.(ii)(unnamed; aff. *A. nervosa*)*
(lvs broad, not conspicuously
*white adaxially as in *A.**
**nervosa*)*

Heteropliton novae-zelandiae
Libertia pulchella
Luzuriaga parviflora
Phormium cookianum

COMPOSITE HERBS

Abrotanella linearis ss.
Anaphalis keniensis ss.
Brachyglottis sp.

DICOT HERBS OTHER THAN

COMPOSITES

Actinotis novae-zelandiae
Anisotome aromatica
Donatia novae-zelandiae
Drosera arcturi
D. spathulata
D. stenopetala
Euphrasia townsonii ss.
Forstera mackayi
Gentiana gracilifolia (incl.
G. vernicosa)
G. montana (incl.
G.m. var. stolonifera, G. patula,
and G. bellidifolia var. pulchella)
G. spencii
Kelleria dieffenbachii
*Nertera sp. (unnamed; aff. *N.**
dichondrifolia) (hairs present
on both surfaces of lf, straight,
*not hooked as in *N. dichondri-**
**folia*)*
Oreostylidium subulatum
Phyllachne colensoi var.
Viola filicaulis

No. of species: 135.

Records not accepted

Calmisia armstrongii
C. dallii
C. incana (incl. *C. discolor*)
C. parva
(*C. armstrongii* X *C. parva*) (unc)
C. similis (unc)
Raoulia grandiflora var.
grandiflora

Hebe odora (Wardle & Buxton, 1985)
[= *H. sp. (unnamed)*]
Uncinia viridis (Wardle & Buxton,
1985) [= *U. involuta*?]
Euphrasia cockayneana (Wardle
& Buxton, 1985) [= *E. townsonii*]

REFERENCES

- Edgar, E. 1970: In "Flora of New Zealand." Government Printer, Wellington. Vol. 2. 354 pp.
- Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans. Roy. Soc. N.Z. Bot. 1:155-73.
- Moore, L.B. 1961: In "Flora of New Zealand." Government Printer, Wellington. Vol. 1. 1085 pp.
- 1966: Australasian asterads (Liliaceae). N.Z.J. Bot. 4: 201-40.
- Wardle, P. 1968: The taxonomy and distribution of the stipulate species of Pseudopanax in New Zealand. N.Z.J. Bot. 6: 226-36.
- Wardle, P.; Buxton, R.P. 1985: List of vascular plants recorded from the Karamea district. Unpublished report, Botany Division, D.S.I.R. 29 pp.
- —

