

HIGHER PLANTS (PSILOPSIDS, LYCOPODS, FERNS, GYMNOSPERMS,
FLOWERING PLANTS) OF WALKER RIDGE, N.W. NELSON,
150-1200 M (500-3960 FT)


†Highest point at grid ref. S8 098807

A.P.Druce, Wellington, and S.Courtney, Dept. of
Conservation, Nelson, Oct. 1987

(revised July 1988, Aug. 1990, July 1991, July 1992,

Based on visits Jan. 1987 and Oct. 1987 Aug 1993)

Rock: marble



INDIGENOUS PLANTS

(unc) - uncommon

GYMNOSPERM TREES AND SHRUBS

- Dactylium cypressinum*
Halocarpus biformis
Lagarostrobos colensoi (unc)
Lepidothamnus intermedius
Libocedrus bidwillii
Phyllocladus sp. (aff. *P. alpinus*)
Podocarpus hallii
Prumnopitys ferruginea
P. taxifolia (unc)

MONOCOT TREES AND SHRUBS

- Cordyline banksii*
C. indivisa (unc)
Rhopalostylis sapida

DICOT TREES AND SHRUBS

- Alectryon excelsum* var. *excelsum* (unc)
Alseuosmia macrophylla (unc)
Archetia traversii (incl. *A. var. australis*)
Aristotelia serrata (unc)
Ascarina lucida (unc)
Brachyglottis hectorii (unc)
B. repanda
B. rotundifolia var. (incl. *B. buchananii* & *B. cockaynei*) (unc)
Carmichaelia sp. (subgenus *Carmichaelia*)
Carpodetus sematus
Cassinia laevis

- Coprosma ciliata* csp. *ciliata* (unc)
C. colensoi s.s.
C. foetidissima
C. grandifolia
C. linariifolia
C. lucida s.s.
C. microcarpa

- C. obconica*
C. propinqua ssp. *propinqua*
C. pseudocuneata ssp.
C. thamnoides (incl. *C. polymorpha*)
C. robusta
C. sp. (unnamed; included in *C. parviflora* by Oliver, 1935, and others) (If undersurface not minutely hairy as is usual in *C. parviflora*; drupes vs. unpigmented, occ. purplish red)
C. propinqua X *C. robusta* (unc)
Coraria arborea (unc)
Cyathodes sempervirens (unc)
C. sp. (*C. juniperina* agg.) (vs small, vs. \angle (cm) x \angle (mm))
Dracophyllum longifolium var. (*D. utileanum*)
D. traversii (incl. *D. pyramidale*)
Elaeocarpus dentatus (juv. only)
E. hookerianus
Gaultheria antipoda (unc)
G. sp. (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962)
G. antipoda X *G. sp.* (unnamed) (unc)
Griselinia littoralis
Hebe gracilis
Hedycarya arborea
Melichrysum aggregatum var. *aggregatum*

<i>Leucopogon fasciculatus</i> var.	<i>Pseudowintera axillaris</i>
<i>Haberla populnea</i> var. (<i>H. ovata</i>)	<i>P. colorata</i>
<i>Kunzea ericoides</i> agg.	<i>Quintinia serrata</i> (incl. <i>Q. acutifolia</i> and <i>Q. elliptica</i>)
<i>Laurilia novae-zelandiae</i>	<i>Sophora microphylla</i> (unc)
<i>Leptospermum scoparium</i>	<i>Streblus heterophyllus</i> (unc)
<i>Lophomyrtus obcordata</i> (unc)	<i>Urtica ferox</i> (unc)
<i>Macropiper excelsum</i> var. <i>excelsum</i>	<i>Wainmannia racemosa</i> var.
<i>Malope simplex</i>	<i>racemosa</i>
<i>Malicytus obovatus</i> var. <i>obovatus</i>	
<i>M. ramiflorus</i> ssp. <i>ramiflorus</i>	
<i>Metrosideros umbellata</i>	
<i>Myrsine australis</i>	MONOCOT LIANES
<i>M. divaricata</i> s.s.	<i>Freylinia banksii</i>
<i>M. salicina</i>	<i>Ripogonum scandens</i>
<i>Neomyrtus pedunculata</i>	
<i>Nothofagus fusca</i>	DICOT LIANES
<i>N. menziesii</i>	<i>Clematis foetida</i> (unc)
<i>N. solandri</i>	<i>C. paniculata</i>
<i>N. truncata</i>	<i>Metrosideros colensoi</i> (incl. <i>M. c.</i> var. <i>pendens</i>)
<i>Olearia arborescens</i> (unc)	<i>M. diffusa</i>
<i>O. avicinaria</i>	<i>M. fulgens</i>
<i>O. lacunosa</i> var. <i>lacunosa</i> (unc)	<i>M. perforata</i>
<i>O. tani</i> var. <i>colorata</i>	<i>Muehlenbeckia australis</i> (unc)
<i>Pennantia corymbosa</i> (unc)	<i>Parsonsia heterophylla</i>
<i>Pittosporum eugenioides</i> (unc)	<i>Rubus cissoides</i> var. (leaflets broader than in var. <i>cissoides</i>)
<i>P. rigidum</i> var. (<i>P. crassicaule</i>)	<i>R. schmideliioides</i> var.
<i>P. tenuifolium</i>	
<i>Pseudopanax anomalous</i>	
<i>P. colensoi</i> s.s.	PSILOPSIDS AND LYCOPODS
<i>P. crassifolius</i>	<i>Tmesipteris elongata</i> (incl. <i>T. e.</i> ssp. <i>robusta</i>)
<i>P. linearis</i>	<i>T. tannensis</i>
<i>P. macintyrei</i>	<i>Lycopodium fastigiatum</i>
<i>P. simplex</i> (incl. <i>P. s.</i> var. <i>sinclairii</i>)	<i>L. variuum</i> (incl. <i>L. billardierei</i> and <i>L. novae-zelandicum</i>)
<i>P. sp. (unnamed; included in <i>P. colensoi</i>, as var. <i>fiordensis</i> and as var. <i>ternatus</i>, by Wardle 1968)</i>	<i>L. volubile</i> (unc)

FERNS

- Adiantum cunninghamii*
Anarhitopteris lanceolata
Asplenium bulbiferum (unc)
A. flaccidum
A. hookerianum
A. lyallii
A. oblongifolium (unc)
A. polyodon (unc)
A. sp. (unnamed; aff. *A. trichomanes*) (unc)
Blechnum chambersii
B. discolor
B. fluviatile agg.
B. fraseri
B. membranaceum (unc)
B. penna-marina (unc)
B. procerum
B. vulcanicum (unc)
B. sp. (unnamed) (common
lowland sp.; lower pinnae re-
duced in length)
- Ctenopteris heterophylla*
Cyathea colensoi
C. dealbata
C. medullaris
C. smithii
Dicksonia squarrosa
Grammitis billardierei
G. ciliata
G. magellanica ssp. *nothofageti*
G. patagonica (unc)
Histeropteris incisa
Hymenophyllum bivalve
H. demissum
H. dilatatum
- H. farrugineum* (unc)
H. flabellatum
H. malinii (unc)
H. multifidum
H. pulcherrimum (unc)
H. ratum (unc)
H. rufescens (unc)
H. sanguinolentum agg.
H. villosum
H. sp. (unnamed; aff. *H. flexuosum*) (unc)
Hypolepis ambigua (unc)
Lastreopsis glabella (unc)
L. hispida (unc)
Leptolepia novae-zelandiae
Leptopteris hymenophylloides
L. superba
L. hymenophylloides × *L. superba* (unc)
Lindsaea trichomanoides
Paesia scabrella (unc)
Pellaea rotundifolia (unc)
Phymatosorus postulatus
P. scandens
Pneumatopteris pennigera
Polystichum richardii (unc)
P. vestitum (unc)
Pteridium esculentum (unc)
Pteris pendula
P. tremula (unc)
Pyrosia eleagnifolia
Sticherus cunninghamii
Trichomanes teniforme
- ORCHIDS**
- Corybas trilobus*
Drymoanthus adversus (unc)
Eatinia mucronata s.s. (unc)
Pterostylis venosa (unc)
Thelymitra sp. (unc)

GRASSES

Chionochloa cheesemanii
C.conspicua ssp. *conspicua* (unc)
Lachnagrostis lyallii (incl. *L. filiformis* var. *samiglabra*)

Colospernum hastatum (unc)
Dianella nigra
Libertia pulchella
L. sp. (cf. *L.grandiflora*)
Luzuriaga parviflora
Phormium cookianum

Microlaena avenacea

SEDGES

Gahnia pauciflora
G. procera
Uncinia astonii
U.filiformis
U.gracilenta
U.laxiflora
U.rupestris
U.silvestris (syn. *U.affinis*)
U.uncinata
U.zotovii

COMPOSITE HERBS

Anaphalis sp. (*Helichrysum bellidoides* s.s.)
Gnaphalium audax (unc)
G.gymnocephalum (unc)
Lagenifera pumila (unc)
Pseudognaphalium sp. (unnamed; aff. *P. luteoalbum*)
Raoulia glabra (unc)

DICOT HERBS OTHER THAN**COMPOSITES**

Acaena australis
Cardamine sp. (*C. debilis* agg.) (unc)
Epilobium nummulariifolium (unc)
E.rotundifolium (unc)
Hydrocotyle moschata (unc)
Myosotis forsteti (unc)

Nicotera depressa (incl. *N.cunninghamii*)
N.sp. (unnamed; aff. *N.dichondrifolia*)
Oreisia lactea ssp. *lactea*

Oxalis magellanica
Ranunculus membranifolius
R.reflexus (unc)

MONOCOT HERBS OTHER THAN**ORCHIDS, GRASSES, SEDGES, RUSHES**

Astelia fragrans
A. trinervia
A.sp.(i) (unnamed; included in
A.nivicola, as var. *moticeae*,
by Moore 1966)

A.sp.(ii) (unnamed; aff. *A.nervosa*)
(lvs broad, not conspicuously
white adaxially as in *A.ner-
vosa*)

Stellaria decipiens (incl. *S.parvi-
flora* (unc))
Urtica incisa (unc)
Viola filicaulis

ADVENTIVE PLANTS
(all uncommon)

GRASSES

Agrostis capillaris
Poa annua

COMPOSITE HERBS

Cirsium vulgare
Hypochaeris radicata
Mycelis muralis

DICOT HERBS OTHER THAN COMPOSITES

Cerastium fontanum ssp. *vulgaris*
Digitalis purpurea
Linum catharticum
Sagina procumbens

No. of indigenous spp.: 220.

No. of adventive spp.: 9.