


INDIGENOUS HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS
FLOWERING PLANTS) OF WAINGARO PK* AND W LOCKETT RA
NW NELSON, 1100-1600M

A.P.Druce, Wellington, May 1990
(revised May 1991, Aug. 1992, Aug. 1993)
(Based on visits 1970-1990)

*Grid ref. NZMS 260 M26 711182

Rocks: argillite, sandstone, quartzite, phyllite, slate
(no carbonate rocks)



(unc)-uncommon

GYMNOSPERM TREES AND SHRUBS

- | | |
|--|---|
| <i>Halocarpus biformis</i> | <i>C. depressa</i> |
| <i>Lepidothamnus laxifolius</i> | <i>C. perpusilla</i> ssp. <i>perpusilla</i> |
| <i>L. intermedius</i> X <i>L. laxifolius</i> | <i>C. pseudocuneata</i> ssp. |
| <i>Phyllocladus alpinus</i> agg. | <i>C. sp.(i)</i> (unnamed; aff. <i>C. pseudocuneata</i>) (low-growing shrubs with small, thick, arcuate lvs) |
| <i>Podocarpus nivalis</i> | <i>C. sp.(ii)</i> (unnamed) (<i>C. parviflora</i> var. <i>dumosa</i> sensu Allan 1961, |

DICOT TREES

- | | |
|--|--|
| <i>Dracophyllum traversii</i> (incl. <i>D. pyramidale</i>) | <i>D. filifolium</i> and <i>D. uniflorum</i> |
| <i>Griselinia littoralis</i> | <i>C. pumila</i> |
| <i>Nothofagus menziesii</i> | <i>C. campetrifolia</i> |
| <i>N. solandri</i> var. <i>cliffortioides</i> | <i>Dracophyllum longifolium</i> var. (<i>D. longifolium</i> and <i>D. oliveri</i>) |
| <i>Pseudopanax linearis</i> | <i>P. simplex</i> (incl. <i>P. s. var. sinclairii</i>) |
| <i>P. sp. (unnamed; included in <i>P. colensoi</i>, as var. <i>fiordensis</i> and as var. <i>ternatus</i>, by Wardle 1968)</i> | <i>D. pubescens</i> var. <i>pubescens</i> |
| | <i>D. uniflorum</i> |
| | <i>Exocarpus bidwillii</i> |
| | <i>Gaultheria crassa</i> |

DICOT SHRUBS

- | | |
|--|--|
| <i>Archeria traversii</i> (incl. <i>A. t. var. australis</i>) | <i>G. depressa</i> , as var. <i>novaeglandiae</i> , by Franklin 1962 |
| <i>Aristotelia fruticosa</i> var. | <i>G. crassa</i> X <i>G. sp. (unnamed)</i> |
| <i>Brachyglottis adamsii</i> | <i>Hebe albicans</i> var. <i>albicans</i> |
| <i>B. bidwillii</i> var. <i>viridis</i> | <i>H. canterburiana</i> |
| <i>B. rotundifolia</i> (incl. <i>B. buchananii</i> and <i>B. cockaynei</i>) | <i>H. ciliolata</i> |
| | <i>H. cockayneana</i> var. |
| | <i>H. mactantha</i> var. <i>brachiphylla</i> |
| <i>Cassinia laevis</i> | <i>H. tetragona</i> var. (<i>H. coarctata</i>) |
| <i>Coprosma cheesemanii</i> ss. (unc) | <i>H. topiaria</i> |
| <i>C. ciliata</i> ssp. <i>ciliata</i> | <i>H. venusta</i> |
| <i>C. colensoi</i> s.s. | <i>H. sp. (<i>Leonohebe masoniae</i>)</i> |
| <i>C. crenulata</i> | |

ORCHIDS

Leptospermum scoparium
Melicytus sp. (unnamed)

Caladenia hyallii
Chiloglottis cornuta
Corybas trilobus s.s.
Microtis oligantha (unc)

Myrsine divaricata s.s.
M. nummularia
Olearia colensoi var. *colensoi*

GRASSES

Pentachondra pumila
Pimelea oreophila
Pittosporum rigidum var. (*P.*
crassicaule)

Agrostis sp. (unnamed; aff. *A.*
dyeri) (panicle not contracted
 as in *A. dyeri*)
Chiachiochloa conspicua ssp.
conspicua
C. pallens ssp. *pallens*

DICOT LIANES AND RELATED**TRAILING PLANTS**

Muehlenbeckia axillaris
Rubus parvus (unc)

C. tubria ssp. *occulta*
C. sp. (unnamed) (*C. flavescentia* avct.
 South I, non *C. f.* Zotov) (unc)

LYCOPODS

*Lycopodium australe*num
L. fastigiatum
L. scariosum
L. variatum (incl. *L. billardierei*
 and *L. novae-zelandicum*)
 (unc)

Deschampsia tenella
Deyeuxia aucklandica
Hierochloe novae-zelandiae
Microlepia colensoi s.s.
Poa colensoi

FERNS

Blachium penna-martini
B. procerum
Grammitis patagonica (unc)
G. poeppigiana

P. dipsacea (unc)
P. sp. (*P. subvestita* or *P. nz.* s.s.)
Rytidosperma nigricans

Triscomia youngii

SEDGES

Hymenophyllum multifidum
Hypolepis millefolium
Polystichum cystostegia
P. vestitum

Carex acicularis
C. echinata var. (*C. stellulata* var.
australis)
C. gaudichaudiana

- C.sinclairii*
C.sp.(unnamed; aff.C.testacea)
 (Ivs broader than in *C.testacea*,
 ls. reddish; uticle margins
 not scabrid)(unc)
- Carpha alpina* s.s.
Gahnia procera
Isolepis habra
Oreobolus pectinatus
O.strictus
Schoenus pauciflorus s.s.
Uncinia divaricata
U.drucei (unc)
U.filiformis
U.fuscovaginata var. (plants
 glaucous)
U.sp.(i) (undetermined)
U.sp.(ii) (undetermined)
- A.nivicola* ss.
A.petriei
A.skottsbergii
A.sp.(i) (unnamed; included in *A.*
nivicola, as var. *moricacea*, by
 Moore 1966)
- A.sp.(ii)* (unnamed; aff. *A.nervosa*)
 (Ivs broad, not conspicuously
 white adaxially as in *A.nervosa*)
Bulbinella hookeri
Libertia pulchella
Luzuriaga parviflora
Phormium cookianum

COMPOSITE HERBS

- Abrotanella linearis* s.s.
Anaphalis sp. (*Helichrysum*
bellidioides s.s.)
Brachyglottis sp. (unnamed; in-
 cluded in *B.bellidioides*, as
 var. *crassa*, by Nordenstam 1978)
Brachyscome radicata

- RUSHES AND ALLIED PLANTS**
- Juncus antarcticus*
J.novae-zelandiae
J.pusillus
Luzula migrata
*L.sp. (unnamed; included
 in *L.crinata*, as var. *petrie-
 ana*, by Edgar 1966)*
Matsippaspermum gracile

- B.sinclairii* s.s.
Celmisia alpina s.s.

- C.bellidioides*
C.dallii
C.gibbsii
C.hiaracifolia ssp.

- MONOCOT HERBS OTHER THAN
 ORCHIDS, GRASSES, SEDGES, RUSHES**
- Astelia graminea*
A.linearis var. *linearis*

<i>C. incana</i> (incl. <i>C. discolor</i>)	<i>L. sp.</i> (unnamed; included in <i>L. squalida</i> , as <i>ssp. mediana</i> , by Lloyd and Webb, 1987)
<i>C. laticifolia</i>	
<i>C. lateralis</i> (incl. <i>C.l.</i> var. <i>villosa</i>)	
<i>C. monroi</i> ssp. (<i>C. coriacea</i> var. <i>lancifolia</i>)	<i>L. sp (iii)</i> (unnamed) (pinnae entire or almost so)
<i>C. petrioi</i> (unc)	
<i>C. rupestris</i>	<i>Leucogenes grandiceps</i>
<i>C. similis</i>	<i>Microseris scapigera</i> var.
<i>C. sinclairii</i> (incl. <i>C. albuni</i> + <i>C. dutielzii</i>)	<i>Raoulia eximia</i> var.
<i>C. spectabilis</i> ssp.	<i>R. glabra</i>
<i>C. traversii</i>	<i>R. grandiflora</i> var. <i>grandiflora</i>
<i>C. similis</i> × <i>C. spectabilis</i> (unc)	<i>Taraxacum magellanicum</i>
<i>Craspedia</i> sp(i) (plants summer-green; lvs green, narrowly elliptic-spathulate, both surfaces densely clad in short glandular hairs; fls white)	
<i>C. sp(ii)</i> (plants summer-green; lvs green, narrowly obovate spathulate, both surfaces densely clad in short non-glandular hairs; fls white)	
<i>C. sp.(iii)</i> (unnamed; included in <i>C. lanata</i> , as var. <i>elongata</i> , by Allan 1961)	
<i>Dolichoglottis scorzonerooides</i>	<i>Acaena ansatinifolia</i>
<i>Gnaphalium latifoliale</i>	<i>A. profunda</i> incisa
<i>G. mackayi</i> s.s	<i>Aciphylla ferox</i>
<i>G. traversii</i>	<i>A. polita</i> var.
<i>Haastia sinclairii</i> ss.	<i>Anisotome aromatica</i>
<i>Laganifera strangulata</i>	<i>A. deltoidea</i>
<i>Leptinella pyrathrifolia</i> agg.	<i>A. filifolia</i>

DICOT HERBS OTHER THAN
COMPOSITES

<i>Cardamine</i> sp. (i) (<i>C. dabilis</i> agg.)
<i>C. sp.(ii)</i> ("Scree Race" of Pitchard 1957)
<i>C. sp.(iii)</i> (unnamed) (lvs mostly simple, occ. with up to 2 pairs of pinnules; blades ± reniform)
<i>Colobanthus affinis</i>
<i>C. canaliculatus</i>
<i>Drosera arcturi</i>
<i>D. spathulata</i>
<i>D. stanopetala</i>

- Epilobium brunnescens* s.s.
E. chlorifolium
E. glabellum s.s.
E. pernitens
E. rubromarginatum
E. tasmanicum
E. sp.(i) (unnamed; included in *E. brunnescens*, as ssp. *minutiflorum*, by Ravene & Engelhorn, 1971)
E. sp.(ii) (unnamed; aff. *E. glabellum*)
Euphrasia townsonii ss.
E. sp.(i) (unnamed; aff. *E. petriei*)
E. sp.(ii) (unnamed; aff. *E. laingii*)
E. sp.(iii) (cf. *E. revoluta*)
Forsiera mackayi
F. sedifolia s.s.
F. tanella
Gentiana bellidifolia
G. gracilifolia (incl. *G. vernicosa*)
G. montana (incl. *G. m. var. stolonifera*, *G. patula*, and *G. bellidifolia* var. *pulchella*)
G. spenceti
G. sp. (unnamed; aff. *G. bellidi-folia*) (plants forming small to large patches; lvs very glossy; fls large, creamy white)
Geranium sp. (unnamed; aff. *G. microphyllum*)
Geum laiospermum
G. uniflorum
G. sp.(i) (*G. parviflorum* agg.) (stock short; terminal leaflet about as long as broad; fls up to 13mm diam.)
G. sp.(ii) (*G. parviflorum* agg.) (stock creeping and branching; terminal flt slightly longer than broad; fls up to 8mm diam.) (unc)
- Gonocarpus aggregatus*
Hydrocotyle sulcata
H. sp. (unnamed; included in *H. novae-zealandiae*, as var. *montana*, by Kirk 1899)
Kelletia dieffenbachii
K. laxa
K. multiflora
K. tessellata
Montia fontana ss.
Myosotis macrantha
Neopaxia sp. (incl. *Montia calycina* and *Claytonia austro-asiatica* var. *biflora*)
Nertera ciliata
Notothlaspi sp. (unnamed) (*N. australe* auct., non *thlaspi australe* Hook.f.)
Oreomythus colensoi ss. (incl. *O. c. var. hispida* and *O. c. var. multifida*)
Oreostylidium subulatum
Ourisia caespitosa ssp. *caespitosa*
O. lactea ssp. *lactea*
O. simpsonii
Oxalis magellanica
Parahabea cheesemanii
P. decora
P. lyallii
Plantago lanigera s.l. (incl. *P. n.z.*)
P. raoulii
Pratia angulata agg.
P. macrodon s.s.
Psychrophila novae-zealandiae

7

Ranunculus foliosus No. of species: 231.
R. insignis
R. verticillatus
Schizolema haastii s.s.
S.sp. (unnamed; included in *S.*
 haastii, as ssp. *cyanopetalum*, by
 Domin (1908))
Stellaria decipiens (incl. *S. parviflora*)
S. gracilenta
Viola cunninghamii
Wahlenbergia albomarginata
var. *albomarginata* (incl. *W.*
 pumila var. *laxa* ^{and} *W. brockiei*)

References

- Allan, H.H. 1961: "Flora of N.Z." Vol. 1.
Domin, K. 1908: Engl. Bot Jb 40: 573-85.
Edgar, E. 1970: In "Flora of N.Z." Vol. 2.
Franklin, D.A. 1962: Trans R. Soc NZ Bot. 1: 155-73.
Kirk, T. 1899: "The Students' Flora of N.Z."
Lloyd, D.G.; Webb, C.J. 1987: N.Z. Jl Bot. 25: 99-105.
Nordenstam, B. 1978: Opera Bot. 44: 1-84.
Pritchard, G.G. 1957: Trans R. Soc. NZ 85: 75-89.
Raven, P.H.; Engelhorn, T. 1971: N.Z. Jl Bot. 9: 345-50.
Wardle, P. 1968: N.Z. Jl Bot. 6: 226-36.
Moore, L.B. 1966: N.Z. Jl Bot. 4: 201-40.
- — —

