

INDIGENOUS VASCULAR PLANTS IN THE VICINITY OF LAKES  
WAIKAREMOANA AND WAIKAREITI, ON THE PANEKIRI AND  
NGAMOKO RANGES, AND ON MANUOHA, UREWERA NATIONAL  
PARK, 1000-4600 FT (300-1400M)


(Based on observations 1949-73; additions from W.B. Shaw)

A.P. Druce, Botany Division, D.S.I.R., 1972

N105 580 295

(revised May 1974,  
Oct 1974)

June 1975, Apr 1976



- 440000 N
- July 1976
- May 1977
- March 1978
- June 1979
- June 1980
- 43
- April 1982
- Oct. 1983
- June 1985
- July 1985
- Sept 1985
- March 1987
- 42
- June 1987
- Nov. 1987
- June 1988

(Additions by W.B. Shaw and B.D. Clarkson, 1984-5)

CYMNOSPERM TREES AND SHRUBS

- Dacrycarpus dactyloides*
- Dacrydium cupressinum*
- Halocarpus bidwillii*
- H. bifornis*
- Lagarostrobos colensoi*
- Lepidothamnus laxifolius*
- Libocedrus bidwillii*
- Phyllocladus alpinus* ss.
- P. glaucus*
- P. trichomanoides* ss.

- P. alpinus* x *P. glaucus* (unc)
- Pedocarpus hallii*
- P. tetara* s.s.
- Prommopitys ferruginea*
- P. taxifolia*
- Nov. 1990
- July 1991
- July 1992
- May 1993

MONOCOT TREES AND SHRUBS

- Cordyline australis*
- C. banksii*
- C. indivisa*
- C. australis* x *C. banksii*
- Rhopalostylis sapida* (unc)

(unc) - uncommon or local

## DICOT TREES

*Alectryon excelsus* var. *excelsus*  
*Aristolelia serrata*  
*Beilschmiedia tawa*  
*Carpodetus serratus*  
*Dracophyllum latifolium* (incl. *D. matthewsii*)  
*Elaeocarpus dentatus* (incl. *E. d. var. obovatus*)  
*E. hookerianus*  
*Fuchsia excorticata*  
*Griselinia littoralis*  
*G. lucida*  
*Hedycarya arborea*  
*Hoheria populnea* var. *lanceolata*  
*Ixerba brexioides*  
*Knightia excelsa*  
*Macropiper excelsum* var. *excelsum*  
*Melicactus lanceolatus* (incl. *M. l. var. latior*)  
*M. ramiflorus* ssp. *ramiflorus*  
*Metrosideros excelsa* (Colenso, 1894, and Cheeseman, 1907, as *M. tomentosa*)  
*M. robusta*  
*Myrsine australis*  
*M. salicina*  
*Nestegis cunninghamii*  
*N. lanceolata*  
*Nothofagus fusca*  
*N. menziesii*  
*N. solandri* var. *cliffortioides*  
*N. s. var. solandri*  
*N. truncata*  
*N. fusca* x *N. solandri*  
*N. solandri* x *N. truncata*  
*Olearia paniculata*  
*O. rani* var. *colorata*

*Pannantia eorymbosa*  
*Pittosporum eugenoides*  
*P. talphii*  
*P. tenuifolium* ssp. *tenuifolium*  
  
*Pseudopanax arboreus* var. *arbor.*  
*P. colensoi* s.s.  
*P. crassifolius*  
*P. edgerleyi*  
*P. simplex* (incl. *P. s. var. sinclairii*)  
*Quintinia serrata* (incl. *Q. acutifolia* and *Q. elliptica*)  
*Schefflera digitata*  
*Sophora tetraptera*  
  
*Weinmannia racemosa* var. *racemosa*

## DICOT SHRUBS

*Alseuosmia pusilla*  
*Aristolelia fruticosa* var. *nictophylla*  
*Brachyglottis kirkii*  
*B. repanda* s.s.  
*Carmichaelia arborea* var. (*C. odorata*)  
*C. sp.* (subgenus *Carmichaelia*)  
*Cassinia leptophylla* var. *leptophylla*  
*Clianthus puniceus* (unc)  
*Coprosma colensoi* s.s.  
  
*C. depressa* (Manuoha)  
*C. foetidissima*  
*C. grandifolia*  
*C. lucida* s.s.  
*C. microcarpa*

*C. propinqua* ssp. *propinqua*  
*C. pseudocuneata* (Manuoha) ssp. *pseudocuneata*  
*C. hamnoides* (incl. *C. polymorpha*)  
*C. rigida*  
*C. robusta*  
*C. rotundifolia*  
*C. rugosa* ssp. *rugosa*  
*C. tenuifolia*  
*C. sp. (i)* (unnamed; included in *C. parviflora* by Oliver, 1935, and others) (If undersurface not minutely hairy as is usual in *C. parviflora*; drupes vs. unpigmented, occ. purplish red)  
*C. sp. (ii)* (unnamed) (*C. parviflora* var. *dumosa* Allan 1961, non *C. p.* var. *dumosa* Cheeseman)  
*C. propinqua* x *C. robusta*  
*Cotania arborea*  
*C. kingiana* (Manuoha)  
*Cotokia cotoneaster* var. *cotoneaster*  
*Cyathodes empetrifolia*  
*C. sp.* (*C. juniperina* ass.)  
*Dracophyllum longifolium* var.

*Gaultheria antipoda*  
*G. sp.* (unnamed; included in *G. depressa*, as var. *novae-zelandiae*, by Franklin 1962)  
*Geniostoma rupestre* var. *ligustri-folium*  
*Hebe macrocarpa* var. (*H. corrigeni*)  
*H. odota* s.s.  
*H. stricta* var. *stricta*  
*H. sp. (i)* (unnamed; included in *H. stricta*, as var. *egmontiana*, by Moore 1961)

*H. sp. (ii)* (*Veronica arborea*)  
*H. macrocarpa* x *H. sp. (a)*  
*Helichrysum aggregatum* var. *aggregatum*  
*Illeostylus micranthus* (unc)  
*Kunzea ericoides* agg.  
*Leptospermum scoparium*  
*Leucopogon fasciculatus* var.  
*L. fraseri* ss.  
*Melicope simplex*  
*Myrsine divaricata* ss.  
*Neomyrtus pedunculata*  
*Olea arborescens*  
*O. capillaris* (Manuoha)  
*O. cheesemani* (Cheeseman 1925, as *O. arborescens* var. *angustifolia*; Poole and Adams, 1963)  
*O. colensoi* var. *colensoi* (Manuoha)  
*O. furfuracea* var.  
*O. ilicifolia* (unc)  
*O. arborescens* x *O. capillaris*  
*O. a.* x *O. ilicifolia*  
*Pentachondra pumila*  
*Peraxilla colensoi*  
*P. tetrapetala*  
*Pimelea longifolia* var. *longifolia*  
*P. prostrata*  
*Pittosporum rigidum* var. *rigidum*

*Pseudopanax anomalus*  
*P. a.* x *P. simplex* var. *sinclairii*  
*Pseudowintera axillaris*  
*P. colorata*  
*Solanum aviculare* var. *aviculare*

*Tupeia antarctica* (unc)  
*Urtica ferox* (unc)

MONOCOT LIANES

*Freyinetia baueriana* ssp. *banksii*  
*Ripogonum scandens*

DICOT LIANES

*Clematis cunninghamii* (Trevarthen & Mason, 1950, as *C. parviflora*)

*C. forsteri*  
*C. paniculata*  
*Metrosideros colensoi* (incl. *M. c. var. pendens*)  
*M. diffusa*  
*M. perforata*  
*M. fulgens* (Trevarthen and Mason, 1950, as *M. scandens*)  
*Muehlenbeckia australis*  
*M. complexa*  
*M. australis* X *M. complexa*  
*Parsonia capsularis*  
*P. heterophylla*  
*Passiflora tetrandra*  
*Rubus cissoides* var. *cissoides*  
*R. schmidlioides* var. *schmidlioides*  
*R. cissoides* X *R. schmidlioides* var. *schmidlioides*

*Azolla filiculoides* (unc)  
*Blechnum chambersii*  
*B. colensoi*  
*B. discolor*  
*B. filiforme*  
*B. fluviatile* ass.  
*B. membranaceum*  
*B. minus*  
*B. penna-marina*  
*B. procerum*  
*B. vulcanicum*  
*B. sp. (i)* (unnamed) (common lowland sp.; lower pinnae reduced in length)  
*B. sp. (ii)* (unnamed) (cliff sp.; lower pinnae not reduced)  
*Botrychium biforme*

*Ctenopteris heterophylla*

*Cyathia colensoi*

*C. dealbata*

*C. medullaris*

*C. smithii*

*Dicksonia fibrosa*

*D. lanata* s.s.

*D. squarrosa*

*Gleichenia dicarpa*

*Grammitis billardierei*

*G. ciliata* (type loc.)

*G. magellanica* ssp. *nolthofaeti*

*G. patagonica*

*Histiopteris incisa*

*Hymenophyllum atrovirens*

*H. bivalve*

*H. demissum*

*H. dilatatum*

*H. ferrugineum*

*H. flabellatum*

*H. flexuosum* (unc)

*H. lyallii*

*H. malingii*

#### PSILOPSIDS AND LYCOPODS

*Tmesipteris elongata* (incl. *T. e. ssp. robusta*)

*T. tannensis*

*Lycopodium australianum*

*L. fastigiatum*

*L. laterale* (incl. *L. ramulosum*?)

*L. scarosum*

*L. varium* (incl. *L. billardierei* and

*L. novae-zelandicum*)

*L. volubile*

#### FERNS

*Adiantum cunninghamii*

*Anarthropteris lanceolata*

*Asplenium bulbiferum*

*A. flaccidum*

*A. gracillimum*

*A. hookerianum*

*A. lyallii*

*A. oblongifolium*

*A. polyodon*

*A. sp.* (*A. trichomanes* ass.) (unc)

H. multifidum  
H. peltatum  
H. pulcherrimum  
H. rarum  
H. revolutum  
H. rufescens  
H. sanguinolentum ass.  
H. scabrum  
H. villosum (type loc.)  
Hypolepis ambigua  
H. millefolium  
H. rufobarbata  
Lastreopsis glabella  
L. hispida

Ophioglossum coriaceum ass.  
Paesia scaberula  
Pellaea rotundifolia  
Pilularia novae-zelandiae (W.R.B.  
Oliver pers. comm.)  
Phymatosorus novae-zelandiae  
P. pustulatus  
P. scandens  
Pneumatopteris pennigera  
Polystichum richardii  
P. silvaticum  
P. vestitum  
Pteridium esculentum  
Pteris pendula  
Pyrrhosia elcagnifolia  
Rumohra adiantiformis  
Schizaea australis

Leptolepia novae-zelandiae  
Leptopteris hymenophylloides  
L. superba  
L. hymenophylloides X L. superba  
Trichomanes colensoi  
T. reniforme  
T. venosum

ORCHIDS

Acianthus sinclairii  
Adenochilus gracilis  
Aporostylis bifolia  
Caladenia carnea agg.  
Chiloglottis cornuta  
Corybus acuminatus  
C. macranthus  
C. pilularis ss.  
C. trilobus ss.  
Dendrobium cunninghamii  
Earina autumnalis  
E. mucronata ss.  
Genoplesium nudum  
Microtis unifolia  
Orthoceras novae-zeelandiae (unc)  
Prasophyllum colensoi  
Pterostylis banksii  
P. montana (Trevorthen and  
Mason, 1950)  
P. patens  
Thelymitra cyanea  
T. hatchii  
T. longifolia  
T. pauciflora agg.

**GRASSES**

- Echinopogon ovatus
- Elymus multiflorus
- E. narduroides
- E. sp. (aff. E. rectisetus)
- Agrostis personata
- Chionochloa cheesemanii
- C. conspicua ssp. cunninghamii
- C. flavicans
- C. rubra ssp. rubra
- C. cheesemanii X C. conspicua ssp. cunninghamii
- C. conspicua ssp. cunninghamii X C. rubra
- Cortaderia fulvida
- C. toetoe
- Deschampsia caespitosa var. macrantha
- D. novae-zelandiae
- Dayeuxia avenoides (incl. D. a. var. brachyantha)
- Dichelachne ciliata
- Festuca multinodis
- Hierochloa redolens
- Lachnagrostis filiformis s.s.
- L. lyallii (incl. L. filiformis var. semiglabra)
- L. striata
- L. sp. (ii) (unnamed) (tall, slender; lvs narrow; awns short)
- Microlaena avenacea
- M. stipoides

- Rytidosperma buchananii
- R. clavatum
- R. gracile
- R. viride (Manuoha)

- Poa anceps ssp. anceps
- P. breviglumis
- P. imbecilla
- Trisetum sp. (unnamed; aff. T. antarcticum)
- Zoysia minima (Zotov 1971)

**SEDGES**

- Baumea articulata (unc)
- B. huttonii
- B. rubiginosa
- B. tenax
- Carex breviculmis
- C. elithosa (unc)
- C. dipsacea var. dipsaccea
- C. dissita s.s.
- C. echinata var. (C. stallulata var. australis)
- C. flagellifera
- C. forsteri
- C. gaudichaudiana
- C. inversa
- C. maorica
- C. secta s.s.
- C. sinclairii
- C. solandri
- C. virgata
- C. sp. (Section Acutae)
- Catpua alpina s.s.
- Cyperus ustulatus
- Eleocharis acuta
- E. gracilis

- E. pusilla
- E. sphacelata
- Gahnia pauciflora
- G. procera
- G. setifolia
- Isolepis habitata
- I. inundata
- I. pottsii
- I. reticulatis
- I. subtilissima
- Lepidosperma australe
- Machaetina sindaitii
- Morelotia affinis (unc)
- Oreobolus pectinatus
- Schoenoplectus validus (unc)
- Schoenus maschalinus
- Uncinia caespitosa (Manuoha)
- U. clavata
- U. distans
- U. ferruginea
- U. filiformis
- U. gracilentata
- U. longifructus
- U. rupestris (incl. U. angustifolia)
- U. scabra
- U. silvestris (incl. U. affinis)
- U. uncinata
- U. zotovii
- U. sp. (i) (unnamed; aff. U. nervosa) (more slender than U. nervosa; see Edgar 1970, p. 231) (Manuoha)

- Luzula migrata
- L. picta s.s.
- L. subclavata
- L. sp. (unnamed; included in L. picta, as var. limosa, by Edgar, 1966)

**MONOCOT HERBS (OTHER THAN ORCHIDS, GRASSES, SEDGES, RUSHES)**

- Arthropodium candidum
- Astelia fragrans
- A. solandri
- A. sp. (unnamed; aff. A. nervosa) (lvs not conspicuously white adaxially as in A. nervosa)
- Collospermum hastatum
- C. microspermum
- Dianella nigra
- Lemna sp. (L. minor auct. N.Z.)
- Libertia grandiflora
- L. pulchella
- Luzuriaga parviflora
- Phormium cookianum
- Potamogeton cheesemanii
- P. suboblongus

> (see foot of next column)

**RUSHES**

- Centrolepis ciliata
- Eleocharis minus
- Juncus gregiflorus
- J. pallidus
- J. planifolius
- J. novae-zelandiae

- Triglochin striata
- Typha orientalis

> U. sp. (ii) (unnamed) (U. affinis sensu Hamlin 1959, non U. riparia var. affinis Col. ex Clarke; see Nooteboom 1978) (Manuoha)

COMPOSITE HERBS

- Brachyscome radicata (unc)
- Calmisia gracilentia ssp. (Manuoha)
- C. spectabilis ssp. spectabilis
- C. sp. (unnamed) (plants strongly rhizomatous)
- C. spectabilis X Olea arborescens (unc)
- Centipeda minima

- Craspedia minor var. minor (incl. C. major)
- C. viscosa

- Anaphalis ketiensis s.s.
- A. sp. (i) (Helichtysum bellidioides s.s.)
- A. sp. (ii) (unnamed) (Helichtysum alpinum of Cockayne 1928)

- Gnaphalium audax
- G. delicatum
- G. gymnocephalum
- G. involucreatum
- G. huahinicum
- G. sphaericum
- G. traversii
- G. sp. (unnamed; aff. G. paludosum) (lvs linear, green on upper surface) (Manuoha)

- Helichtysum filicaule
- Lagenifera pumila
- L. strangulata
- Leptiniella squalida s.s.
- Pseudognaphalium sp. (unnamed; aff. P. luteoalbum) (lvs mostly linear-spathulate to linear, acute)
- Raoulia albosericea
- R. glabra
- R. tenuicaulis (incl. R. t. var. dimorpha and R. t. var. pusilla)

- Senecio banksii var.
- S. glaucophyllus ssp. tota
- S. glomeratus
- S. minimus

- S. rufiglandulosus var. rufiglandulosus
- Vitadinia australis agg.

DICOT HERBS (OTHER THAN COMPOSITES)

- Acaena ansetinifolia
- Aciphylla sp. (unnamed; aff. A. squarrosa)
- Cardamine sp. (i) (C. debilis agg) ("Long Style" of Pritchard, 1957)
- C. sp. (ii) (C. debilis agg) ("Glossy Leaf" of Pritchard, 1957)
- Callitriche petriei ssp. petriei
- Centella uniflora
- Crassula pedunculatis (unc)
- Dichondra repens var. (unc)
- Drosera binata
- D. peltata ssp. auriculata
- D. spathulata
- Elatine gratioloidea (unc)
- Epilobium alsinoides s.s.
- E. atriplicifolium s.s.
- E. brunnescens s.s.
- E. cinereum
- E. cockayneanum
- E. insulare
- E. Komarovianum
- E. linnaeoides
- E. microphyllum
- E. netteroides
- E. nummulariifolium
- E. pubens
- E. rotundifolium
- E. tenuipes s.s.
- E. sp. (unnamed; included in E. brunnescens, as ssp. minutiflorum, by Raven and Engelhorn, 1971)
- Euphrasia cuneata
- Galium propinquum
- G. trilobum
- Gentiana grisebachii (Manuoha)



G. sp. (unnamed; aff. G. spenceri) (discussed by Allan 1961, p. 769)

Geranium sp. (aff. G. microphyllum)

G. potentilloides var. potentilloides

G. sessiliflorum var. novae-zealandiae

Glossostigma elatinioides

Gonocarpus aggregatus

G. micranthus ssp. micranthus

Gratiola sexdentata

Gunnera monoica

Haloragis erecta ssp. erecta

Hydrocotyle dissecta (Trevorthen and Mason, 1950)

H. elongata

H. heteromeria

H. hydrophila

H. microphylla

H. moschata

H. novae-zealandiae ssp.

H. pterocarpa

H. sulcata

Hypericum japonicum ass.

Hypsela rivalis

Jovellana repens

Lilaeopsis ruthiana

Limosella lineata

Linum monogynum

Mentha cunninghamii

Myosotis forsteri

M. saxosa (incl. M. amabilis) (Panekiti Ra.)

M. venosa

Myriophyllum pedunculatum

ssp. novae-zealandiae

M. propinquum

M. triphyllum

Nertera ciliata

N. depressa (incl. N. cunninghamii)

N. scapanioides

N. sp. (unnamed; aff. N. dichondrifolia) (hairs present on both surfaces of lf, straight, not hooked as in N. dichondrifolia)

Oreomyrrhis edenae s.s. (incl. O. c.

var. hispida and O. c. var. multifida)

O. sp. (unnamed; aff. O. ramosa)

Oursia lactea ssp. drucei

O. macrophylla ssp. robusta

Oxalis exilis

O. magellanica

Parahebe diffusa ssp. diffusa

Plantago raoulii

Potentilla anserinioides

Pratia angulata agg.

P. perpusilla

Ranunculus amphitrichus

R. reflexus

R. sp. (cf. R. royi)

Rorippa palustris

Rumex flexuosus

Scandia sp. (S. rosifolia ass.) (lfts ovate-lanceolate)

Schizeilema coleusoi

S. trifoliolatum

Stellaria decipiens (incl. S. parviflora)

Urtica incisa

U. lineatifolia (Allan 1961)

Viola cunninghamii

V. filicaulis

V. lyallii

Wahlenbergia albomarginata

var. pygmaea

W. coleusoi

W. rupestris

Doubtful and erroneous records not incorporated in the main list:

Record

Probable identity of plant on which record based

*Aciphylla squarrosa* var. *flaccida*

(Druce, 1954) ----- *A. sp.* (unnamed)

*Agrostis* sp. (*A. dyeri*?) (Trevarthen  
and Mason, 1950)

*Carex ochrosaccus* (T. and M., 1950,  
as *C. dissita* var. *ochrosaccus*)

*C. subdola* (T. and M., 1950) ----- *C. sinclairii*

*C. ternaria* (T. and M., 1950) ----- *C. sp.* (Section *Acutae*)

"*Deyeuxia forsteri* var. *humilior*" (Cheeseman, 1929) *Lachnagrostis striata*

*Gahnia lacera* (T. and M., 1950)

*Gaultheria depressa* (T. and M., 1950) - *G. sp.* (unnamed)

{ *Coprosma areolata* (T. and M., 1950)

{ *C. parviflora* (T. and M., 1950) ----- *C. sp.* (i) (unnamed)

*Hebe salicifolia* (T. and M., 1950) - - - *H. stricta*

*Isolepis cernua* (T. and M., 1950, as *Scirpus cernuus*)

*Juncus pauciflorus* (T. and M., 1950)

*Libertia ixioides* (T. and M., 1950)

*Lindsaea trichomanoides* (T. and M.,  
1950, as *L. coneata*)

*Nertera dichondrifolia* (T. and M., 1950) - - - *N. sp.* (unnamed)

*Oplismenus imbecillus* (T. and M., 1950,  
as *O. undulatifolius*)

{ *Pittosporum crassifolium* (T. and M.,  
1950) -----

*P. ralphi*

*Phormium tenax* (T. and M., 1950)

*P. cookianum* X *P. tenax* (T. and M.,  
1950)

{ *Plagianthus regius* (T. and M., 1950,  
as *P. betulinus*)

- Pteris tremula* (T. and M., 1950)  
*Sparganium subglobosum* (T. and M., 1950, as *S. antipodum*)  
*Toroniatoru* (T. and M., 1950, as *Persoonia toru*)  
*Thelymitra hatchii* X *T. longifolia*  
 (T. and M., 1950, as *T. pachyphylla* X *T. longifolia*)  
*T. pachyphylla* (Hatch, 1952) - - - - - *T. hatchii*  
 [*T. longifolia* var. *insularis* (T. and M., 1950) NOMEN NUDUM]  
*Trisetum antarcticum* (T. and M., 1950) - *T. sp.* (unnamed)  
*Uncinia caespitosa* (T. and M., 1950) - - *U. zotovii*  
  
*Vitex lucens* (T. and M., 1950)

No. of species (excluding doubtful records): 494.

REFERENCES

- Allen, H.H. 1961: "Flora of New Zealand". Government Printer, Wellington. Vol. 1. 1085 pp.
- Cheeseman, T.F. 1907: Contributions to a fuller knowledge of the flora of New Zealand. Trans. N.Z. Inst. 39: 439-50.
- 1925: "Manual of the New Zealand Flora". Ed. 2. Edited by W.R.B. Oliver. Government Printer, Wellington. 1163 pp.
- Cockayne L. 1928: "The Vegetation of New Zealand." Ed. 2. Englemann, Leipzig. 456 pp.
- Coleenso, W. 1895: Notes and reminiscences of early crossings of the romantically-situated Lake Waikaremoana, County of Hawke's Bay, of its neighbouring country, and of its peculiar botany; performed in the years 1841 and 1843. Trans. N.Z. Inst. 27: 359-82.
- Druce, A.P. 1954: Plant distribution records (2). Bull. Wellington bot. Soc. 27: 4-9.
- Edgar, E. 1966: Luzula in New Zealand. N.Z. J. Bot. 4: 159-84.
- 1970: In "Flora of New Zealand". Government Printer, Wellington. Vol. 2. 354 pp.
- Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Bernettya). Trans. R. Soc. N.Z. Bot. 1: 155-73.
- Oliver, W.R.B. 1935: The genus Coprosma. Bull. Bishop Mus. Honolulu, 132 pp.
- Pool, A.L.; Adams, N.M. 1963. "Trees and Shrubs of New Zealand". Government Printer, Wellington. 250 pp.
- Pritchard, G.G. 1957: Experimental taxonomic studies on species of Cardamine Linn. in New Zealand. Trans. R. Soc. N.Z. 85: 75-89.
- Hatch, E.D. 1952. The New Zealand forms of Thelymitra J.R. & G. Forster and appendices. Trans. R. Soc. N.Z. 79: 386-402.

- Raven, P. H.; Engelman, T. 1971: new taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. J. Bot. 9: 345-50.
- Trevarthen, C.; Mason, G. 1950: The vegetation of the Lake Waikaremoana area. Tane 3: 24-9.
- Zotov, V. D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.
-

