

HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS, FLOWERING PLANTS) OF KAWEKA RANGE<sup>1</sup>, INCLUDING BLACK BIRCH RANGE<sup>2</sup>

3000 - 5657 FT (900 - 1725 M)


- <sup>1</sup> Highest point: Kaweka, 5657 ft, grid ref. N123 829650
- <sup>2</sup> Highest point: Ngahere Hill, 3650 ft, grid ref. N123 870642

A.P. Druce, Botany Division, D.S.I.R., 1972

(revised May 1974, Dec 1974, July 1975, March 1976, July 1976, May 1977, June 1979, May 1980, Oct 1981, April 1984, May 1984, June 1984, May 1985, July 1985, March 1986, Apr. 1986, June 1990, Aug. 1992)

(Based on visits 1957, 1965, 1975, 1980, 1984, 1985, 1986, 1989)

(Additions by S. Beadel 1989)



THIS LIST IS FOR LAND ABOVE 3000 FT (900 M) - FOR A COMPLETE LIST OF HIGHER PLANTS FROM KAWEKA RA. AND ADJACENT AREAS SEE LIST 333

P - seen in Te Pukeohikarua (dissected plateaux) area<sup>†</sup>  
K - seen in Kaweka (main range) area  
B - seen in Black Birch area  
(species not marked are assumed to be widespread)

(<sup>†</sup> for boundaries see title page)  
(unc) - uncommon or local

### INDIGENOUS SPECIES

#### GYMNOSPERM TREES AND SHRUBS

<i>Halocarpus bidwillii</i> (P)	<i>Carpodetus serratus</i>
<i>Lepidothamnus laxifolius</i> (K, P)	<i>Elaeocarpus hookerianus</i> (unc)
<i>L. intermedius</i> x <i>L. laxifolius</i> (unc, K)	<i>Fuchsia excorticata</i>
<i>Phyllocladus alpinus</i> s.s. (B, K, P)	<i>Griselinia littoralis</i>
	<i>Nothofagus fusca</i>
	<i>N. menziesii</i> (P)
<i>Podocarpus hallii</i>	<i>N. solandri</i> var. <i>cliffortioides</i> (B, K, P)
<i>P. nivalis</i> (K, P)	<i>N. fusca</i> x <i>N. solandri</i> (unc)
<i>P. hallii</i> x <i>P. nivalis</i> (unc, K)	<i>Pittosporum ralphii</i> (unc, B and K)
<i>Prumnopitys ferruginea</i> (unc)	<i>P. tenuifolium</i> ssp. <i>colensoi</i> (unc, K and P)

#### MONOCOT TREES

<i>Cordyline australis</i> (unc)	<i>Pseudopanax arboreus</i> var. <i>arboreus</i>
<i>C. indivisa</i> (unc)	<i>P. colensoi</i> s.s. (K, P)
	<i>P. crassifolius</i> (unc)
	<i>P. simplex</i> (incl. <i>P. s.</i> var. <i>sinclairii</i> )
	<i>Schefflera digitata</i> (unc)

#### DICOT TREES

*Amstotelia serrata*

**DICOT SHRUBS**

- Alepis flavida* (unc)  
*Aristolelia fruticosa* var. *microphylla* (unc, K and P)  
*Brachyglottis bidwillii* var. *bidwillii* (K, P)  
*B. repanda* s.s. (unc)  
*B. rotundifolia* var. (*B. elaeagni folia*) (unc, K)  
*B. bidwillii* X *B. rotundifolia* (unc)  
*Cassinia vauvilliersii* (B, K, P)  
*Coprosma acerosa* ssp. (unc, K and P)  
*C. cheesemani* s.s. (B, K)  
*C. colensoi* s.s.  
*C. depressa* (K, P)  
*C. foetidissima*  
*C. grandifolia* (unc)  
*C. lucida* s.s. (unc)  
*C. microcarpa*  
*C. perpusilla* ssp. *perpusilla* (B, P, K)  
*C. pseudocuneata* ssp. *pseudoc.* (K, P)  
*C. thamnoides* (incl. *C. polymorpha*)  
  
*C. robusta* (unc)  
*C. rugosa* ssp. *rugosa* (K)  
*C. tenuifolia*  
  
*C. sp. (i)* (unnamed; included in *C. parviflora* by Oliver, 1935, and others) (If undersurface not minutely hairy as is usual in *C. parviflora*; drupes vs. unpigmented, occ. purplish red)  
  
*C. sp. (ii)* (unnamed; aff. *C. cheesemani*) (lvs broader than in *C. cheesemani*) (B, K, P)  
*C. robusta* X *C. tenuifolia* (unc)  
  
*Coriaria arborea*  
*C. kingiana* (B, K, P)  
*C. plumosa* (K, P)  
*C. pteridoides* (K, P)  
*C. arborea* X *C. kingiana*  
*C. a.* X *C. pteridoides*  
*C. kingiana* X *C. pteridoides*  
*Cyathodes dealbata* (incl. *C. pumila*) (F)  
*C. empetrifolia* (B, K, P)  
*C. sp. (C. juniperina agg.)* (lvs vs. <1cm X <1mm)  
*Dracophyllum recurvum* (B, K, P)  
*D. subulatum* (B, K, P)  
*D. recurvum* X *D. subulatum*  
*Epaemis alpina* (B, K, P)  
*Gaultheria antipoda*  
*G. colensoi* (B, K, P)  
*G. sp. (unnamed; included in G. depressa, as var. novae-zelandiae, by Franklin 1962)* (B, K, P)  
*G. antipoda* X *G. colensoi*  
*G. a.* X *G. sp. (unnamed)*  
*G. colensoi* X *G. sp. (unnamed)*  
*Hebe colensoi* (incl. *H. c. var. hillii*) (unc, K and P)  
*H. macrocarpa* var. (*C. cortiganii*) (B, K, P)  
*H. odora* s.s. (B, K, P)  
*H. stricta* var. *stricta*

- H. tetragona var. tetragona (B, K, P)
- H. venustula (incl. H. brachysiphon) (B, K, P)
- H. sp. (i) (unnamed; included in H. stricta, as var. egmontiana and as var. lata, by Moore 1961) (K)
- H. sp. (ii) (Veronica squakida) var. (K)
- H. macrocarpa x H. venustula (?) (unc)
- P. rigidum var. rigidum

Helichrysum aggregatum var. aggregatum

**DICOT LIANES AND RELATED**

**TRAILING PLANTS**

- Kunzea ericoides agg.
- Leptospermum scoparium
- Leucopogon fasciculatus var.
- L. fraseri s.s.
- Melicytus alpinus s.s. (K, P)
- Myrsine divaricata s.s.
- M. nummularia (K, P)
- Neomyrtus pedunculata
- Olearia arborescens
- O. furfuracea var.

- Clematis forsteri
- Muehlenbeckia axillaris
- Rubus cissoides var. (leaflets narrower than in var. cissoides)

**LYCOPODS**

- O. ilicifolia (unc) (Cunningham 1974)
- O. nummulariifolia var. nummul. (B, K, P)

- Lycopodium australianum (K)
- L. fastigiatum
- L. scariosum
- L. varium (incl. L. billardierei and L. novae-zelandicum)
- L. volubile (unc)

O. arborescens x O. furfuracea (unc)

**FERNS**

- O. a. x Celmisia spectabilis (unc)
- Pentachondra pumila (K, P)
- Petaxilla tetrapetala (unc)
- Pimelea buxifolia (K, P)
- P. longifolia var. longifolia (unc, K)
- P. microphylla (unc, K)
- P. prostrata
- P. suteri var. (P. oreophila) (K, P)
- Pittosporum divaricatum (Cunningham 1974) (B, K)

- Asplenium bulbiferum (unc)
- A. flaccidum
- A. fitchardii (unc, P)
- A. terrestre ssp. terrestre
- Blechnum chambersii (unc)
- B. colensoi (unc)
- B. discolor
- B. fluviatile agg.
- B. penna-marina

*B. procerum*  
*B. vulcanicum*  
*B. sp. (i)* (unnamed) (common sp.;  
 lower pinnae reduced in length)  
*B. sp. (ii)* (unnamed) (mountain sp.;  
 lower pinnae not reduced) (K)  
*B. procerum* x *B. sp. (i)*  
*Cyathea colensoi* (unc, K)  
*Dicksonia fibrosa* (unc)  
*D. lanata* s.s. (unc)  
*Gleichenia dicarpa* (B)  
*Grammitis billardierei*  
*G. magellanica* ssp. *notofageti*  
*G. patagonica* (unc, K)  
*G. poeppigiana* (K)  
*Histiopteris incisa*  
*Hymenophyllum cypressiforme* (unc, K)  
*H. minimum* (unc, K)  
*H. multifidum*  
*H. pettatum* (unc)  
*H. rarum* (unc)  
*H. revolutum* (unc)  
*H. sanguinolentum* agg.  
*Hypolepis ambigua*  
*H. millefolium* (K)  
*H. rufo barbata* (unc)  
*Leptopteris hymenophylloides*  
*L. superba* (unc)  
*Ophioglossum coriaceum* agg.  
*Paesia scaberula*  
*Polystichum vestitum*  
*Pteridium esculentum*  
*Sticherus cunninghamii*

## ORCHIDS

*Adenochilus gracilis* (unc, K and P)  
*Aporostylis bifolia* (B, K, P)  
*Caladenia carnea* agg.  
*C. lyallii* (unc, K and P)  
*Chiloglottis cornuta*  
*Corybas cryptanthus* (unc, K)  
*C. maceranthus*  
*C. trilobus* s.s.  
*Gastrodia cunninghamii*  
*Microtis unifolia*  
*Orthoceras novae-zeelandiae*  
*Prasophyllum colensoi* (B, K, P)  
*Genoplesium nudum* (unc, B)  
*Pterostylis humilis* (unc, P)  
*P. patens*  
*P. venosa* (unc, B and K)  
*Thelymitra cyanea*  
*T. decora*  
*T. hatchii*  
*T. longifolia*

## GRASSES

*Agrostis dycii* s.s. (K, P)  
*A. muelleriana* (K)  
*A. muscosa* (K)  
*A. personata* (K)  
*Chionochoa conspicua* (Cunningham  
 1974)  
*C. pallens* ssp. *pallens* (K, P)  
*C. rubra* ssp. *rubra* (B, P)  
*C. pallens* x *C. rubra* (unc, B and P)  
*Cortaderia fulvida*

- Deschampsia novae-zelandiae* (K)
- D. tenella* (K, P)
- Deyeuxia aucklandica* (K, P)
- D. avenoides* (incl. *D. a.* var. *brachyantha*)
- Dichelachne cuneata*
- Echinopogon ovatus*
- Elymus hardyioides* (unc, P)
- E. sp.* (aff. *E. rectisetus*) (Series 2, Group Waiouru, of Löve & Connor, 1982)
- Festuca sp.* (unnamed; aff. *F. rubra*) (K, P)
- Hierochloa recurvata* (K, P)
- H. redolens*
- Koeleria cheesemani* (K, P)
- Lachnagrostis sp.* (i) (unnamed; included in *L. filiformis*, as var. *semiglabra*, by Zotov 1965) (B, K, P)
- L. sp.* (ii) (unnamed) (tall, slender plants, with narrow lvs; glumes up to 4mm long; awn short or absent) (B)
- Microlaena avenacea*
- Poa anceps* ssp. *anceps*
- P. breviglumis*
- P. cita* ssp.
- P. colensoi* (B, K, P)
- P. lindsayi* (unc, K)
- P. sp.* (unnamed; aff. *P. novae-zel.*) (K)
- Rytidosperma gracile*
- R. pulchrum* (K, P)
- R. setifolium* (B, K, P)
- R. viride* (K)
- Trisetum sp.* (i) (unnamed; aff. *T. antarcticum*) (If sheath densely hairy) (B, K)
- T. sp.* (ii) (unnamed) (plants slender; If sheath glabrous or nearly so) (K)

- T. sp.* (iii) (unnamed; aff. *T. antarcticum*) (plants robust; If sheath glabrous at in upper part; If blade margins scabrid)
- SEDGES**
- Carex acicularis* (K)
- C. breviculmis*
- C. coriacea*
- C. dipsacea* var. (*C. tahoata*) (B)
- C. dissita* ss.
- C. echinata* var. (*C. stellulata* var. *australis*) (B)
- C. forsteri* (unc)
- C. gaudichaudiana* (unc, K)
- C. patrici* (unc, K)
- C. pyrenaica* var. *cephalotes* (Cunningham 1974)
- C. solandri*
- Carpha alpina* ss. (B, K, P)
- Eleocharis acuta*
- E. gracilis* (B, P)
- Isolepis aucklandica* (K)
- I. caligenis* (unc, K)
- I. crassiuscula* (P)
- I. habra*
- I. pottsii*
- I. subtilissima* (B, P)
- Lepidosperma australe*
- Oreobolus pectinatus* (B, K, P)
- Schoenus pauciflorus* ss.
- S. sp.* (unnamed; aff. *S. pauciflorus*) ("robust-form" of Edgar 1970, p. 174)
- Uncinia caespitosa* (K, P)
- U. clavata* (K)
- U. divaricata* (K)
- U. ferruginea* (unc)
- U. fuscovaginata* var. *fuscovag.* (K, P)
- U. rubra* (B, K)

- U. rupestris* (incl. *U. angustifolia*)
- U. silvestris* (*Cunninghamii* 1974) (K)
- U. uncinata*
- U. viridis* (K)
- U. sp. (i)* (unnamed) (*U. affinis* sensu Hamlin 1959, non *U. riparia* var. *affinis* Col. ex Clarke; see Nootboom 1978) (B, P)

*U. sp. (ii)* (unnamed; aff. *U. nervosa*) (more slender than *U. nervosa*; see Edgar 1970, p. 231 (unc, B))

**RUSHES**

- Centrolepis ciliata* (P)
- Empodisma minus* (B, P)
- Juncus gregiflorus*
- J. novae-zelandiae* (B, K, P)
- J. planifolius*
- Luzula colensoi* (K)
- L. decipiens* (unc, K)
- L. migrata* (B, K, P)
- L. rufa* s.s. (K)
- L. subclavata*
- L. sp.* (unnamed; included in *L. picta*, as var. *limosa*, by Edgar 1966)
- L. colensoi* x *L. migrata* (unc)

**COMPOSITE HERBS**

- Anaphalis* spp. (see below)
- Brachyglottis lagopus* var. *lagopus* (B, K, P)
- Brachyscome radicata* (K)
- Calmisia glandulosa* ssp. *glandulosa* (incl. *C. g.* var. *latifolia*) (B, K, P)
- C. gracilentata* ssp. (B, K, P)
- C. incana* (incl. *C. discolor*) (B, K, P)
- C. spectabilis* ssp. *spectabilis* (B, K, P)
- C. sp.* (unnamed) (strongly rhizomatous (B, P))
- C. gracilentata* x *C. incana* (unc)
- C. g.* x *C. spectabilis* (unc)
- C. incana* x *C. spectabilis* (unc)
- Craspedia minor* var. *minor* (incl. *C. major*)
- C. viscosa* (B, K)
- C. sp. (i)* (unnamed) (summer-green bog plants; glomerules 0.8-1.5 cm diam.) (B, P)
- C. sp. (ii)* (unnamed) (summer-green plants of seepages; glomerules 1.8-2.5 cm diam.) (K)
- Gnaphalium audax*
- G. limosum* s.s.
- G. paludosum* s.s. (P)
- G. tuahinicum*

**MONOCOT HERBS (OTHER THAN ORCHIDS)**

**GRASSES, SEDGES, RUSHES**

- Astelia fragrans*
- A. sp.* (unnamed; aff. *A. nervosa*) (lvs not conspicuously white axially as in *A. nervosa*) (unc, P)
- Herpolition novae-zelandiae* (B)
- Luzuhaga parviflora* (unc, K)
- Phormium cookianum*
- Potamogeton suboblongus* (B)

(insert above)

*Halichtysum filicaule*

- Anaphalis keteaensis* s.s. (unc)
- A. sp. (i)* (*Halichtysum bellidioides* s.s.) (B, K, P)
- A. sp. (ii)* (unnamed) ("*Halichtysum alpinum*" of Cockayne 1928) (lvs and capitula larger than in *A. sp. i*) (B, K, P)

Lagenifera pinnatifida (P)

L. pumila

L. strangulata

Leucogenes (contopodium ss. (K)

Microseris scapigera var. (P)

Pseudognaphalium sp. (aff. P. luteo-  
album) (lvs mostly linear-spathulate  
to linear, acute)

Raoulia albosericea (B, K, P)

R. glabra (B, K, P)

R. grandiflora var. grandiflora (K)

R. tenuicaulis (incl. R. t. var. dimorpha  
and R. t. var. pusilla) Drosera arcturi (K, P)

R. albosericea x R. glabra (unc)

Senecio minimus

S. rufiglandulosus ssp. rufigland-  
ulosus (K)

S. sp. (unnamed; included in S. glauco-  
phyllus, as ssp. discoideus, by  
Ornduff 1960) ssp. (K)

Taraxacum magellanicum (K)

C. sp. (iii) ("Tussock Race" of Pritchard) (K)

Centella uniflora

Colobanthus acicularis (K)

C. apetalus (incl. C. a. var. alpinus) (K)

C. strictus (K)

D. spathulata (P)

Epilobium astonii (K)

E. atriplicifolium s.s. (B, K, P)

E. brunnescens s.s.

E. chlorifolium (K, P)

E. cockayneanum (B, K)

E. glabellum s.s. (K)

E. macropus (K)

E. melanocaulon (K)

E. nummularifolium

E. pedunculare agg.

E. pernitens (unc, K)

E. pubens

E. pycnostachyum (K, P)

E. rotundifolium

E. tenuipes s.s. (B, K)

E. sp. (unnamed; included in E.  
brunnescens, as ssp. minutiflorum,  
by Raven and Engelhorn, 1971) (K)

E. melanocaulon x E. pycno-  
stachyum (unc, K)

Euphrasia cuneata (B, K, P)

E. zelandica agg. (K, P)

DICOT HERBS OTHER THAN

COMPOSITES

Acaena anserinifolia

A. amittens (unc, K and P)

A. profundincisa (K, P)

Aciphylla colensoi var. colensoi (B, K, P)

Asp. (unnamed; aff. A. squarrosa) (K)

Anisotome aromatica (B, K, P)

Cardamine sp. (i) (C. debilis agg.)  
(cf. "Long Style" of Pritchard 1957)

C. sp. (ii) ("Sneeze Race" of Pritchard  
1957) (K)



*Forstera bidwillii* (incl. F.b. var. <sup>*dansifolia*</sup>) (K,P)  
*Galium parviflorum* ss. (K,P)  
*Gentiana bellidifolia* var. <sup>*bellidifolia*</sup> (B,K,P)  
*G. grisebachii* (B,K,P)  
*G. montana* (incl. *G. m.* var. *stolonifera*, *G. patula*, and *G. bellidifolia* var. *pulchella*) (K,P)  
*Geranium potentillodes* var. *pot.*  
*G. sessiliflorum* var. *novae-zelandiae*  
*G. sp.* (unnamed; aff. *G. microphyllum*) (B,K,P)  
*Geum sp.* (*G. parviflorum* agg.) (stock short; terminal leaflet about as long as broad; fls up to 13mm diam) (K)  
*Gonocarpus aggregatus* (B)  
*G. micranthus* ssp. *micranthus* (B,P)  
*Hydrocotyle heteromera* (unc)  
*H. microphylla*  
*H. moschata* (unc)  
*H. sp.* (unnamed; included in *H. n.z.*, as var. *montana*, by Kirk 1899) <sup>var.</sup> (B,K)  
*Hypericum japonicum* agg. (B)  
*Jovellana repens* (unc, K)  
*Kelleria dieffenbachii* (K)  
*K. laxa* (unc P)  
*Lilium monogynum* (K)  
*Montia fontana* s.s. (K)

*Myosotis forsteri*  
*M. sp. (i)* (*M. australis* agg.) (fls yellow) (K,P)  
*M. sp. (ii)* (unnamed; included in *M. pygmaea*, as var. *drucei*, by Moore 1961) (K,P)

*Neopaxia sp.* (*Montia calycina*) (K)  
*Nertera ciliata*  
*N. depressa* (unc)  
*N. sp.* (unnamed; aff. *N. dichondrifolia*) (hairs present on both surfaces of lf, straight, not hooked as in *N. dichondrifolia*)  
*Oreomyza colensoi* s.s. (incl. *O. c.* var. *hispida* and *O. c.* var. *multifida*) (K,P)  
*O. sp.* (unnamed; aff. *O. ramosa*)  
*Oursia caespitosa* ssp. *caespitosa* (unc, K)  
*O. colensoi* (K)  
*O. lactea* ssp. *drucei* (K)  
*O. macrophylla* ssp. *robusta* (B,K)  
*Oxalis magellanica*  
*Parakebe diffusa* ssp. *diffusa* (K,P)  
*P. olsenii* (K,P)  
*P. spathulata* (K)  
*Phyllachne colensoi* var. *colensoi* (K)  
*Plantago novae-zelandiae* (K)  
*P. raouli*  
*P. triandra* s.s. (P)  
*P. uniflora* (K)  
*Pratia angulata* agg.  
*Ranunculus cheesemanii* var. (*R. carsei*) (K)  
*R. insignis* (K)  
*R. nivicola* (K)  
*R. reflexus*  
*R. royi* (unc, K)  
*R. sp. (i)* (cf. *R. glabrifolius*) (B)  
*R. sp. (ii)* (unnamed; included in *R. lappaceus*, as var. *macrophyllus*, by Kirk 1899) (B,K,P)  
*Psychrophila novae-zelandiae* (K)

Hordeum murinum (unc, near hut)  
Poa pratensis

### RUSHES

Juncus effusus  
J. tenuis

### COMPOSITE HERBS

Achillea millefolium  
Cirsium arvense  
Crepis capillaris  
Hieracium pilosella  
Hypochoeris radicata  
Leontodon taraxacoides  
Mycelis muralis  
Taraxacum officinale s.l.

Scandia sp. (*S. rosifolia* agg.) (leaf-lets ovate-lanceolate) (K)

Schizilema colensoi (K, P)

*S. nitens* (K)

*S. sp.* (unnamed) (lvs cut  $\frac{1}{3}$ - $\frac{2}{3}$  to base; scattered hairs on upper surface; involueral bracts obtuse; fr = or  $<$  its pedicel) (K)

Scleranthus uniflorus (K)

Stachys minima (K)

Stellaria decipiens (incl. *S. parviflora*)

*S. gracilentia* (K, P)

Urtica incisa

Viola cunninghamii (B, K, P)

*V. filicaulis* (B, K, P)

Wahlenbergia albomarginata var. pygmaea (B, K, P)

### ADVENTIVE SPECIES

(excluding planted spp.)

### TREES AND SHRUBS

Pinus contorta

Lupinus arboreus

Ulex europaeus (unc)

### GRASSES

Agrostis capillaris

Aira caryophylla

Anthoxanthum odoratum

Dactylis glomerata

Festuca nigrescens

Holcus lanatus

### DICOT HERBS OTHER THAN

#### COMPOSITES

Anagallis arvensis  
Cerastium fontanum ssp. vulgare  
Euphrasia nemorosa  
Linum catharticum  
Lotus pedunculatus  
Plantago major (unc)  
Prunella vulgaris  
Rumex acetosella  
Trifolium repens  
Veronica arvensis

No. of indigenous spp.: 377.

No. of adventive spp.: 31.

DOUBTFUL AND ERRONEOUS RECORDS NOT INCLUDED IN MAIN LIST

<u>Record</u>	<u>Probable identity of plant on which record based</u>
<i>Aciphylla squarrosa</i> (Cunningham 1974)	<i>Aciphylla</i> sp. (unnamed)
<i>Astelia nervosa</i> (Cunningham 1974)	<i>Astelia fragrans</i>
<i>Carex diandra</i> (Cunningham 1974)	<i>Carex dipsacea</i>
<i>C. ternaria</i> (Cunningham 1974)	<i>C. cotiacea</i>
" <i>Celmisia graminifolia</i> X <i>C. spectabilis</i> " (Giv- <i>Chionochoa flavescens</i> [en 1984])	<i>Celmisia gracilentia</i> X <i>C. spectabilis</i> <i>Chionochoa pallens</i>
(Elder 1959, as <i>Danthonia flavescens</i> )	
<i>C. rigida</i> (Elder 1959, as <i>Danthonia</i> <i>rigida</i> )	<i>C. tubra</i>
<i>Coprosma parviflora</i> (Cunningham 1974)	<i>Coprosma</i> sp. (i) (unnamed)
<i>Cotiarca angustissima</i> (Cunningham 1974)	<i>Cotiarca plumosa</i>
<i>C. sarmentosa</i> (Cunningham 1974)	<i>C. arborea</i> X <i>C. pteridoides</i>
<i>Lepidothamnus intermedius</i> (Cunning- ham 1974)	<i>Lepidothamnus intermedius</i> X <i>L. laxifolius</i>
<i>Dracophyllum longifolium</i> var. (Cunningham 1974, as <i>D. filifolium</i> )	
<i>Gaultheria depressa</i> (Cunningham 1974)	<i>Gaultheria</i> sp. (unnamed)
<i>G. rupestris</i> (?) (Cunningham 1974)	
<i>Hebe diosmifolia</i> (Cheeseman 1925; <i>Hebe venustula</i> as <i>Veronica diosmifolia</i> )	
<i>Meliccytus crassifolius</i> (Cunning- ham 1974, as <i>Hymenanthera crass.</i> )	<i>Meliccytus alpinus</i>
<i>Kirkianella novae-zelandiae</i> (Cunn- ingham 1974)	
<i>Koeleria novozelandica</i> (Cunning- ham 1974, as <i>K. kurtzii</i> )	<i>Deyeuxia avenoides</i>
<i>Lagenifera petiolata</i> (Cunning- ham 1974)	<i>Lagenifera strangulata</i>
<i>Myosotis pygmaea</i> (Cunningham 1974)	<i>Myosotis</i> sp. (unnamed)

*Nertera dichondrifolia* (Cunning- *Nertera* sp. (unnamed)  
ham 1974)

*Olearia pachyphylla* (Carse 1929) *Olearia furfuracea*

*Oreomyza rigida* (Cunningham  
1974)

*Pimelea aridula* (Cunningham 1974) *Pimelea suteri* var.

*P. lyallii* (Elder 1959) *P. suteri* var.

*P. pseudolyallii* (Cunningham 1974, p. 39) *P. suteri* var.

*Plantago triantha* (Cunninghamii) *Plantago novae-zelandiae*  
1974)

"*Poa laevis*" (Elder 1959, p. 23) *Rytidosperma setifolium*

*Raoulia australis* (Cunningham  
1974) *Raoulia albosericea*

*Rytidosperma nigricans* (Cunning- *Rytidosperma gracile*  
ham 1974, as *Notodanthonia*  
*nigricans*)

#### REFERENCES

Carse, H. 1928: Botanical notes, new species and varieties. Trans.  
N.Z. Inst. 59: 315-6.

Cockayne, L. 1928: "The Vegetation of New Zealand". 2nd ed. Engelmann,  
Leipzig. 456 pp.

Cunningham, A. 1974: Headwaters of the Tutakahi catchment: study  
of an erosion problem in Hawke's Bay. N.Z. Forest Service Tech. Paper  
No. 62. N.Z. Forest Service, Wellington. 59 pp.

- 17) Jovan, V.K. 1964: A taxonomic revision of Leimisia subgenus Leimisiatae sectia Petiolatae (Compositae-Astereae). N.Z. J. Bot. 22: 139-58.
- 
- Edgar, E. 1966: Luzula in New Zealand. N.Z. J. Bot. 4: 159-84.
- 1970: Cyperaceae. Pp. 167-285 in "Flora of New Zealand," Government Printer, Wellington. Vol. 2. 354 pp.
- Elder, N.L. 1959: Vegetation of the Kaweka Range. Trans. Roy. Soc. N.Z. (Bot.) 87: 9-26.
- Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pernettya). Trans. R. Soc. N.Z. Bot. 1: 155-73.
- 27) Kirk, T. 1899: "The Students' Flora of New Zealand." Government Printer, Wellington. 408 pp.
- 37) Moore, L.B. 1961: Boraginaceae, Oursia, Hebe (except Whipcord spp.). Pp. 806-33, 86-70, 885-952 in "Flora of New Zealand" Government Printer, Wellington. Vol. 1. 1085 pp.
- 47) Oliver, W.R.B. 1935: The genus Coprosma. Bishop Mus. Bull. 132. 207 pp.
- 57) Pritchard, G.G. 1957: Experimental taxonomic studies on species of Cardamine Linn. in New Zealand. Trans. R. Soc. N.Z. 85: 75-89.
- Raven, P.H.; Engelhorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. J. Bot. 9: 345-50.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.
- 

- 37) Löve, A.; Connor, H.E. 1982: Relationships and taxonomy of New Zealand wheat grasses. N.Z. J. Bot. 20: 169-86.
- 27) Hamlin, B.G. 1959: A revision of the genus Urcinia (Cyperaceae-Caricoideae) in New Zealand. Dom. Mus. Bull. 19: 106 pp.
- 47) Nootboom, H.P. 1978: A taxonomic revision of the Malesian and Australian species of Urcinia (Cyperaceae). Blumea 24: 511-20.
- 57) Arnduff, R. 1960: An interpretation of the Senecio laetus complex in N.Z. Trans. R. Soc. N.Z. 80: 63-77.

