

(Supplement to list III)
Revised Oct. 1987,

INDIGENOUS VASCULAR PLANTS PRESENT IN THE TAPUAE-O-
UENUKU AREA, BETWEEN AWATERE AND CLARENCE VALLEYS,
INLAND KAIKOURA RA., BUT NOT IN THE BEN MORE-
CHALK RA. AREA†

Dec. 1987,
June 1988,
June 1991,
July 1992

(† for plants of this area see list 107)

DICOT TREES AND SHRUBS

Brachyglottis cassinioides
Carmichaelia menziesii
C. sp. (unnamed) (erect shrubs;
pods 1-1.5 cm long) (unc) [= *C. sp. (a)*]
Coprosma atropurpurea
C. cheesemanii s.s.
C. perpusilla ssp. *perpusilla*
C. sp. (unnamed; aff. *C. pseudo-*
cuneata) (low-growing shrubs
with small, thick, arcuate lvs)
(unc) [= *C. sp. (d)*]
Coriaria plumosa
Diacophyllum longifolium
(incl. *D. filifolium* and *D.*
olivieri)
D. uniflorum
Gaultheria crassa

Hebe carnosula
H. cheesemanii
H. epactidea var. *epactidea*
H. laastii var. (*H. ramosissima*)

H. pimeleoides s.s. (incl. *H. p. var. minor*) (unc)
H. rakaiensis (unc)
Helichrysum coralloides

Leucopogon suaveolens (incl.
Cyathodes colensoi)
Meliccytus sp. [= *M. sp. (b)*]
(unnamed; aff. *M. alpinus*) (low-
growing ^{rhizomatous} shrubs with recurved
spinous branches and narrowly
linear-oblongate lvs) (unc)
Muehlenbeckia astenii (Martin,
1932 and 1938) (Winterton R.,
2000 ft)
Pachystegia sp. ("*Pachystegia B*"
of Molloy and Simpson 1980) [= *P. sp. (c)*]
Pentachondra pumila
Pimelea concinna
P. sericeovillosa
Pitiosperum anomalum (unc)

DICOT LIANES

Clematis quadribracteolata (unc)
Convolvulus fractosaxosa

FERNS

Pellaea calidicrupium
Polystichum cystostegia

GRASSES

Agrostis muelleriana
A. muscosa (unc)

A. personata (unc)
A. sp. (unnamed; aff. *A. dyeri*)
(panicle spreading, not contracted as in *A. dyeri*) [= *A. sp. (b)*]

Isolepis Aucklandica
I. caligenis
I. habra
Oiceobolus pectinatus

Deschampsia novae-zelandiae **RUSHES**

Elymus enysii
E. tenuis

Luzula pumila
L. travelsii s.s.

L. sp. (i) (unnamed; included in *L. ennita*, as var. *petrieana*, by Edgar 1966) [= *L. sp. (b)*]

Microchloa novae-zelandiae

Keeleria cheesemani

L. sp. (ii) (unnamed; included in *L. rufa*, as var. *albicomans*, by Edgar 1966) (unc) [= *L. sp. (c)*]

Poa buchananii

P. dipsacea

P. lindsayi

P. manicata (unc)

P. subvestita

P. sp. (unnamed; aff. *P. novae-zelandiae*) (lvs narrower than in *P. n.z.*) [= *P. sp. (c)*]

Rytidosperma australe (unc)

R. merum (unc)

R. pumilum

Trisetum sp. (*T. spicatum* agg.)

COMPOSITE HERBS

Brachyglottis bellidioides ss.

Brachyscome radicata

Calmisia bellidioides

C. sessiliflora

C. sinclairii (incl. *C. allanii* and *C. durietzii*)

Dolichoglossis lyallii

Ewartia sinclairii

Gnaphalium laterale [formerly *G. sp. (c)*]

G. mackayii ss.

SEDGES

Carex acicularis (unc)

C. berggrenii (unc)

C. comans (unc)

C. coriacea

C. enysii (unc)

C. hectorii (unc)

C. inversa (unc)

C. maerica

C. muelleri (unc)

C. sp. (unnamed; aff. *C. wakatipu*) (robust plants; lvs up to 6mm wide; utricles up to 4mm long) [= *C. sp. (i)*]

Haastia pulvinaris ss.

H. recurva

H. sinclairii ss.

H. sp. (unnamed; included in H. pulvinaris, as var. minor, by Haung 1912) [= H. sp. (b)]

Lagenifera barkeri (unc)

Leptinella deudyi

L. pectinata s.s. (unc)

Leucogenes grandiceps (unc)

Pseudognaphalium sp. (unnamed; aff. P. luteoalbum) [P. sp. (b)]

Raoulia apicinigra s.s.

R. bryoides

R. cinerea (unc)

R. grandiflora var. grandiflora

R. sp. (unnamed) (stems forming open patches; lvs rigid, recurved, subacute; upper surface grey-green, lower white) [R. sp. (c)]

Chionohebe pulvinatis

Colobanthus buchananii

C. canaliculatus (unc)

Crassula multicaulis (unc)

Epilobium angustum (unc)

E. cocharyneanum (unc)

E. crassum

E. forbesii

E. ktulleanum

E. porphyrium

E. pycnostachyum

E. tasmanicum

E. tenuipes s.s.

Euphrasia zelandica var. (fls small, 2-3 mm wide)

Gentiana bellidifolia s.s.

G. tenuifolia (unc)

Gingidia decipiens

Gunnera densiflora (unc)

Hydrocotyle sulcata

Ischnocarpus novae-zelandiae (B.C. Aston MS)

Kalleria dieffenbachii

Lobelia roughii

Mazus radicans

Montia calycina var. calycina (incl. Claytonia australasica var. biflora)

Myosotis macrantha (unc)

M. traversii var. traversii

M sp. (i) (unnamed; included in M. pygmaea, as var. drucei, by Moore 1961) [= M sp. (e)]

DICOT HERBS

Aciphylla menziesii

Anisoteme pilifera (unc)

A: sp. (unnamed; included in A. umbicata, as var. prostrata, by Dawson 1961) [= A. sp. (b)]

Galium sp. (unnamed; aff. G. perpusillum) (lvs wider than stipules, elliptic-lanceolate; apex rounded to subacute) (unc) [G. sp. (b)]

Cardamine bilobata

C. sp. ("Scree Race" of Hitchcock 1957) [C. sp. (i)]

Cheesemanianthus enysii (Martin 1932, as Nasturtium enysii) (Mt. Glacstone)

Chenopodium detestans (Martin 1938) (Tapuaenuku)

M.sp.(ii)(unnamed)(plants forming cushions c. 5cm diam.; lvs spatulate, up to 1cm long, sub-acute, upper surface and margins and margins with long, spreading hairs, lower surface with few hairs or glabrous; fls solitary, 4-5mm diam.; anthers wholly below corolla scales) (unc) [= M.sp.(f)]

Neopaxia sp. (incl. Montia calycina and Claytonia australasica var. biflora) [= N.sp.(b)]

Nertera ciliata (unc)

Notothlaspi australe s.s. (incl. N. rosulatum) N.sp. (unnamed) (N. australe auct., non Thlaspi australe Hook.f.) [= N.sp.(b)]

Plantago lanigera

P. obconica (unc)

P. triandra ss.

Psychrophila obtusa (unc)

Ranunculus haastii ssp. haastii (unc)

Rumex flexuosus (unc)

Stellaria roughii

Swainsona novae-zelandiae (St Bernara Spur)

Wahlenbergia cartilaginea (unc)

No. of spp. in Tapuaenuku area: 447.

No. of spp. in Bai Mote-Chalk Ra.

area : 566

No. of spp. common to the two

areas : 308

No. of spp. confined to Tapuae-

nuku area : 139

No. of spp. confined to Bai Mote-

Chalk Ra. area : 258

No. of species: 138.