

PLANTS MENTIONED BY COMMON NAMES IN THIS PAPER

acaena	<i>Acena anserinifolia</i>
ake-ake (Craig)	<i>Olearia paniculata</i>
akiaki (Greville)	<i>Olearia paniculata</i> ?
black maire	<i>Nestegis (Olea) cunninghamii</i>
bracken fern	<i>Pteridium aquilinum</i> var. <i>esculentum</i>
catsear	<i>Hypochaeris radicata</i>
cassinia	<i>Cassinia leptophylla</i>
clematis	<i>Clematis</i> spp.
cocksfoot	<i>Dactylis glomerata</i>
danthonia	<i>Notodanthonia</i> spp.
five-finger	<i>Pseudopanax arboreum</i>
fog	<i>Holcus lanatus</i>
flax	<i>Phormium tenax</i>
fuchsia	<i>Fuchsia excorticata</i>
hangehange	<i>Geniostoma ligustrifolium</i>
hard fern	<i>Paesia scaberula</i>
heketara	<i>Olearia rami</i>
hinau	<i>Elaeocarpus hookerianus</i>
horehound	<i>Marrubium vulgare</i>
honeysuckle (Craig)	<i>Knightsia excelsa</i>
horopito	<i>Pseudowintera axillaris</i>
Italian thistle	<i>Carduus pycnocephalus</i>
kahikatea	<i>Podocarpus dacrydioides</i>
kaikomako	<i>Pennantia corymbosa</i>
kamahi	<i>Waimanania racemosa</i>
kanuka	<i>Leptospermum ericoides</i>
karaka	<i>Corynocarpus laevigatus</i>
kawakawa	<i>Macropiper excelsum</i>
knio (Craig)	<i>Myoporum laetum</i> ?
kohekohe	<i>Dysoxylum spectabile</i>
kohukohu	<i>Pitiosporum tenuifolium</i>
konini (Craig)	<i>Fuchsia excorticata</i>
lancewood	<i>Pseudopanax crassifolium</i>
lemon tree (Craig)	<i>Pitiosporum eugenioides</i> ?
mahoe	<i>Meliccytus ramiflorus</i>
mako (Craig)	<i>Aristolelia serrata</i> ?
manuka	<i>Leptospermum scoparium</i>
mapou	<i>Myrsine australis</i>
matai	<i>Podocarpus spicatus</i>
matipo (Greville and Craig)	<i>Pitiosporum</i> or <i>Myrsine</i> ?
miro	<i>Podocarpus ferrugineus</i>

1967]

ESLER - KAPITI ISLAND

393

ngaio	<i>Myoporum laetum</i>
nikau	<i>Rhopalosiphis sapida</i>
northern rata	<i>Metrosideros robusta</i>
ongaonga	<i>Urtica ferox</i>
pate	<i>Schefflera digitata</i>
pigeonwood (pigeontree, Craig)	<i>Hedycarya arborea</i>
plumegrass	<i>Dichelachne crinita</i>
prairie grass	<i>Bromus catharticus (uniloides)</i>
pukatea	<i>Laurelia novae-zelandiae</i>
putaputaweta	<i>Carpodetus serratus</i>
rangiora	<i>Brachyglottis repanda</i>
raukawa	<i>Pseudopanax edgerleyi</i>
raurekau	<i>Coprosma australis</i>
rewarewa	<i>Knightsia excelsa</i>
rice grass	<i>Microlaena stipoides</i>
rimu	<i>Dacrydium cupressinum</i>
ryegrass	<i>Lolium perenne</i>
shining broadleaf	<i>Griselinia lucida</i>
supplejack	<i>Rhipogonum scandens</i>
sweet vernal	<i>Anthoxanthum odoratum</i>
tall oatgrass	<i>Arrhenatherum elatius</i>
tarata	<i>Pitiosporum eugenioides</i>
tauhinu (Greville)	<i>Cassinia leptophylla</i>
taupata	<i>Coprosma repens</i>
tawa	<i>Betischmidia tawa</i>
tioki	<i>Alectryon excelsus</i>
toru	<i>Myrsine salicina</i>
totara	<i>Podocarpus totara</i>
wharangi	<i>Melicope ternata</i>
white clover	<i>Trifolium repens</i>
white maire	<i>Nestegis (Olea) lanceolata</i>
wineberry	<i>Aristolelia serrata</i>
variegated thistle	<i>Silybum marianum</i>

Note—*Poa caespitosa* referred to in this paper is the species now bearing the name *Poa laevis* (see V. D. Zotov, 1965: *Rec. Dom. Mus.* 5: 106.