



Oporua Scenic Reserve

Area: 1.6919 ha

OPOURA SCENIC RESERVE FOR: [] scenery [] bush ✓ scrub / tussock etc [] wetland [] plants ✓ animals [] geology / soil [] history / archaeology [] water supply LOCATION Situated off Kaitutara Road, 20 km SE of Featherston, 4.5 km S of the Lake Wairarapa shoreline. ACCESS By a 3m wide "right-of-way" from Kaitutara Road, which is unfenced and crosses 200m of pasture, gate in northern-most corner of reserve. OUTLINE & TOPO The recently fenced reserve is a square block of flat reclaimed swamp near the lake with pole stand regenerating Kahikatea. Exposure to strong southerly winds has resulted in wind-scooping and shaping of the canopy.										GRID REF S27 268600 AREA ha 0.88 1.6919 L.D. No. lot 1 DP 49734 Block XII, Wairarapa SD STATUS Scenic RES sub-Catag CONTROL CCL GAZETTED 1981 RESERVED HISTORY GIFTED to Crown by Mr. Reginald Matthews for preservation in 1976																																																																					
SURROUNDINGS It is completely surrounded by pasture, with scattered cabbage trees and stock-damaged kahikatea stands to the north. The extensive Te Horai Basin Drainage Scheme includes this reserved land.										AIR PHOTO 5497 M/10 date 16.10.79. nearby town 17 km Martinstown nearest reserve 1.5 km Holmes SR public road Kaitutara Road sea (km) ALTITUDE m NN 3 - 4																																																																					
SPM Gravel deposit brought down by Ruamahanga River, overlain by wind-blown loess.										ASPECT (Plat)																																																																					
SOIL Central recent soils from alluvium: Karimea and Manawatu.										RAIN mm yr 964 (approx)																																																																					
DRAINAGE Developed, often once poorly.										NAT FERTILITY High																																																																					
RES NOT DIS ✓✓ EN ✓✓										MAIN HABITATS and FEATURES																																																																					
<p>1. Predominantly immature kahikatea with some titoki, kumuka, kowhai, and cabbage tree over-shrubbying shrubs understorey. Canopy height 6-8m.</p> <p>2. Bordering fringe of exotic grasses and weeds - some phormium and cabbage trees.</p>										<table border="1"> <thead> <tr> <th>% area</th> <th>area ha</th> <th>native 1^o</th> <th>native 2^o</th> <th>ind</th> <th>ex</th> <th>regen</th> <th>NAT</th> <th>trend</th> <th>ecistics</th> </tr> </thead> <tbody> <tr> <td>85</td> <td>1.4</td> <td>✓</td> <td></td> <td></td> <td>M</td> <td>B</td> <td>I</td> <td>L</td> <td></td> </tr> <tr> <td>15</td> <td>0.3</td> <td>✓</td> <td>✓</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>H</td> </tr> <tr> <td colspan="10"><hr/></td> </tr> <tr> <td colspan="10"></td> <td>100</td> <td>1.7</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>										% area	area ha	native 1 ^o	native 2 ^o	ind	ex	regen	NAT	trend	ecistics	85	1.4	✓			M	B	I	L		15	0.3	✓	✓						H	<hr/>																				100	1.7								
% area	area ha	native 1 ^o	native 2 ^o	ind	ex	regen	NAT	trend	ecistics																																																																						
85	1.4	✓			M	B	I	L																																																																							
15	0.3	✓	✓						H																																																																						
<hr/>																																																																															
										100	1.7																																																																				
CONDITION The canopy is in good condition and the understorey is recovering very well since fencing. It has been heavily logged and stock-damaged in the past but there is little human impact now other than drainage. Wind damage is evident on the exposed south and west sides although not as severe as in neighbouring stock-damaged stands. A number of native trees have been planted.																																																																															
FENCING The reserve was fenced on all sides in 1972.																																																																															
PRIORITY Low at present but standard must be maintained.																																																																															
RATING 0-10 SCIENTIFIC 4... Pole stand of kahikatea of very small area, exposed to prevailing winds. Drainage likely to modify composition. Very low native bird numbers probably due to presence of magpies. SCENIC 2... Low canopy height; distant from road, on farmland. RECREATION 0... Awkward access across roadside ditch and rough farmland.																																																																															
CONCLUSIONS and RECOMMENDATIONS Although the reserve is very small, it is the only protected remnant of what was once a continuous belt of regenerating Kahikatea forest on the north banks of the Ruamahanga River. Other remaining patches are rapidly deteriorating as a result of stock damage and wind scouring. The regeneration of the reserved bush to its original local composition is now in doubt as the land is incorporated in the Te Horai Basin Drainage Scheme. If the reserve were to dry out further, some species may die out, possibly the Kahikatea itself. With this in mind it becomes more urgent to fence off the nearby groups of Kahikatea for preservation to encourage a more extensive bird habitat and seed source for further regeneration.										(Continued..)																																																																					
.... hours ... fine conditions										IG 1981																																																																					
NATURNESS										: TREND: D deteriorating; S steady; I improving																																																																					

CONCLUSIONS AND RECOMMENDATIONS (Continued)

There are three other reserves within 7km of Oporua Scenic Reserve: E.C. Holmes Memorial Scenic Reserves is a group of exotics and 'natives' amongst original kahikatea and tikoki; Kahutara Scenic Reserve is a deteriorated, species-depleted stand of titoki and kowhai on drier, drained ground, and Tuhitarata Scenic Reserve is outstanding as the last remnant of true kahikatea/pukatea swamp forest in the lowlands east of Lake Wairarapa. In this intensive pastoral landscape small stands of bush like these and Oporua Scenic Reserve become significant records of original vegetation types, and what effect drainage for pastoral use has had on the forest composition. Although the Oporua stand of young kahikatea has been reserved, it is saddening to watch similar remnants deteriorating in neighbouring paddocks. It may be too late to negotiate fencing these stands as suggested in Recommendations, but the possibility should be discussed immediately by the appropriate expertise.

The reserve is really only of botanical interest. There are weeds throughout, ragwort and Jerusalem cherry are the only noxious weeds present and do not seem to be posing a threat to the undergrowth. The forest floor is well lit so the undergrowth is prolific, in particular sedges, buttercup, grasses, *Parsonsia* and ferns. Divaricating shrubs and juveniles are concentrated at the bush margins. The canopy contains elements of semi-swamp forest (*kahikatea* and cabbage tree) and drier *titoki*, *kanuka* and *kowhai*, a mixture typical of Southern Wairarapa basin forests.

REFERENCES

Tall conifer: kahikatea, kauri, matai, miro, rimu, totara

2 remaining 11 tree-sized spp

3 broadleaved

* e.g. flax

⁵ incl. rough grassland, pasture etc.

NATIVE SPECIES

Trees and shrubs

Alectryon excelsus
Carmichaelia arborea
Coprosma areolata
C. propinqua
C. rigida
C. robusta
C. propinqua X C. robusta

Cordyline australis
Hoheria angustifolia
H. populnea var. lanceolatum
(incl sexstylosa)
Leptospermum ericoides
Melicope simplex
Melicytus micranthus

Melicytus ramiflorus
Myrsine divaricata
Pennantia corymbosa
Pittosporum tenuifolium
Plagianthus betulinus
Podocarpus dacrydioides
Sophora microphylla

Climbers, Lianes etc.

Muehlenbeckia australis
M. complexa

Parsonsia heterophylla
Passiflora tetrandra

Ripogonum scandens
Rubus schmidelioides

Grasses, rushes and like plants

Carex geminata
C. lambertiana
C. lessoniana

Carex virgata
C. sp.
Cyperus ustulatus

Microlaena stipoides
Uncinia uncinata

Herbs

Galium sp.
Haloragis erecta

Hydrocotyle americana
H. novae-zelandiae

Oxalis lactea
Polygonum sp.

Ferns

Asplenium bulbiferum
Blechnum fluviatile

Paesia scaberula
Polystichum vestitum

Pteris tremula

buttercup
 Californian thistle
Cyperus vegetus

hawthorn
 Jerusalem cherry
 plantain

ragwort
 tutsan
 grasses

fantail
 greywarbler

BIRDS
 magpie

silvereye